



## **Pulicaria uniseriata nom. nov. (Compositae, Inuleae)**

Author: Kilian, Norbert

Source: Willdenowia, 33(1) : 106

Published By: Botanic Garden and Botanical Museum Berlin (BGBM)

URL: <https://doi.org/10.3372/wi.33.33109>

---

BioOne Complete ([complete.BioOne.org](https://complete.BioOne.org)) is a full-text database of 200 subscribed and open-access titles in the biological, ecological, and environmental sciences published by nonprofit societies, associations, museums, institutions, and presses.

Your use of this PDF, the BioOne Complete website, and all posted and associated content indicates your acceptance of BioOne's Terms of Use, available at [www.bioone.org/terms-of-use](https://www.bioone.org/terms-of-use).

Usage of BioOne Complete content is strictly limited to personal, educational, and non - commercial use. Commercial inquiries or rights and permissions requests should be directed to the individual publisher as copyright holder.

---

BioOne sees sustainable scholarly publishing as an inherently collaborative enterprise connecting authors, nonprofit publishers, academic institutions, research libraries, and research funders in the common goal of maximizing access to critical research.

NORBERT KILIAN

## *Pulicaria uniseriata* nom. nov. (*Compositae*, *Inuleae*)

### Abstract

Kilian, N.: *Pulicaria uniseriata* nom. nov. (*Compositae*, *Inuleae*). – Willdenowia 33: 106. 2003. – ISSN 0511-9618; © 2003 BGBM Berlin-Dahlem.

The nomen novum *Pulicaria uniseriata* is validated to provide a legitimate binomial in *Pulicaria* for the N Somalian species originally described as *Sclerostephane longifolia*.

Comparative micromorphological and anatomical studies of achene and pappus (Kilian 1999) established a sister group relationship between the N Somalian genus *Sclerostephane*, a segregate of *Pulicaria* with a peculiar, strongly derived achene type, and a group of SW Asian true *Pulicaria* species. Consequently, *Sclerostephane* has been included in *Pulicaria* as a section that includes the five former *Sclerostephane* species along with the *P. glaucescens* group, five closely related species from S Arabia, SE Iran and S Pakistan (Kilian 1999: 174).

When making the required transfers to *Pulicaria*, I overlooked that the combination based on *Sclerostephane longifolia* Wagenitz & E. Gamal-Eldin is a younger homonym of *P. longifolia* Boiss. (1856), a synonym of *P. inuloides* (Poir.) DC. (see Gamal-Eldin 1981: 96). Henk Beentje (Kew) kindly pointed this out to me and gave me the opportunity to validate a nomen novum:

***Pulicaria uniseriata*** N. Kilian, **nom. nov.** ≡ *Sclerostephane longifolia* Wagenitz & E. Gamal-Eldin in Bot. Jahrb. Syst. 104: 106. 1983 ≡ *Pulicaria longifolia* (Wagenitz & E. Gamal-Eldin) N. Kilian in Willdenowia 29: 176. 1999, nom. illeg. [non Boiss., Diagn. Pl. Orient., ser. 2, 3: 16. 1856]. – The epithet refers to the autapomorphic lack of the outer, corona-like pappus in this species.

### References

- Boissier, E.: 1856: Diagnoses plantarum orientalium novarum, ser. 2, **3**. – Genève, etc.  
 Gamal-Eldin, E. 1981: Revision der Gattung *Pulicaria* (*Compositae* - *Inuleae*) für Afrika, Makaronesien und Arabien. – Phanerog. Monogr. **14**.  
 Kilian, N. 1999: Studies in the *Compositae* of the Arabian Peninsula and Socotra – 1. *Pulicaria gamal-eldinae* sp. nova (*Inuleae*) bridges the gap between *Pulicaria* and former *Sclerostephane* (now *P. sect. Sclerostephane*). – [Willdenowia 29: 167-185.](https://doi.org/10.1007/s11597-003-0001-1)

Address of the author:

N. Kilian, Botanischer Garten und Botanisches Museum Berlin-Dahlem, Freie Universität Berlin, Königin-Luise-Str. 6-8, 14191 Berlin, Germany; e-mail: n.kilian@bgbm.org