

# **On flower anatomy and embryology of Lophiocarpus polystachyus (Lophiocarpaceae)**

Author: Hakki, Madjit Ismail

Source: Willdenowia, 43(1) : 185-194

Published By: Botanic Garden and Botanical Museum Berlin (BGBM)

URL: https://doi.org/10.3372/wi.43.43120

BioOne Complete (complete.BioOne.org) is a full-text database of 200 subscribed and open-access titles in the biological, ecological, and environmental sciences published by nonprofit societies, associations, museums, institutions, and presses.

Your use of this PDF, the BioOne Complete website, and all posted and associated content indicates your acceptance of BioOne's Terms of Use, available at www.bioone.org/terms-of-use.

Usage of BioOne Complete content is strictly limited to personal, educational, and non - commercial use. Commercial inquiries or rights and permissions requests should be directed to the individual publisher as copyright holder.

BioOne sees sustainable scholarly publishing as an inherently collaborative enterprise connecting authors, nonprofit publishers, academic institutions, research libraries, and research funders in the common goal of maximizing access to critical research.

To the memories of Jules Parisinos BSc., Dept. of Agriculture, Nicosia, Cyprus, and Fritz and Gertrud Pommerencke, teachers of the German Language, Goethe Institute, Nicosia, Cyprus

## MADJIT ISMAIL HAKKI<sup>1</sup>

# **On flower anatomy and embryology of** *Lophiocarpus polystachyus* **(***Lophiocarpaceae***)**

#### **Abstract**

Hakki M. I.: On flower anatomy and embryology of *Lophiocarpus polystachyus* (*Lophiocarpaceae*). – Willdenowia 43: 185–194. June 2013. – Online ISSN 1868-6397; © 2013 BGBM Berlin-Dahlem. Stable URL: http://dx.doi.org/10.3372/wi.43.43120

Flower anatomy and embryology of *Lophiocarpus polystachyus* was studied, based on over one thousand slides left by Theo Eckardt and prepared from material of a wild source in South Africa chiefly in the 1970s. The all bisexual flowers of the polytelic inflorescences possess a well-developed nectary, are promoted on the abaxial side, have a perianth of five tepals, a 4-merous or, more rarely, by fusion of two adaxial stamens, 3-merous androecium, a somewhat stipitate, unilocular gynoecium of two fused carpels with one basally inserted ovule. Wall formation of the introrse and tetrasporangiate anther conforms to the Monocotyledonous type. Meiosis of the microsporocytes during which 9 bivalents are visible leads after simultaneous delimitation of the microspores to tetrahedral and decussate pollen tetrads. The tricolpate pollen is 3-celled when shed. The ovule is campylotropous, bitegmic and crassinucellate, as is common in the *Centrospermae*. Embryo sac development is of the monosporic, *Polygonum* type. Embryogenesis follows the Caryophyllad type of Johansen. The fruit is a 1-seeded drupe. The flower anatomy of *Lophiocarpus* confirms an affinity to the *Phytolaccaceae* s.l. and its distance from the *Chenopodiaceae* because of the missing gynophore in the latter. A comparison with the embryological data for *Corbichonia* does not corroborate an intimate relationship of the two genera.

Additional key words: *Caryophyllales*, *Phytolaccaceae*, *Corbichonia*, systematics

## **Introduction**

The small southern African genus *Lophiocarpus* (comprising four species), originally placed in the *Chenopodiaceae*, was transferred by Brown (1909) into the *Phytolaccaceae*, and there even merged with the Caribbean-South American genus *Microtea*. Its placement in the *Phytolaccaceae*, but again as a separate genus, was maintained by later authors such as Heimerl (1934), Eckardt (1964), Nowicke (1968), Takhtajan (1969), Behnke (1974) and Rohwer (1993), mostly in a separate subfamily *Rivinoideae* or *Microteoideae*, while authors splitting up *Phytolaccaceae*, such as Hutchinson (1959, 1973) and Brown & Varadarajan (1985), placed it with *Microtea* in the segregate family *Petiviaceae.* In all classifications, *Lophiocarpus*, however, is considered as an anomalous genus and usually included therefore with reservations only. Bortenschlager (1973), based on his palynological investigations, was the first who separated *Lophiocarpus* in a family of its own, but a valid publication of the family name *Lophiocarpaceae* was only provided by Doweld & Reveal (2008). Molecular phylogenetic analyses

<sup>1</sup> Dr Madjit Ismail Hakki, c/o Botanischer Garten und Botanisches Museum Berlin-Dahlem, Freie Universität Berlin, Königin-Luise-Str. 6–8, 14195 Berlin, Germany; e-mail: mahakki@hakki.de

by Cuénoud & al. (2002), Brockington & al. (2009) and Schäferhoff & al. (2009) unambiguously confirm *Phytolaccaceae* to be polyphyletic and support the separation of *Lophiocarpus* from both *Phytolaccaceae* and *Petiviaceae. Lophiocarpaceae* are recognized as a separate family also in the latest update of the Angiosperm Phylogeny Group classification (APG III 2009). Schäferhoff & al. (2009), moreover, show that *Lophiocarpus* and the Caribbean-South American *Microtea*, are by no means closely related; while *Microtea* constitutes a separate family in an isolated position of the caryophyllid clade, *Lophiocarpus* is sister to the *Aizoaceae-Nyctaginaceae-Phytolaccaceae* clade. A relationship of *Lophiocarpus* with *Corbichonia* (the latter formerly misplaced in *Molluginaceae*) as revealed in the molecular analyses is also supported by both genera sharing betacyanins (Cuénoud & al. 2002). Both genera are accepted to constitute the family *Lophiocarpaceae* by Stevens (2001+).

Theo Eckardt, when he was working at the end of the 1950s on the familial classification of the *Centrospermae* for the 12th edition of Engler's "Syllabus" (Eckardt 1964), developed an interest in the then still little known *Phytolaccaceae*, especially in *Phytolacca*, in those days the so-called "genus primordioides" of the *Centrospermae* (Bittrich 1993), and subsequently also in *Lophiocarpus*. When Eckardt in 1977 unexpectedly passed away, he had been studying *Lophiocarpus* for some years (Eckardt 1974), without having, however, investigated its embryology, but he left behind more than one thousand anatomical slides of *Lophiocarpus polystachyus.* Since an embryological study of *Lophiocarpus*, when compared to the available data on other *Centrospermae* taxa (Rocen 1927; Mauritzon 1934; Johanson 1950; Kajale 1954; Narayana & Lodha 1963; Davis 1966; Hakki 1972; Johri & al. 1992) might help to clarify the phylogeny and systematic position of this odd genus, I worked up this material during the last years. The results of these flower anatomical and embryological studies are presented here.

# **Material and methods**

This study is based on more than one thousand slides of *Lophiocarpus polystachyus* left by Theo Eckardt that had been sectioned by Maria Gerstenberger, Monika Lüchow (née Schröder, Berlin) as well as by various students of him during the years 1969 to 1977. Much of the material had been collected in South Africa near Vioolsdrif (28°54'S, 17°44'E) and fixed in FPA (formaldehyde, proprionic acid and ethyl alcohol) and CRAF V (chromic acid, ethyl alcohol and formaldehyde) by Hans-Dieter Ihlenfeldt and Heidrun Hartmann (Institut für Allgemeine Botanik und Botanischer Garten Hamburg) in 1969 and 1971, respectively (for details see Eckardt 1974: 16). Some plants were grown from seed in the Botanic Garden Berlin-Dahlem and their inflorescences were fixed in CRAF V and serial sections were made by Monika Lüchow. Prior to embedding in Malinol the serial sections of 6 to 10 µm thickness were stained either with Heidenhain's haematoxiylin and counterstained with Fast Green FCF or, for the sake of rendering the vascular system more clearly, with Fuchsin and Fast Green.

For an embryological study of *Lophiocarpus*, haematoxylin is definitely better but older pollen is to such an extent overstained that it is not possible to detect the nuclei contained therein. For clear pictures of the contents of the pollen grains as well as for critical anatomical work Fuchsin has proved to be more suitable. Both stains have kept perfectly well. This is perhaps explained by the fact that the slides were constantly kept in darkness, being no more viewed since 1977. I used a Leitz-Dialux microscope for the study of the sections and in addition I also studied the herbarium collections at B. All figures have been drawn by the present author.

# **Results**

## **Flower morphology**

All flowers of the polytelic inflorescences are bisexual. They possess a well-developed nectary and are promoted on the abaxial side. Five tepals make up the perianth, with the adaxial member being a little wider than the



Fig. 1a–1j. Transverse sections of a flower with five tepals and a 4-merous androecium, showing the vascular bundles.



Fig. 2a–i. Transverse sections of a flower with five tepals and a 4-merous androecium, showing the vascular bundles; 3a–f: Transverse sections of a triplet of flowers, the middle flower has a 3-merous androecium due to fusion of the two adaxial stamens.

others, see Fig. 1–4. The androecium is either 4-merous (Fig. 1–3) or to a lesser extent 3-merous (Fig. 4). The latter condition is due to fusion of the two adaxial stamens. Also the somewhat stipitate and unilocular gynoecium with one basally inserted ovule is oligomerous, showing two intimately fused carpels, the hairy sutures of which form two narrow bands of inner stigmatic tissue, leading down to the foot of the funicle and acting as obturator (Fig. 39). Only the abaxial carpel produces the ovule, the adaxial one being sterile, see Fig. 1–4.

#### **The nectarium**

The nectarium is situated on the filaments of the stamens, right at the foot of the androecium (Fig. 5), building a conspicuous ring of densely staining tissue around the base of the somewhat stipitate ovary. Viewed in the early stages of flower development, it originates from a small group of cells of the filaments. In longitudinal sections of the earliest stage about 8 such cells destined to build the nectarium are discernible (Fig. 6) at the base of each filament. At the time when the ovule has become hemitropous (Fig. 5, 5a) about 18 such cells can be seen. When

Downloaded From: https://bioone.org/journals/Willdenowia on 27 Jul 2024 Terms of Use: https://bioone.org/terms-of-use

the ovule is completely bent just after RT II with the fertile macrospore (Fig. 25, 25a) the number of the future secretory cells is almost four times that of the earliest stage, showing many cells with 2 smaller and other cells with single but much larger nuclei. This perhaps indicates nuclear fusion and polyploidization. The already enlarged nectarium tightly embraces the base of the ovary showing no gap between the two. The flower is still a closed bud and I noticed no sign of nectar secretion. During double fertilization (Fig. 36) the flower is open, the nuclei of the nectarium cells have become still bigger and the cytoplasm of the cells is granular and very dense. There is a small gap between the nectarium and ovary wall. When the embryo sac shows a 2- or 3-celled embryo and 4 endosperm nuclei (Fig. 41, 41a), the nectarium has reached the peak of its development consisting of about 100 cells and the gap between nectarium and ovary is large. After that, the number of secretory cells does not increase; on the contrary the nectarium declines gradually, showing signs of degeneration due to lysis and absorption of its cells. Finally, at the 9-celled embryo stage, it is almost completely lost, showing vacuolated cells with very little cytoplasm.



Fig. 4a–j. Transverse sections of a flower with five tepals and a 3-merous androecium. Study of the vasculature; 5: Longitudinal section of a young flower, showing the nectariferous tissue (Fig. 5a) on the base of the anther filaments; 6: Longitudinal section of the nectariferous tissue, the youngest stage observed.



Fig. 7–20. Development of the anther wall, microsporogenesis and microgametogenesis. – 7–9: Sections of parts of three microsporangia showing the development of the wall layers and microspore mother cells (stippled). – 10: Transverse section of one half of an anther with the prominent sporogenous tissue in the middle of each of the two sporangia just after all 4 surrounding wall layers have been completed. – 11: One binucleate tapetum cell and a microsporocyte in prophase of the reduction division. – 12: Microsporocyte in meiosis I. – 13: Microsporocyte in meiosis I showing 9 chromosomes. – 14: One binucleate tapetum cell and a microsporocyte in meiosis II. – 15: Two microspore tetrads. – 16: A microspore on the left just after liberation from the tetrad and an older one with thicker wall on the right. – 17: Uninucleate microspore prior to the onset of the first mitosis in the pollen grain. – 18: Binucleate microspore with the lenticular generative cell lying on the cell wall. - 19: Binucleate microspore after migration of the generative into the vegetative cell. – 20: Trinucleate ripe pollen grain before liberation from the microsporangium.

#### **Anther wall formation, microsporogenesis and microgametogenesis**

The introrse anther is dithecous and tetrasporangiate (Fig. 4a). Its wall develops after the monocotyledonous type of Davis (1966), comprising 4 layers, i.e. epidermis, endothecium, one middle layer and a single-layered tapetum (Fig. 9). Immediately after their completion all wall layers show uninucleate cells. The tapetal layer is of dual origin and although the nuclei of its cells multiply to 4 or 5 during meiosis I of the pollen mother cells, yet these nuclei fuse later and mostly 2 large nuclei remain

Downloaded From: https://bioone.org/journals/Willdenowia on 27 Jul 2024 Terms of Use: https://bioone.org/terms-of-use

in each cell until maturity and thereafter (Fig. 11, 14). The mature tapetum is thus binucleate. After meiosis and simultaneous cytokinesis the microsporocytes (Fig. 14) develop into tetrahedral and decussate tetrads of microspores (Fig. 15) that are nourished by the glandular tapetum. Ubisch bodies (orbicules) of differing sizes are always produced. They are most abundant when the tapetum has largely declined and the uninucleate pollen grains are highly vacuolated, being in the so called "signet-ring" stage (Fig. 17). After the first mitosis, a large vegetative and a small lenticular generative cell are produced in the pollen grain. At first the small lenticular cell is placed on the inner wall of the pollen grain (Fig. 18);



Fig. 21–25a. Development of the ovule and megasporogenesis. Longitudinal sections. – 21: Very young open gynoecium with the orthotropous ovule protruding just after production of a parietal cell and initiation of the inner integument. – 22: The curving ovule after initiation of inner and outer integuments and first periclinal divisions in the nucellus epidermis. – 23: Part of nucellus showing fertile chalazal megaspore and degenerating micropylar megaspore and dyad cell. – 24: Whole nucellus with fertile megaspore and three degenerating micropylar spores. – 25: The closed gynoecium just after RT2 (same stage as Fig. 24) showing pollen-transmitting tissue and fully bent campylotropous ovule with inner integument forming the micropyle and a prominent air space between the two integuments, the stippled area below is the nectary. – 25a: Part of the nectary enlarged, note several cells showing 2 nuclei – sign of high mitotic activity!



Fig. 26–31. Megagametogenesis and development up to 4-nucleate embryo sac stage. Longitudinal sections of micropylar part of the nucellus (Fig. 26, 27, 29–31) and overview of whole ovule (Fig. 28). – 26: Uninucleate embryo sac. – 27: First mitosis leading to 2-nucleate stage. – 28: A not completely bent ovule carrying a 2-nucleate embryo sac, which is rather unusual. – 29: Two-nucleate embryo sac from a normally bent ovule. – 30: The normal-sized 4-nucleate embryo sac. – 31: A rather smaller 4-nucleate embryo sac.

later it is engulfed by the vegetative cell and comes to lie right in the middle of the pollen grain (Fig. 19). Already before anther dehiscence, at the stage of the young 8-nucleate embryo sac, the generative nucleus divides in the pollen grain, producing two vermiform nuclei (gametes). The mature pollen grains are 3-nucleate (Fig. 20), showing also three conspicuous colpi. The endothecium (fibrous layer) develops the usual thickenings and anther dehiscence is longicidal. The pollen grains germinate monosiphonously on the ripe stigma, during which the two vermiform gametes move into the developing pollen tube. I could not detect the tube nucleus.

#### **Megasporogenesis and gametophytogenesis**

In *Lophiocarpus*, like in all other members of the *Centrospermae*, ovules are orthotropous when initiated. They bend in the course of further development, becoming campylotropous. At maturity they are bitegmic and crassinucellate with the inner integument forming the micropyle and also showing a distinct air space between the two integuments at the chalazal side of the ovule (Fig. 21–25).

Fig. 22 shows the longitudinal section of a fairly young ovule a short time after both integuments have been initiated. The nucellus of the ovule already displays one to two macrosporocytes and one layer of parietal cells. The pollen mother cells in the anthers of the same flower are in the meiosis I stage of development. Fig. 23, 24 and 26–34 show the development of the monosporic *Polygonum* type embryo sac.

#### **Fertilization, endosperm and embryo development**

The ripe 3-celled pollen grain germinates monosiphonously on the stigma. Stained with haematoxylin, the two sperm cells are well visible, whereas I could not localize the tube nucleus in the pollen tube. The latter traverses at first the short style, reaching the inner surface at the top



Fig. 32–36. Development of 8-nucleate embryo sac up to early double fertilization. Longitudinal sections of micropylar part of nucellus. – 32: The not-yet organized 8-nucleate stage with a small deposit of starch granules. – 33–34: The semi-organized (Fig. 33) and completely organized (Fig. 34) 8-nucleate embryo sac. – 35: The 7-nucleate ready-to-fertilize embryo sac with ample starch granules around the fused polar nuclei. – 36: Embryo sac during double fertilization. Right above the burst pollen tube on the right are the 2 synergids, the nucleus of one of which is degenerating. The sperm cell on the left has not yet reached the larger egg nucleus and above these are the huge central nucleus and the second sperm cell in the process of fusion. The 3 antipodals are still present. Note the multitude of starch grains, some of which have been delivered by the pollen tube.



Fig. 37–40. Late stages of double fertilization, young zygote and early endosperm development, pollen tube transmitting tissue and trichomes. Longitudinal sections of parts of nucelli (Fig. 37, 38). – 37: Late double fertilization, fusion of male gamete with central nucleus is already completed, syngamy of egg cell nucleus and second male gamete is not yet finished. – 38: Embryo sac showing the young zygote, rest of the still visible pollen tube and first division of the endosperm nucleus. – 39: Longitudinal section of the middle flower of a triplet, the gynoecium is cut transversely right at the plane of fusion of the two carpels, showing both the external as well as the internal pollen tube transmitting tissue. Deep below the stigmatic crests of the style, concealed in the gynoecium, two narrow bands of hairy tissue extend along the margins of the two intimately fused carpels from the top of the ovary downward up to the base of the funicle of the ovule, forming an efficient transmitting tissue (obturator) for the pollen tubes. – 39a: Detail of pollen-tube-transmitting tissue comprising 2- or 3-celled hairs the long-end cells of which are multinucleate and rich in cytoplasm. – 40: Trichome (epidermal vesicle) swollen above the level of the other epidermal cells on the tip of a young tepal.

of the ovary. From there it makes its way down to the micropyle of the single basal ovule. At the inside of the ovary the pollen tube is guided by one of the two narrow bands of inner stigmatic tissue (obturator, Fig. 25, 39) that run on the suture of the two intimately fused carpels, right from the very top of the ovary up to the base of the funicle of the ovule. Fig. 39a shows the inner stigmatic tissue which is made up of a tuft of loosely interwoven 2 to 3-celled trichomes, the end cells of which are multinucleate and rich in cytoplasm. Fig. 35 and 36 show double fertilization. Fig. 37, 38, 41 and 44 present the early and Fig. 50 and 53 the later endosperm development. Finally, Fig. 54–64 document the development of the embryo.

#### **Development of the seed coat**

Seen in longitudinal sections of ovules at the time when the tetrad of megaspores is formed, the outer integument situated in the height of the fertile megaspore consists of two layers of cells, whereas at the chalazal end three layers are present. During further development the number of layers in the outer integument may increase on the chalazal side and especially at the tip in the vicinity of the micropyle. The inner integument consists of two layers of cells except at the tip where it is much thicker (5–6 layers).



Fig. 41–43. Flowers of *Lophiocarpus polystachyus*, *Phytolacca acinosa* and *Chenopodium bonus-henricus*. Longitudinal sections. Overview of ovule, ovary and dwindling nectary; 41: *Lophiocarpus polystachyus.* Somewhat stipitate (gynophore) ovary after fertilization with the dwindling nectary (stippled), pollen-tube-transmitting hairs (obturator) at the foot of the funicle and embryo sac with 2-celled embryo and 4 endosperm nuclei; 41a: *Lophiocarpus polystachyus.* Nectary in higher magnification showing the vacuolated, already degenerating cells; 42: Anthetic flower of *Phytolacca acinosa* (stamens and tepals not drawn) showing gynoecium, nectary and gynophore; 43: Flower of *Chenopodium bonus-henricus* shortly before anthesis, without gynophore



Fig. 44–48. Development of nuclear endosperm and of the integuments; 44: Longitudinal section of part of nucellus showing embryo sac with 2-celled embryo and 4 endosperm nuclei; 45–48: Longitudinal sections of parts of integuments at about half height of the embryo sac of ovules in different developmental stages. – 45: From ovule at the time of uninucleate embryo sac (see Fig. 27). – 46: From ovule with embryo sac showing double fertilization (Fig. 37). – 47: From ovule with embryo sac showing 4 endosperm nuclei and 2-celled embryo (Fig. 41, 44). – 48: From ovule with embryo at "heart stadium" (see Fig. 53).

Both outer as well as the inner integument participate in the formation of the seed coat, which comprises all the layers of the outer integument plus the innermost layer of the inner integument in the still not fully mature seed (oldest examined stage). The outer layer of the inner integument becomes crushed, is resorbed (except at the tip) and disappears during development of the testa (Fig. 45–48). The cells of the outer layer of the outer integument become very conspicuous by excessive radial elongation. Also cells of the remaining (extant) inner layer of the inner integument elongate radially and in both of these cells dark brown deposits occlude (fill in) the cell lumina. Cells of the two inner cell layers of the outer integument have some starch-like deposits and healthy-looking nuclei but no brown deposits. Especially the outer tangential cell walls ("roof" of the cells, building the surface of the seed) of the outer layer of the outer integument show a tremendous increase in thickness.

The innermost layer of the testa is derived from the inner layer of the inner integument.

## **Conclusions and discussion**

*Lophiocarpus polystachyus* has polytelic inflorescences with all flowers being bisexual. The flowers possess a well developed nectary and are promoted on the abaxial side. Five tepals make up the perianth with the adaxial member being a little wider than the others. The androecium is either 4-merous or, more rarely, due to fusion of the two adaxial stamens, 3-merous. Also the somewhat stipitate and unilocular gynoecium with one basally inserted ovule is oligomerous, showing two intimately fused carpels the hairy sutures of which form two narrow bands of inner stigmatic tissue leading down to the foot of the funicle and acting as obturator. Only the abaxial carpel produces the ovule, the adaxial one being sterile. Wall formation of the introrse and tetrasporangiate anther conforms to the Monocotyledonous type and four wall layers including a glandular tapetum with binucleate cells are built. Meiosis of the microsporocytes during which 9 bivalents are visible leads after simultaneous delimitation of the microspores to tetrahedral and decussate pollen tetrads. The tricolpate pollen is 3-celled when shed. The ovule is campylotropous, bitegmic and crassinucellate. Early in its ontogeny a parietal cell is cut, later also a many-layered nucellus epidermis (nucellar cap) develops. The inner integument alone forms the micropyle (endostome) and there is an air space between the inner and outer integument at the chalazal side. Embryo sac development is of the monosporic, *Polygonum* type and at maturity the megagametophyte is 7-celled. Soon after double fertilization the 3 antipodals and the intact synergid degenerate. During early embryogenesis endosperm development is free nuclear and an aggregate of larger nuclei and cytoplasm with coarse-grained contents functions as chalazal haustorium, digesting the tissue around the massive core of the nucellus. After starch deposition, the latter forms the major food reserve (perisperm) of the circular embryo. In late embryogenesis the endosperm becomes cellular and in addition to the perisperm also a scanty residue of cellular endosperm remains clothing the radicle of the embryo. The seed coat comprises all the layers of the outer integument and the inner layer of the inner integument. The outer layer of the inner integument disappears during development of the testa. The fruit is a 1-seeded drupe.

The flower anatomy of *Lophiocarpus* confirms an affinity to the *Phytolaccaceae* s.l. and its distance from the *Chenopodiaceae* because of the missing gynophore in the latter (Fig. 41–43). The ovule in *Lophiocarpus* is built as usual in *Centrospermae*, the embryo sac development is of the common *Polygonum* type, present in all *Centrospermae*, and embryogenesis follows the Caryophyllad type of Johansen (1950). A comparison of the flower morphology and embryology of *Lophiocarpus* with the



Fig. 49–52a. Older stages of endosperm, development of perisperm, of the seed coat and fruit wall. Transverse sections (Fig. 49, 51, 52) and longitudinal section (Fig. 50); 49: Part of the fruit wall and of the integuments, stage between Fig. 47 and 48. – 50: Overview of ovule at the time of the 9-celled embryo with free endosperm nuclei lining the walls of the embryo sac, a basal aggregation of haustorially active coarse granular plasma and larger endosperm nuclei at the chalaza and a circular tissue right in the middle of the ovule that will remain in the ripe seed as perisperm – a starch deposit for the future use of the adult embryo during germination. – 51: Overview of nearly ripe seed showing the massive perisperm with still living cells and nuclei in the middle and the sectioned circular embryo with the two cotyledons above and the radicle below, both surrounded by a ring of scanty cellular endosperm. The seed wall (testa), especially the outer wall of the outer integument is already heavily incrusted with dark brownish black substances. – 52–52a: Transverse section of 150-celled embryo surrounded by cellular endosperm from the micropylar side of an ovule and in Fig. 52a the haustorially active nuclear endosperm and coarse plasma aggregation from the chalaza of the same ovule.



Fig. 53–64. Development of the endosperm and embryo. Longitudinal sections; Fig. 53: Overview of ovule with embryo shortly before initiation of the cotyledons surrounded by cellular endosperm reaching to one third of the length of the embryo sac followed by free nuclear endosperm lining the walls of the rest of the embryo sac. The nuclear endosperm at the chalaza is composed of a multitude of bigger nuclei and coarse granular plasma forming an effective haustorial basal aggregation. – 53a–b: micropylar (Fig. 53a) and chalazal (Fig. 53b) part of Fig. 53 in higher magnification; Fig. 54–64: Embryogenesis from zygote to the 31-celled stage, small size numbers with the prefix  $(E = endosperm)$  give the number of endosperm nuclei accompanying that embryo in the embryo sac.  $-54$ : Zygote, E = 2 (accompanied by 2 endosperm nuclei). – 55–56: Embryo, 2-celled, both with  $E = 4. -57$ : Embryo, 3-celled,  $E = 8$  and 3 antipodals. – 58: Embryo, 4-celled, E = 22. – 59: Embryo, 7-celled, E = 34. – 60: Embryo, 7-celled, E = 37. – 61: Embryo, 9-celled, E = 42. – 62: Embryo, 21-celled, E = 88. – 63: Embryo, 19–21-celled, E  $= 63. - 64$ : Embryo, 33-celled, E = 135 free nuclei.

data known for *Corbichonia* (Narayana & Lodha 1963 under *Orygia decumbens*), to which *Lophiocarpus* presumably has a closer relationship (Cuénoud & al. 2002), indicates that they are not intimately related because of the following differences between them:

(1) Flower morphology:

*Corbichonia*: Flowers with 5 free tepals. Androecium of numerous stamens of which those in the outer 2 or 3 whorls are petaloid staminodia. The gynoecium is pentacarpellary with 2 rows of ovules in each locule on axile placenta.

*Lophiocarpus*: Flowers with 5 free tepals. Androecium of only 4 or 3 stamens. The gynoecium is bicarpellate and unilocular with only one basally inserted ovule. (2) Ovule:

*Corbichonia*: The ovule displays the primordium of an arillus.

*Lophiocarpus*: The ovule shows no arillar primordium.

(3) Obturator:

*Corbichonia*: Epidermal cells of the placenta serve as a placental obturator.

*Lophiocarpus*: Epidermal cells of the funicle act as a funicular obturator.

(4) Development of the embryo:

*Corbichonia*: Embryogenesis conforms to the *Linum* variation, Solanad type.

*Lophiocarpus*: Embryogenesis conforms to the Caryophyllad type of Johansen (1950).

# **Acknowledgements**

The author is grateful to the reviewers, Dietmar Behnke (Heidelberg) and Hiroshi Tobe (Tokyo), for their valuable comments on a previous version of the manuscript, and to the editor, Norbert Kilian, who reviewed the manuscript and improved the Introduction substantially.

# **References**

- APG III 2009: An update of the Angiosperm Phylogeny Group classification for the orders and families of flowering plants. – [Bot. J. Linn. Soc.](http://dx.doi.org/10.3372/wi.39.39201) **[161:](http://dx.doi.org/10.3372/wi.39.39201)** [105–121](http://dx.doi.org/10.3372/wi.39.39201) [online at[: http://dx.doi.org/10.3372/wi.39.39201\].](http://dx.doi.org/10.3372/wi.39.39201])
- Behnke H. D. 1974: Electronenmikroskopische Untersuchungen an Siebrohren-Plastiden und ihre Aussage über die systematische Stellung von *Lophiocarpus*. – Bot. Jahrb. Syst. **94:** 114–119.
- Bittrich V. 1993: Introduction to *Centrospermae.* Pp. 13–19 in: Kubitzki K., Rohwer J. G. & Bittrich V. (ed.), The families and genera of vascular plants **2.** Flowering plants. Dicotyledons. Magnoliid, Hamamelid and Caryophillid families. – Berlin: Springer.
- Bortenschlager S. 1973: Morphologie pollinique des *Phytolaccaceae*. – Pollen & Spores **15:** 227–253.
- Brockington S. F., Alexandre R., Ramdial J., Moore M. J., Crawley S., Dhingra A., Hilu K., Soltis D. S. & Soltis P. S. 2009: Phylogeny of the *Caryophyllales* sensu lato: revisiting hypotheses on pollination biology and perianth differentiation in the core *Caryophyllales*. – [Int. J. Pl. Sci.](http://dx.doi.org/10.1086/597785) **[170:](http://dx.doi.org/10.1086/597785)** [627–643.](http://dx.doi.org/10.1086/597785)
- Brown G. K. & Varadarajan G. S. 1985: Studies in *Caryophyllales* I. Re-evaluation of classification of *Phytolaccaceae* s.l. – [Syst. Bot.](http://dx.doi.org/10.2307/2418434) **[10:](http://dx.doi.org/10.2307/2418434)** [49–63.](http://dx.doi.org/10.2307/2418434)
- Brown N. E. 1909: List of plants collected in Ngamiland and the northern part of Kalahari Desert, chiefly in the

neighbourhood of Kwebe and along Botltle and Lake rivers. – Kew Bull. **1909:** 89–146.

- Cuénoud P., Savolainen V., Chatrou L. W., Powell M., Grayer R. J. & Chase M. W. 2002: Molecular phylogenetics of *Caryophyllales* based on nuclear 18S rDNA and plastid rbcL, atpB, and matK DNA sequences. [– Amer. J. Bot.](http://dx.doi.org/10.3732/ajb.89.1.132) **[89:](http://dx.doi.org/10.3732/ajb.89.1.132)** [132–144.](http://dx.doi.org/10.3732/ajb.89.1.132)
- Davis G. L. 1966: Systematic embryology of the angiosperms. – New York: Wiley.
- Doweld A. & Reveal J. L. 2008: New suprageneric names of vascular plants. – Phytologia **90:** 416–417.
- Eckardt T. 1964: *Centrospermae.* Pp. 79–101 in: Melchior H. (ed.), A. Engler's Syllabus der Pflanzenfamilien, ed. 12, **2.** – Berlin: Borntraeger.
- Eckardt T. 1974: Vom Blütenbau der Centrospermen-Gattung *Lophiocarpus* Turcz. – Phyton **16:** 13–27.
- Hakki M. I. 1972: Blütenmorphologische und embryologische Untersuchungen an *Chenopodium capitatum* und *Chenopodium foliosum* sowie weiteren *Chenopodiaceae*. – Bot. Jahrb. Syst. **92:** 178–330.
- Heimerl B. 1934: *Phytolaccaceae.* Pp. 161–163 in: Engler A. & Prantl K. (ed.), Die natürlichen Pflanzenfamilien, ed. 2, 16c. – Leipzig: Engelmann.
- Hutchinson J. 1959: The families of flowering plants, ed. 2. – Oxford: Clarendon.
- Hutchinson J. 1973: The families of flowering plants, ed. 3. – Oxford: Clarendon.
- Johansen D. A. 1950: Plant embryology. Waltham, Mass.: Chronica Botanica.
- Johri B. M., Ambegaokar K. B. & Srivastava P. S. 1992: Comparative embryology of angiosperms 1. – Berlin: Springer.
- Kajale L. B. 1954: A Contribution to the Embryology of the *Phytolaccaceae*. II. Fertilization and the development of Embryo, Seed and Fruit in *Rivina humilis* Linn. and *Phytolacca dioica* Linn. – J. Indian Bot. Soc. **33:** 206–225.
- Mauritzon J. 1934: Ein Beitrag zur Embryologie der Phytolaccaceen und Cactaceen. – Bot. Notiser **1934:**  111–135.
- Narayana H. S., & Lodha B. C. 1963: Embryology of *Orygia decumbens* Forsk. (*Aizoaceae*). – Phytomorphology **13:** 54–59.
- Nowicke J. W. 1968: Palynotaxonomic study of the *Phytolaccaceae*. – [Ann. Missouri Bot. Gard.](http://dx.doi.org/10.2307/2395128) **[55:](http://dx.doi.org/10.2307/2395128)** [294–364.](http://dx.doi.org/10.2307/2395128)
- Rocen T. 1927: Zur Embryologie der Centrospermen. Diss. Uppsala.
- Rohwer J. G. 1993: *Phytolaccaceae*. Pp. 506–515 in: Kubitzki K., Roher J. G. & Bittrich V. (ed.), The families and genera of vascular plants **2.** – Berlin: Springer.
- Stevens P. F. 2001+: Angiosperm Phylogeny Website. Version 12, July 2012 [and more or less continuously updated since]. – Published at [http://www.mobot.org/](http://www.mobot.org/MOBOT/research/APweb/) [MOBOT/research/APweb/](http://www.mobot.org/MOBOT/research/APweb/) [accessed 26 Oct 2012].