

## **From India to Madeira and back again: a new combination for a wide-ranging *Spergularia* (Caryophyllaceae)**

Author: Turner, Ian M.

Source: *Willdenowia*, 47(3) : 213-216

Published By: Botanic Garden and Botanical Museum Berlin (BGBM)

URL: <https://doi.org/10.3372/wi.47.47303>

---

BioOne Complete ([complete.BioOne.org](https://complete.BioOne.org)) is a full-text database of 200 subscribed and open-access titles in the biological, ecological, and environmental sciences published by nonprofit societies, associations, museums, institutions, and presses.

Your use of this PDF, the BioOne Complete website, and all posted and associated content indicates your acceptance of BioOne's Terms of Use, available at [www.bioone.org/terms-of-use](https://www.bioone.org/terms-of-use).

Usage of BioOne Complete content is strictly limited to personal, educational, and non-commercial use. Commercial inquiries or rights and permissions requests should be directed to the individual publisher as copyright holder.

---

BioOne sees sustainable scholarly publishing as an inherently collaborative enterprise connecting authors, nonprofit publishers, academic institutions, research libraries, and research funders in the common goal of maximizing access to critical research.

IAN M. TURNER<sup>1\*,2</sup>

## From India to Madeira and back again: a new combination for a wide-ranging *Spergularia* (*Caryophyllaceae*)

Version of record first published online on 9 November 2017 ahead of inclusion in December 2017 issue.

**Abstract:** A recently published phylogenetic analysis strongly supports the inclusion of *Spergula fallax* (Lowe) E. H. L. Krause in the genus *Spergularia* (Pers.) J. Presl & C. Presl (*Caryophyllaceae*), where it was originally described. This species of spurrey, which occurs across North Africa, the Middle East and N India, has also been named *Arenaria flaccida* Roxb., which is a later homonym and therefore illegitimate. *Stipularia flaccida* Madden is an overlooked replacement name for *A. flaccida* and is the earliest legitimate name for the species. The new combination *Spergularia flaccida* (Madden) I. M. Turner is proposed here. In addition, lectotypes are designated for *A. flaccida* and its taxonomic synonyms *Lepigonum eximium* Kindb., *Spergula flaccida* f. *purpurascens* Maire & Sauvage, *Spergula pentandra* var. *intermedia* Boiss. and *Spergularia fallax* Lowe.

**Key words:** Ascherson, *Caryophyllaceae*, lectotypification, Madden, Middle East, new combination, nomenclature, North Africa, Roxburgh, *Spergula*, *Spergularia*

**Article history:** Received 29 August 2017; peer-review completed 9 October 2017; received in revised form 11 October 2017; accepted for publication 12 October 2017.

**Citation:** Turner I. M. 2017: From India to Madeira and back again: a new combination for a wide-ranging *Spergularia* (*Caryophyllaceae*). – Willdenowia 47: 213–216. doi: <https://doi.org/10.3372/wi.47.47303>

### Introduction

William Roxburgh's magnum opus, his *Flora Indica*, was never published in the format the author intended. He accumulated manuscript descriptions of many plant species and had drawings made of a large proportion of these. But by his death in 1815, the flora was still only a manuscript. It was not until 1832 (Roxburgh 1832), when the second edition of the posthumous *Flora Indica* appeared, that all Roxburgh's seed plant taxa were finally published (though still without the illustrations). One of the many new species first validated in the 1832 *Flora Indica*, was a small *Caryophyllaceae* species that Roxburgh called *Arenaria flaccida*. He reported that "it is only found during the cold season as a weed in gardens about Calcutta".

In 1856, Richard Lowe, described *Spergularia fallax* from Madeira (Lowe 1856). This is some 10,000 km from where Roxburgh described *Arenaria flaccida* in Bengal. Yet, the following year, Lowe (1857: 75) noted that his species was the same as Roxburgh's. The species, ranging from Macaronesia, across North Africa, the Middle East and on to North India, has been recognized by many authors, generally in the genus *Spergula* L. However, a recently published phylogeny (Kool & Thulin 2017), based on molecular data, has demonstrated that the species is sister to the rest of *Spergularia* (Pers.) J. Presl & C. Presl, but definitely lies outside a monophyletic *Spergula*. The authors recommended that the species be treated as a member of *Spergularia*. The aim of this paper is to briefly review the nomenclature of this spurrey species of extensive distribution.

1 Singapore Botanical Liaison Officer, Identification and Naming, Herbarium, Royal Botanic Gardens, Kew, Richmond, Surrey TB9 3AE, U.K.; \*e-mail: [i.turner@kew.org](mailto:i.turner@kew.org)

2 Herbarium, Singapore Botanic Gardens, NParks, 1 Cluny Road, Singapore 259569.

## Nomenclature

Ascherson (1889) first transferred Roxburgh's name to *Spergula*. However, Roxburgh's supposed basionym is actually a later homonym and thus illegitimate, as first noted by Milne-Redhead (1951) at the instigation of J. E. Dandy. Therefore, *Spergula fallax* (Lowe) E. H. L. Krause, based on *Spergularia fallax* Lowe, appears to be the correct name for the plant in *Spergula* and has been used by many authors (Burt & Lewis 1952: 349; Nair 1964; Rechinger 1964: 229; Zohary 1966: 122; Ratter 1980: 28; Greuter & al. 1984: 282; Ratter 1990: 149; Townsend & al. 2016: 23), though with a notable exception still using *S. flaccida* (Raffaelli 1983). In researching Roxburgh's botanical output, I found that there is an overlooked publication of a combination based on *Arenaria flaccida*. Madden (1849) published supplementary notes on the flora of Kumaon in N India, in which he proposed *Stipularia flaccida* based on "*Arenaria flaccida* of Roxburgh". As Roxburgh's name is illegitimate, Madden effectively provided for *Arenaria flaccida* a replacement name, which is the oldest legitimate name for the taxon at specific rank. While Ascherson did not cite Madden's name in publishing *Spergula flaccida*, Art. 41.4 of the *International Code of Nomenclature for algae, fungi, and plants* (McNeill & al. 2012) applies, as Ascherson's name would otherwise be validly published as that of a new taxon supported by the diagnosis provided by Ascherson's key to species. Ascherson's intention of making a new combination based on Roxburgh's name is clear. Therefore, *Spergula flaccida* (Madden) Asch. is the correct name in *Spergula* for *Spergula fallax* (Lowe) E. H. L. Krause. But following the recommendation of Kool & Thulin (2017), the species should be considered a member of *Spergularia*, as indicated by its tricarpetate, rather than pentacarpetate, flowers. The binomial *Spergularia flaccida* has appeared in print on several occasions (Battandier & Trabut 1911: 625; Kunkel 1967: 25; Hansen 1979: 110), I suspect mostly with *Spergularia* in error for *Spergula*, but none seems to even approach being validly published as a new combination. I therefore provide the required new combination below.

## Typification

Surprisingly for a very wide-ranging taxon that has been treated in various floras, there do not seem to have been any published typifications for these names and other synonyms. Forman (1997) did not list any Roxburgh herbarium specimens for *Arenaria flaccida*, and I have not located any subsequently. There is a Roxburgh drawing of the species; I therefore designate here the copy at Kew as the lectotype.

Lowe's herbarium was divided between K and BM after his death. All the material in K seems to have been collected after the original publication of *Spergularia fallax*, but BM has some earlier specimens, among which

one from a locality mentioned in the protologue is designated here as the lectotype. Lowe appears to have used taxon numbers rather than specimen numbers for his collections. Most of the BM collections of *S. flaccida* are numbered 857 despite being gathered from different locations and on different dates.

The original material (at least four syntypes) for Boissier's *Spergula pentandra* var. *intermedia* is present in the Boissier Herbarium in Geneva. I here select the Griffith specimen from Afghanistan as the lectotype. It seems likely that Griffith's number here is a taxon number (Lamond 1970). There are two specimens with the same number in K. One is from Bentham's herbarium that is marked as originally coming from Lemann's herbarium, which is also the source of Boissier's specimen. I therefore consider it more likely that these two specimens are true duplicates (i.e. obtained from the division of a specimen Lemann received from Griffith) and cite the K specimen as a possible isolectotype.

One of the two syntypes cited by Kindberg is designated here as the lectotype of *Lepigonium eximium* Kindb., and a syntype in Montpellier is designated as the lectotype of *Spergula flaccida* f. *purpurascens* Maire & Sauvage.

- Spergularia flaccida* (Madden) I. M. Turner, comb. nov.**  
 ≡ *Stipularia flaccida* Madden in J. Asiat. Soc. Bengal 18: 621. 1849 ≡ *Arenaria flaccida* Roxb., Fl. Ind., ed. 1832, 2: 447. 1832, nom. illeg. (Art. 53.1) [non *A. flaccida* All., Auct. Syn.: 35. 1773, nec *A. flaccida* Clairv., Man. Herbor. Suisse: 149. 1811, nec *A. flaccida* Schleich., Cat. Pl. Helv., ed. 3: 46 (add.). 1815] ≡ *Spergula flaccida* (Madden) Asch. in Verh. Bot. Vereins Prov. Brandenburg 30: XLIII. 1889 ≡ *Spergula pentandra* subsp. *flaccida* (Madden) Jafri, Fl. Karachi: 124. 1966, – **Lectotype (designated here):** [icon] Icones Roxburghianae No. 1141 (K) – Fig. 1.  
 – *Spergula indica* Royle, Ill. Bot. Himal. Mts.: 78, 464. 1834, nom. inval. (Art. 38.1(a)).  
 = *Spergula pentandra* var. *intermedia* Boiss., Diagn. Pl. Orient., ser. 2, 1: 93. 1854. – **Lectotype (designated here):** Afghanistan, W. Griffith 1629 (G-BOIS G00547728; possible isolectotype: K 000723332).  
 = *Spergularia fallax* Lowe in Hooker's J. Bot. Kew Gard. Misc. 8: 289. 1856 ≡ *Lepigonum fallax* (Lowe) Lowe, Fl. Salvag. Tent.: 9. 1869 ≡ *Spergula fallax* (Lowe) E. H. L. Krause, Deutschl. Fl. [Sturm], ed. 2, 5: 19. 1901. – **Lectotype (designated here):** Madeira, [Ilhas Desertas, Bugio,] top of S. Dez.<sup>a</sup>, 18 Jun 1855, R. T. Lowe 857 (BM 000072083).  
 = *Lepigonum eximium* Kindb., Monogr. Gen. Lepigon.: 32. 1863. – **Lectotype (designated here):** Palestine, 1855, T. Kotschy iter Syriacum 516 (S S17-522233; isolectotype: UPS).  
 = *Spergula flaccida* f. *purpurascens* Maire & Sauvage in Bull. Soc. Hist. Nat. Afrique N. 32: 204. 1941. – **Lectotype (designated here):** Libya, Tripolitania, in arenosis amnis Ganima, 14 Apr 1938, R. Maire & M. Weiller 274 (MPU 004482).



Fig. 1. Scan of the lectotype of *Arenaria flaccida* Roxb. – Icones Roxburghianae No. 1141 (K). – Reproduced with the kind permission of the Director and the Board of Trustees, Royal Botanic Gardens, Kew.



## Acknowledgements

Many thanks to Julia Buckley (K) for providing the scan of the Roxburgh drawing, Jovita Yesilyurt (BM), Laurence Loze (G), Arne Anderberg (S) and Mats Hjertson (UPS) for providing images of specimens from their respective herbaria. The reviewers, Richard Rabeler (MICH) and Abigail Moore (BRU, MJG, OKL), provided helpful suggestions that led to significant improvements in the paper.

## References

- Ascherson P. 1889: [Adventivpflanzen der Oelfabrik in Mannheim (Fr. Lutz), namentlich *Spergula flaccida* (Roxb.) Aschers. (*Spergula fullax* [sic!] Lowe) – Title of paper only appears in the volume contents list]. – Verh. Bot. Vereins Prov. Brandenburg **30**: XXXI–XLIV.
- Battandier J.-A. & Trabut L. 1911: Contribution à la flore du pays des Touaregs. – Bull. Soc. Bot. France **58**: 623–629.
- Burt B. L. & Lewis P. 1952: On the flora of Kuwait: II. – Kew Bull. **7**: 333–352.
- Forman L. 1997: Notes concerning the typification of names of William Roxburgh's species of phanerogams. – Kew Bull. **52**: 513–534.
- Greuter W., Burdet H. M. & Long G. 1990: Med-checklist **I**. – Genève: Conservatoire et Jardin botaniques, Ville de Genève.
- Hansen A. 1979: Some floristic notes for the Canary Islands. – Vieraea **8**: 107–112.
- Kool A. & Thulin M. 2017: A giant spurrey on a tiny island: on the phylogenetic position of *Sanctambrosia manicata* (Caryophyllaceae) and the generic circumscriptions of *Spergula*, *Spergularia* and *Rhodalsine*. – Taxon **66**: 615–622.
- Kunkel G. 1967: Plantas vasculares: nuevas adiciones para la flora de Gran Canaria. – Cuad. Bot. Canaria **2**: 23–27.
- Lamond J. M. 1970: The Afghanistan collections of William Griffith. – Notes Roy. Bot. Gard. Edinburgh **30**: 159–175.
- Lowe R. T. 1856: Species plantarum maderensium quaedam novae, vel hactenus ineditae, breviter descriptae. – Hooker's J. Bot. Kew Gard. Misc. **8**: 289–302.
- Lowe R. T. 1857: On *Ranunculus Creticus*, *cortusaefolius*, and *grandifolius* of authors. – Hooker's J. Bot. Kew Gard. Misc. **9**: 65–75.
- Madden E. 1849: Supplementary notes to "The Turae and Outer Mountains of Kumaoon;" Journal of the Asiatic Society, Bengal, May and June, 1848. – J. Asiat. Soc. Bengal **18**: 603–644.
- McNeill J., Barrie F. R., Buck W. R., Demoulin V., Greuter W., Hawksworth D. L., Herendeen P. S., Knapp S., Marhold K., Prado J., Prud'homme van Reine W. F., Smith G. F., Wiersema J. H. & Turland N. J. 2012: International Code of Nomenclature for algae, fungi, and plants (Melbourne Code) adopted by the Eighteenth International Botanical Congress Melbourne, Australia, July 2011. – Königstein: Koeltz Scientific Books. – Regnum Veg. **154**.
- Milne-Redhead E. 1951: *Caryophyllaceae*. – Kew Bull. **5**: 338.
- Nair N. C. 1964: Note on *Spergula fallax* in northern India. – Bull. Bot. Survey India **9**: 277.
- Raffaelli M. 1983: Il genere «*Spergula*». – Webbia **37**: 1–12.
- Ratter J. A. 1980: *Spergula*. – Pp. 27–28 in: Chaudhri M. N., Ratter J. A. & Rechinger K. H., Flora iranica **144**. – Graz: Akademische Druck- u. Verlagsanstalt.
- Ratter J. A. 1990: *Spergula* L. – Pp. 145–149 in: Castroviejo S., Laínz M., López González G., Montserrat P., Muñoz Garmendia F., Paiva J. & Villar L. (ed.), Flora iberica **II**. – Madrid: Real Jardín Botánico, CSIC.
- Rechinger K. H. 1964: Flora of lowland Iraq. – Weinheim: Verlag von J. Cramer.
- Roxburgh W. 1832: Flora indica; or, descriptions of Indian plants [ed. 1832]. **1–3**. – Serampore: W. Thacker and Co.; Parbury, Allen and Co.
- Townsend C. C., Melzheimer V., Kandemir A., Ghazanfar S. A. & Haloob A. 2016: *Caryophyllaceae*. – Pp. 6–123 in: Ghazanfar S. A. & Edmondson J. R. (ed.), Flora of Iraq **5(1)**. – Richmond: Royal Botanic Gardens, Kew.
- Zohary M. 1966: Flora palaestina **1**. Text. *Equisetaceae* to *Moringaceae*. – Jerusalem: Israel Academy of Sciences and Humanities.

## Willdenowia

Open-access online edition [www.bioone.org/loi/will](http://www.bioone.org/loi/will) 

Online ISSN 1868-6397 · Print ISSN 0511-9618 · Impact factor 0.680

Published by the Botanic Garden and Botanical Museum Berlin, Freie Universität Berlin

© 2017 The Author · This open-access article is distributed under the CC BY 4.0 licence