

Multivariate morphometric analysis of Petrorhagia subsect. Saxifragae (Caryophyllaceae) in Greece, with a new species from SE Peloponnisos: P. laconica

Authors: Trigas, Panayiotis, Kougioumoutzis, Konstantinos, Ermidou, Aikaterini, and Kalpoutzakis, Eleftherios

Source: Willdenowia, 48(1): 137-146

Published By: Botanic Garden and Botanical Museum Berlin (BGBM)

URL: https://doi.org/10.3372/wi.48.48110

BioOne Complete (complete.BioOne.org) is a full-text database of 200 subscribed and open-access titles in the biological, ecological, and environmental sciences published by nonprofit societies, associations, museums, institutions, and presses.

Your use of this PDF, the BioOne Complete website, and all posted and associated content indicates your acceptance of BioOne's Terms of Use, available at <u>www.bioone.org/terms-of-use</u>.

Usage of BioOne Complete content is strictly limited to personal, educational, and non - commercial use. Commercial inquiries or rights and permissions requests should be directed to the individual publisher as copyright holder.

BioOne sees sustainable scholarly publishing as an inherently collaborative enterprise connecting authors, nonprofit publishers, academic institutions, research libraries, and research funders in the common goal of maximizing access to critical research.



PANAYIOTIS TRIGAS^{1*}, KONSTANTINOS KOUGIOUMOUTZIS¹, AIKATERINI ERMIDOU¹ & ELEFTHERIOS KALPOUTZAKIS²

Multivariate morphometric analysis of *Petrorhagia* subsect. *Saxifragae* (*Caryophyllaceae*) in Greece, with a new species from SE Peloponnisos: *P. laconica*

Version of record first published online on 5 April 2018 ahead of inclusion in April 2018 issue.

Abstract: The intra-generic relationships within the intriguing genus *Petrorhagia* are currently debated, as the genus is underrepresented in recent taxonomic studies. In this study, we investigate the relationships among the species of *Petrorhagia* [sect. *Petrorhagia*] subsect. *Saxifragae* occurring in Greece. Based on a combination of ordination methods and discriminant analyses of 36 macromorphological characters, five species are recognized in Greece. The analyses showed that all currently known species from Greece (*P. fasciculata, P. graminea, P. phthiotica* and *P. saxifraga*) represent distinct taxonomic entities. According to our results, two recently collected populations from SE Peloponnisos represent a fifth, new *Petrorhagia* species, which is first described and illustrated here as *P. laconica*. It is related to the widespread *P. saxifraga* and the Greek endemic *P. graminea*, but clearly distinct from both species by a combination of morphological characters, including indumentum, inflorescence, calyx, capsule and seed characters.

Key words: *Caryophyllaceae*, Greece, morphology, multivariate morphometric analysis, new species, Peloponnisos, *Petrorhagia*, *Petrorhagia* subsect. *Saxifragae*, taxonomy

Article history: Received 18 December 2017; peer-review completed 7 February 2018; received in revised form 14 February 2018; accepted for publication 20 February 2018.

Citation: Trigas P., Kougioumoutzis K., Ermidou A. & Kalpoutzakis E. 2018: Multivariate morphometric analysis of *Petrorhagia* subsect. *Saxifragae* (*Caryophyllaceae*) in Greece, with a new species from SE Peloponnisos: *P. laconica.* – Willdenowia 48: 137–146. doi: https://doi.org/10.3372/wi.48.48110

Introduction

Caryophyllaceae include approximately 3000 species, distributed in about 100 genera (Hernández-Ledesma & al. 2015). The family has a primarily N-temperate distribution with a diversity centre in the E Mediterranean and Irano-Turanian regions, while presence in the tropics and the S hemisphere is limited and mostly confined to higher elevations (Bittrich 1993; Greenberg & Donoghue 2011; Rabeler & Hartman 2005; Hernández-Ledesma & al. 2015). The phylogenetic relationships within *Caryophyllaceae* is a field of active research (Fior & al. 2006; Harbaugh & al. 2010; Greenberg & Donoghue 2011;

Hernández-Ledesma & al. 2015, etc.) and genera delimitation has been altered in some cases, reflecting the results of recent molecular studies (e.g. Dillenberger & Kadereit 2014).

Petrorhagia (Ser.) Link (*Caryophylloideae*, *Caryophylleae*) is a small genus of c. 33 species, distributed in the Mediterranean area, Europe and W Asia (from the Canary Islands east to Kashmir), with a diversity centre in Greece and Turkey (Ball & Heywood 1964; Georgiou 1997). In the past, it was placed in different genera such as *Dianthus* L., *Gypsophila* L., *Kohlrauschia* Kunth and *Tunica* Scop. Evidence from both traditional and phylogenetic systematics support its recognition as a distinct

¹ Laboratory of Systematic Botany, Department of Crop Science, Agricultural University of Athens, Iera Odos 75, 118 55, Athens, Greece; *e-mail: trigas@aua.gr (author for correspondence).

² Division of Pharmacognosy and Chemistry of Natural Products, School of Pharmacy, National and Kapodistrian University of Athens, 157 71 Athens, Greece.

taxon at the genus rank, which clusters as sister to a clade including *Dianthus* and *Velezia* L. (Ball & Heywood 1964; Harbaugh & al. 2010; Greenberg & Donoghue 2011; Pirani & al. 2014; Hernández-Ledesma & al. 2015; Hilooğlu & al. 2016).

Recent data on the infrageneric classification of Petrorhagia is still scarce, as the genus is rather poorly sampled. The available data (Greenberg & Donoghue 2011; Hilooğlu & al. 2016) are fragmentary and do not allow a reliable infrageneric classification. As a result, most recent treatments of the genus follow the monograph of Ball & Heywood (1964) that recognizes five sections within Petrorhagia, distinguished by petal and seed characters, life-cycle, leaf venation and the presence/absence of epicalyx bracts. Out of the five sections, P. sect. Petrorhagia includes ten species (seven of them occurring in Greece and five endemic to the country) distributed in two subsections: P. subsect. Saxifragae Ball & Heywood and P. subsect. Thessalae Ball & Heywood. The two subsections are distinguished by stem branching (much branched vs simple or with few branches), inflorescence (flowers solitary or fasciculate vs capitate) and bract (lanceolate or ovate, membranous, 1-veined vs broadly ovate or suborbicular, membranous and 1-veined or brown-scarious and many-veined) morphological characters.

In May 2014, during field work in SE Peloponnisos, we detected a population of an interesting *Petrorhagia* species, growing on coastal sand-dunes, an unusual habitat for the members of the genus. Careful examination of the collected specimens revealed that plants from SE Peloponnisos belong to *P*. sect. *Petrorhagia*, and they resemble *P. saxifraga* (L.) Link, a widely distributed species in C and S Europe and SW Asia. After a detailed morphological study, it became apparent that plants from SE Peloponnisos are clearly distinct from *P. saxifraga*, as well as from all known *Petrorhagia* species and they belong to a new, undescribed species. Dr E. Kalpoutzakis had also found the same species in May 2013 in another locality of SE Peloponnisos.

The morphological characters of the new species from SE Peloponnisos, described here as Petrorhagia laconica, support its classification in P. subsect. Saxifragae. Species delimitation within P. subsect. Saxifragae has never been studied in detail. All species of P. subsect. Saxifragae, except P. riphaea (Pau & Font Quer) Ball & Heywood that is the only species of this group distributed in Africa (Morocco), are distributed in Greece. Thus, the scope of this study is manifold; it aims to resolve the apparent ambiguities within P. subsect. Saxifragae and to evaluate the taxonomic validity of the two recently discovered deviating Petrorhagia populations. More specifically, we aim: (1) to clarify the patterns of morphological variation within P. subsect. Saxifragae in Greece, (2) to disclose the level of morphological differentiation for recognized species, (3) to indicate the most informative characters for the identification of the species, and (4) to describe the newly discovered species.

Material and methods

Plant material

Data for the multivariate analyses were recorded from herbarium specimens deposited in UPA and ATH (acronyms follow Thiers 2017+) belonging to Petrorhagia fasciculata (Margot & Reut.) P. W. Ball & Heywood, P. graminea (Sm.) P. W. Ball & Heywood, P. phthiotica (Boiss. & Heldr.) P. W. Ball & Heywood and P. saxifraga. We collected several specimens of P. laconica in the field. All specimens included in the analyses were selected so as to represent the entire distribution range of the taxa in Greece, as well as the morphological variation in each species. Only well-preserved and intact specimens were considered in the analyses. The total number of herbarium specimens included in the analyses was 97. A list of the examined specimens is provided in Table S1 (see Supplementary Material online). Each specimen was preliminary identified as one of the five aforementioned species. Species identification and nomenclature are according to Ball & Heywood (1964), Georgiou (1997) and Dimopoulos & al. (2013).

Characters recorded

Twenty-seven quantitative and 9 qualitative (binary) morphological characters were scored for the herbarium specimens included in our analyses (Table 1). The selected characters represent those previously quoted as diagnostic and are the ones most commonly used in the latest comprehensive monographs, regional and local floras (Ball & Heywood 1964; Coode & Cullen 1967; Georgiou 1997). Capsule and seed morphological characters, although with a diagnostic value in the genus Petrorhagia (Ball & Heywood 1964), were excluded from the morphometric analyses as most studied specimens were in flower, lacking capsules and seeds. These characters, however, were used for the documentation of morphological differences among P. laconica and its closest relatives. All the characters in the herbarium specimens were measured under a stereomicroscope using a ruler with the precision of 0.5 mm.

Morphometric analyses

Basic statistical parameters (mean, minimum and maximum value, standard deviation, 5th and 95th percentiles) were calculated for each species included in the analyses. We used a combination of ordination methods and discriminant analyses in the morphometric analyses of the specimens (Marhold 2011).

We first tested whether any character deviated from a normal distribution using the Shapiro-Wilk statistic

Table 1.	Characters	used in	the morp	hometric	analyses
			1		2

Character abbreviation	Detailed character definition			
al	Number of flowering stems			
a2	Length of flowering stems (cm)			
a3	Length of the unbranched part of flowering stem (cm)			
a4	Length of the inflorescence (cm)			
a5	Glandular hairs present at the lower 1/3 of the stem			
a6	Glandular hairs present at the middle 1/3 of the stem			
a7	Glandular hairs present at the upper 1/3 of the stem			
a8	Eglandular hairs present at the lower 1/3 of the stem			
a9	Eglandular hairs present at the middle 1/3 of the stem			
a10	Eglandular hairs present at the upper 1/3 of the stem			
a11	Number of flowers per flowering stem			
a12	Number of lateral branches per flowering stem			
a13	Length of lateral branches per flowering stem (cm)			
a14	Number of flowers per lateral branch			
a15	Number of flowers per terminal unit of the inflorescence			
a16	Length of lower leaves (mm)			
a17	Width of lower leaves (mm)			
a18	Length of middle leaves (mm)			
a19	Width of middle leaves (mm)			
a20	Length of internodes at the middle of the stem (mm)			
a21	Number of leaves at the middle of the stem			
a22	Ratio of leaves/internodes length at the middle of the sten			
a23	Number of veins on leaves			
a24	Length of petals (mm)			
a25	Width of petals (mm)			
a26	Length of calyx (mm)			
a27	Length of calyx teeth (mm)			
a28	Width of calyx teeth at their base (mm)			
a29	Length of pedicels (mm)			
a30	Bract length (mm)			
a31	Ratio pedicel length/calyx length			
a32	Ratio petal length/calyx length			
a33	Number of veins on sepals			
a34	Glandular hairs present on calyx			
a35	Eglandular hairs present on calyx			
a36	Epicalyx bracts present			

and we log₁₀-transformed any characters that deviated from normality. Then, we computed the correlation coefficients between each character pair in order to reveal highly correlated characters and to ensure that no high correlations (> 0.95 – Španiel & al. 2017) were present that could potentially distort the discriminant analysis. As a second step, we performed several stepwise canonical and classificatory Discriminant Analyses (DAs) in order to: (1) identify the variables having the highest potential as diagnostic characters and (2) test the effectiveness of the discrimination rules by using the Leave-One-Out cross-validation (LOO). The value of the inclusion and the exclusion criterion in the stepwise DAs was set at F = 0.05 and F = 0.1, respectively. Wilk's lambda was used to assess the significance of each variable and Discriminant Function (DF). The Maximum Chance Criterion (MCC) and the Proportional Chance Criterion (PCC) were used to determine whether the prediction equation was better than random chance (Huberty & Olejnik 2006). The DAs were checked for outliers. Finally, we examined the morphological variation in our dataset in relation to the species boundaries by analysing the ranges of the characters using core functions in the R computing environment (R Core Team 2015). All the analyses were carried out using IBM SPSS 24. The correlation coefficients were calculated with the "usdm" (Naimi & al. 2014) package, while the stepwise discriminant analysis plots were visualized via the ggplot2 (Wickham, 2009) package in the R computing environment. The Maximum and Proportional Chance criteria were ran under the Zclass algorithm kindly provided by John D. Morris (Florida Atlantic University).

Results

Characters scored

The basic statistical parameters for all species are given in Table S2. The only character that had a correlation coefficient above the threshold was a10 (eglandular hairs present at the upper $\frac{1}{3}$ of the stem) and was thus excluded from further analyses.

Multivariate analyses

The DA clearly discriminated the five *Petrorhagia* species. Sixteen of the characters emerged as statistically significant (Table 2), with LOO showing a 100 % correct classification, being significantly better than random chance for both the PCC and MCC (P < 0.001, Table S3). These characters showed no to little overlap across the five species (Fig. 1). The first (DF1), the second (DF2), the second (DF2),

the third (DF3) and the fourth (DF4) discriminant function explained 86.6%, 11.5%, 1.3% and 0.6% of the total variation, respectively (Table 2, Fig. S1). Among the morphological characters used, the presence and the type of the indumentums are the most informative characters for species distinction in P. subsect. *Saxifragae*. The highest discriminative function (DF1) is primarily influenced by the presence of eglandular hairs at the middle part of the stem (a9). The presence of glandular hairs on calyx (a34) and eglandular hairs at the lower part of the stem (a8) have also high discriminant validity in DF1 (Table 2). The presence of eglandular hairs on calyx (a35) and glandular hairs at the middle/upper part of the stem (a6/a7) are the characters loading most heavily on DF2 and DF3, respectively (Table 2). The five Petrorhagia species included in the analyses are clearly distinct from one another (Table S3, Fig. 1, S1, S2).



Fig. 1. Stepwise discriminant analysis plot for the five *Petrorhagia* species. The squares indicate the function group centroid of each species.

Discussion

Multivariate morphometrics provide a powerful mean in order to assess the variation patterns at various taxonomic levels and is especially useful in disentangling the boundaries between closely related and/or poorly differentiated taxa (e.g., Kougioumoutzis & al. 2015), as well as in unveiling the most informative and discriminating characters that differentiate among them (Marhold 2011). In the present study, the application of multivariate morphometrics resulted in important conclusions with taxonomic consequences. More specifically, the results of morphometric analyses enabled the delimitation of five well-separated species and the elucidation of the identity of the deviating *Petrorhagia* populations from SE Peloponnese: the latter constitute a new, distinct and rather rare taxon, herein described as *P. laconica*.

Petrorhagia subsect. *Saxifragae* is a taxonomically complicated group with several overlapping characters. Our morphometric analyses, however, showed that all species are clearly distinguished from each other. Their distinction can be achieved even without using morphological seed characters, an important diagnostic feature in the genus *Petrorhagia*. The latter were not included in the morphometric analyses, as most specimens used were not in fruit.

The presence and type of hairs on stem and calyx have emerged as important diagnostic characters. The presTable 2. Stepwise discriminant analysis results for the five Petrorhagia species included in the present study based on 14 morphological characters, as well as the eigenvalues of the discriminant functions and the proportion of variance (PV) explained by each discriminant function. DF1, DF2, DF3 and DF4 indicate the first, the second, the third and the fourth discriminant function, respectively. F, T and WD indicate the F-test P values, the tolerance values, as well as the Wilk's lambda values of the discriminant functions and characters retained in the stepwise discriminant analysis, respectively. SML indicates the loadings from the stepwise discriminant analysis (i.e. the correlation of the characters with the discriminant functions). Character abbreviations follow Table 1. Higher values are shown in bold.

Character	Т	F	WD	DF1	DF2	DF3	DF4
Eigenvalue	_	_	_	205.23	27.22	3.12	1.38
PV	-	-	_	86.6	11.5	1.3	0.6
WD	-	_	_	0.00	0.00	0.10	0.42
					SML		
a9	0.85	0.01	0.00	0.35	0.09	-0.06	0.11
a34	0.42	0.00	0.00	0.26	0.06	0.07	0.16
a15	0.58	0.00	0.00	-0.06	0.22	0.03	0.09
a35	0.51	0.00	0.00	-0.06	0.23	0.69	-0.23
a31	0.09	0.00	0.00	0.06	-0.03	-0.36	0.02
a8	0.45	0.00	0.00	0.20	0.03	0.34	0.32
a29	0.08	0.00	0.00	-0.09	-0.13	-0.29	0.03
a28	0.67	0.01	0.00	-0.06	-0.16	0.25	-0.14
a25	0.60	0.03	0.00	-0.07	-0.15	0.24	-0.02
a7	0.28	0.00	0.00	-0.05	0.18	-0.19	0.53
a6	0.16	0.00	0.00	-0.05	0.18	-0.19	0.53
a5	0.24	0.00	0.00	-0.09	0.38	-0.21	0.44
a14	0.38	0.00	0.00	-0.13	0.00	-0.08	0.37
a4	0.69	0.03	0.00	-0.15	0.00	0.02	0.28
a27	0.23	0.00	0.00	-0.05	-0.04	0.11	-0.26
a30	0.60	0.03	0.00	0.03	0.03	-0.04	-0.11

ence of eglandular hairy stems throughout their length is a unique feature of *Petrorhagia laconica*. *Petrorhagia fasciculata* and *P. phthiotica* are the only species with an indumentum of glandular hairs on stems, while the stems of *P. graminea* are covered by a mixture of glandular and eglandular hairs, at least at base. All examined specimens of *P. saxifraga* had glabrous stems. Furthermore, *P. laconica* is the only species with a glandular hairy calyx. The calyx of *P. graminea* is predominantly covered by eglandular hairs, often mixed with few glandular hairs. All examined specimens of *P. fasciculata* had eglandular hairy calyces, while all calyces in *P. phthiotica* and *P. saxifraga* were glabrous.

Length of the inflorescence, number of lateral branches and flowers per flowering stem, width of lower leaves and the ratio pedicel/calyx length further differentiate *Petrorhagia laconica* from all other species of *P*. sect. *Saxifragae* (Table S2, Fig. S1). Moreover, *P. laconica*, together with *P. saxifraga*, are the only species with calyces enclosed by epicalyx bracts; in the specimens of all other species (including *P. fasciculata*) there are no bracts enclosing the calyx. The small petals and the large number of flowers per terminal inflorescence unit easily distinguish *P. fasciculata* from all other species. *Petrorhagia graminea* and *P. saxifraga* are the only species with 3-veined sepals and they also have the largest petals. The 3-veined leaves and the short stems further differentiate *P. phthiotica* from all other members of *P.* sect. *Saxifragae*.

The results of our analyses have revealed a diagnostic character not previously used in the genus *Petrorhagia*. Although the length of leaves is similar in all species examined, leaf width varies among species. *Petrorhagia laconica* and *P. saxifraga* have narrow basal leaves compared to the other species. Furthermore, cauline leaves are diminishing in size toward the apex in all species, but diminishing in leaf length only, while leaf width remains stable, has been observed only in *P. laconica* (Table S2, Fig. S1).

Taxonomy

Petrorhagia laconica Trigas, Kalpoutz. & Kougioum., **sp. nov.** – Fig. 2.

Holotype: Greece, Peloponnisos, sand dunes c. 3.5 km NW of Neapoli village, 36°31.721'N, 23°01.273'E, 2 m, 22 May 2017, *Trigas 6311* (ACA; isotypes: B, UPA).

Diagnosis — Related to *Petrorhagia saxifraga*, from which it differs in the following morphological characters: underground stock rooting at nodes; stems procumbent to ascending or erect, glaucous-green, simple or rarely with few (1–3[–6]) lateral branches at upper ¹/₃, minutely papillose-scabridulous except glabrous at 1–4 upper internodes; leaves closely appressed to stem, rigid; flowers solitary or rarely paired, 1(or 2) per flowering stem, rarely up to 6; calyx minutely glandular pubescent; ribs purplish brown, obscurely 1-veined; capsule 2.5–3 mm long, sparsely tuberculate at proximal ¹/₂; seeds with flat, non-thickened margin.

Description — Plants perennial, with branched, slender, woody *underground stock* rooting at nodes, producing few to several flowering stems. *Flowering stems* procumbent to ascending or erect, glaucous-green to purplish brown, 7–40 cm long, simple or rarely with few (1-3[-6]) lateral branches at upper $\frac{1}{3}$, minutely papillose-scabridulous ex-



Fig. 2. *Petrorhagia laconica* – A: sand-dune habitat NW of Neapoli village in SE Peloponnisos with evergreen shrubs (locus classicus), 22 May 2017; B: individual with simple, procumbent-ascending stems at the locus classicus, 16 May 2014; C: middle part of stem with leaves; D: upper part of stem with flower; E: flower, lateral view; F: flower, apical view; G: immature capsule; H: seed, dorsal face; I: seed, ventral face. – Scale bars: C–F: 3 mm; G: 1 mm; H, I: 500 µm. – All photographs by Panayiotis Trigas except G by Eleftherios Kalpoutzakis.

cept glabrous at 1-4 upper internodes; internodes markedly unequal, larger at middle of stem, diminishing in size toward base and apex. Leaves linear-subulate, 1/3-11/4 as long as internodes, rigid, 1-veined, united at base into a sheath 1–1.5 mm long, with a purplish brown ring at base, apex acuminate to caudate; *lower leaves* $7-19 \times$ 0.5-1 mm, minutely roughly papillose at margin, otherwise glabrous; cauline leaves closely appressed to stem, $3-16 \times 0.5-1$ mm, glabrous. *Flowers* solitary or rarely paired at apices of stems and branchlets, 1(or 2) per flowering stem, rarely up to 6, subtended by 4 straw-coloured, lanceolate, hyaline, glabrous epicalyx bracts, sometimes purplish-tinged at midrib. Calyx cylindric, 3-6 mm long, minutely glandular pubescent; ribs purplish brown, obscurely 1-veined; teeth oblong-triangular, 0.5-1 mm long, apex obtuse. Petals white, $4-8 \times 1.5-3$ mm, abaxial surface purple-veined, adaxial surface with three longitudinal purplish stripes at base, with glabrous claw not distinctly delimited from emarginate lamina. Anthers white. Ovary oblong-ovoid, c. 1 mm long; anthophore green, c. 0.5 mm long, glabrous. Capsule pale brown, ovoid, 2.5–3 mm long, smooth or sparsely tuberculate at proximal ¹/2. Seeds blackish brown, ovate-oblong, 1-1.2 \times 0.7–0.8 mm, both surfaces reticulate, margin flat, not thickened, apex long mucronate.

Distribution — *Petrorhagia laconica* is distributed in the Malea Peninsula in SE Peloponnisos. Two populations have been discovered so far; the first one is located at the Gulf of Neapoli, right across Elafonisos Island, while the second one was found close to Sykea village, c. 25 km N-NW of the first population. It is the southernmost distributed species of *P*. subsect. *Saxifragae* in Europe.

Ecology — Petrorhagia laconica is known to grow in two different habitats. In Neapoli, all individuals were observed to grow in the inner sand dune zone, usually surrounded by sclerophyllous shrubs, such as Anthyllis hermanniae L., Ceratonia siliqua L., Pistacia lentiscus L., Pyrus spinosa Forssk. and Smilax aspera L. Petrorhagia laconica grows in stabilized sand dunes among the shrubs, together with other sand dune adapted taxa, such as Anthemis tomentosa L., Asphodelus ramosus L., Centaurea sonchifolia L., Elytrigia juncea (L.) Nevski, Linaria tenuis (Viv.) Spreng. and Scirpoides holoschoenus (L.) Soják.

Close to Sykea village, *Petrorhagia laconica* grows in small remnant patches of macchie and phrygana formations that form a mosaic among the abundant olive groves. The population occupies a small area at 120–140 m a.s.l., on flysch. Shrub species composition of macchie is similar to that of the locus classicus, with *Lavandula stoechas* L., *Olea europaea* L., *Phlomis fruticosa* L., *Pistacia lentiscus* and *Pyrus spinosa* the predominant species. *Petrorhagia laconica* usually grows in small patches among the shrubs, together with *Ballota acetabulosa* (L.) Benth., *Cistus salviifolius* L., *Globular*- *ia alypum* L., *Hypericum triquetrifolium* Turra, *Sarcopoterium spinosum* (L.) Spach, *Silene nocturna* L., etc.

Conservation status — In Neapoli Lakonias, *Petrorhagia laconica* grows on a large sandy beach c. 3.5 km long, between Neapoli and Viglafia village. The width of sand dunes along this beach ranges from 30 to 150 m, and the species is exclusively confined to the inner sand dune zone, where it is fairly rare. It usually forms small groups of 5–30 individuals, but single remote plants were also observed. The total population of Neapoli is estimated to include 300–400 individuals, mainly concentrated at the E part of the beach. The Sykea population counts c. 500 individuals distributed in an area of c. 0.1 km².

Habitat loss and touristic development are the main threats that *Petrorhagia laconica* faces at its locus classicus. Sand dunes have been partly turned into cultivated fields at the E half of the beach, almost certainly leading to a significant population decline. The habitat also hosts thousands of swimmers during summer, being an unpredictable risk factor for its long term survival. The whole area, however, belongs to the Natura 2000 network of protected areas (Periochi Neapolis kai Nisos Elafonisos, GR2540002), and numerous human activities that could create additional risks for the population of *P. laconica* in Neapoli area are fortunately forbidden.

Habitat loss is also the main threat that *Petrorhagia laconica* faces in Sykea area, as more than 50% of the area previously covered with natural vegetation have been turned into cultivated fields. As a result, it is irrefutable that the species has suffered a severe population decline in the near past, which can be estimated as probably approaching the percentage of habitat loss.

The flora of SE Peloponnisos has been intensively explored during the last decades (e.g. Kalpoutzakis & Constantinidis 2005, 2006; Greuter 2012; Kalpoutzakis & al. 2012). The recent discovery of Petrorhagia laconica indicates that the species should not be common in this area. The habitat types in which it grows (especially the mosaic with macchie and phrygana formations), however, are abundant in SE Peloponnisos. Consequently, the existence of additional populations of this tiny and easily overlooked species cannot be ruled out. The extent of occurrence (EOO) does not exceed 25 km² and since *P. laconica* is a very local species, its area of occupancy (AOO) is much smaller, apparently less than 1 km². Ergo, due to: (1) the restricted EOO and AOO, (2) the current existence of max. 900 mature individuals distributed in two populations and (3) the high possibility that the extant localities and populations could be eradicated as a result of human interference, P. laconica is assigned to the Endangered (EN) IUCN (2001) category, following criteria B1ab(i,ii,iii,iv)+2ab(i,ii,iii,iv). Petrorhagia laconica should be carefully monitored, cultivated ex situ in botanical gardens and special conservation measures be taken to safeguard its populations.

Character	P. laconica	P. saxifraga	P. graminea
Underground stock	rooting at nodes	not rooting at nodes	not rooting at nodes
Stems	procumbent to ascending or erect, glaucous-green, simple or rarely with few (1–3[–6]) lateral branches at upper ¹ / ₃ , minutely papillose- scabridulous except glabrous at 1–4 upper internodes	arcuately ascending to erect, green, usually branched to much-branched, glabrous to shortly papillose or scabrid-pubescent at least in lower part	ascending to erect, green, sparingly branched, glabrous to densely pubescent
Leaves	closely appressed to stem, linear- subulate, rigid	± spreading, linear to linear- lanceolate, herbaceous	± spreading, linear, herbaceous
Inflorescence	flowers solitary or rarely paired, 1(or 2) per flowering stem, rarely up to 6	flowers solitary or in fascicles, 2–59 per flowering stem	flowers solitary or in fascicles of 2 or 3, 2–37 per flowering stem
Epicalyx bracts	present	present	absent
Calyx	minutely glandular pubescent; ribs purplish brown, obscurely 1-veined	glabrous or occasionally sparsely eglandular pubescent; ribs green, 3-veined	densely eglandular pubescent to glabrous; ribs green, 3-veined
Capsule	2.5–3 mm long, smooth or sparsely tuberculate at proximal ½	3.5–5 mm long, smooth	3–5 mm long, smooth
Seeds	ovate-oblong, 1–1.2 × 0.7–0.8 mm, margin flat, not thickened	ovate, 0.9–1.6 × 0.6–1.1 mm, margin curved, thickened	oblong to suborbicular, 0.8–1.5 × 0.7–1 mm, margin ± curved, slightly thickened
Flowering period	May-August	May–August	September–January

Table 3. Main morphological differences among Petrorhagia laconica, P. saxifraga and P. graminea.

Etymology — The specific epithet refers to Laconia (Lakonia, also known as Lacedaemonia), a region that covers almost the entire SE part of the Peloponnisos, where the new species was collected. This name goes back to the historical times, when Laconia was the principal region of the Spartan state.

Taxonomic relationships — Petrorhagia laconica undoubtedly belongs to P. sect. Petrorhagia, as it is indicated by its 1-veined leaves, the presence of epicalyx bracts, the not abruptly clawed petals, the reticulate, blackishbrown seeds and the perennial life-cycle. The thickened seed margin is not a common feature of all species in P. sect. Petrorhagia, as it is indicated by Ball & Heywood (1964). Petrorhagia dianthoides (Sm.) P. W. Ball & Heywood, P. fasciculata, P. grandiflora Iatrou and P. thessala (Boiss.) P. W. Ball & Heywood have a flat, thin seed margin, similar to that of P. laconica.

Within *Petrorhagia* sect. *Petrorhagia*, *P. laconica* shows close affinities to the members of *P.* subsect. *Saxi-fragae*. The solitary or paired flowers and the lanceolate, 1-veined epicalyx bracts advocate for the inclusion of *P. laconica* within *P.* subsect. *Saxifragae*. Flower and seed morphological characters further support this taxonomic view. The simple or sparingly branched stems, however, resemble the members of *P. subsect. Thessalae*, which otherwise have significant morphological differences from *P. laconica* (e.g. capitate inflorescence, broadly ovate or sub-orbicular bracts, larger and different petals, etc.).

Petrorhagia laconica appears to be one of the most distinct species in *P*. sect. *Petrorhagia*. The results of multivariate morphometric analyses (Tables 2, S3, Fig. 1, S1, S2) support its distinct taxonomic position with-

in P. subsect. Saxifragae. The closest related species seem to be P. saxifraga and P. graminea, which, however, show significant morphological differences from P. laconica (see also Table 3). The possibility of vegetative reproduction (underground stolons connecting different individuals have been observed in the field) and the tuberculate capsule have not been recorded in other Greek Petrorhagia species, and they probably represent unique features within the whole genus. The discovery of P. laconica is in line with Trigas & al. (2007, 2012), which state that SE Peloponnese is one of the important regions in Greece in terms of endemic plant species richness and conservation, as well as a significant diversity centre for Petrorhagia, because more than 30 % of the species diversity of the whole genus is hosted in this small region.

Additional specimens examined — GREECE: Peloponnisos, c. 3.5 km S-SE of Sykea village, 36°43.970'N, 22°57.079'E, 130 m, macchie and phrygana among olive groves, on flysch, 6 May 2013, *Kalpoutzakis 4294* (ACA); ibid., 29 Jul 2017, *Kalpoutzakis 4959* (ACA); c. 2.5 km S-SE of Sykea village, 36°44.552'N, 22°57.209'E, 120 m, macchie and phrygana among olive groves, on flysch, 22 May 2017, *Trigas 6312* (ACA).

Acknowledgements

We would like to thank Prof. Emer. Arne Strid and an anonymous reviewer for insightful comments that greatly improved the manuscript. We are grateful to the curators of UPA and ATH for loaning specimens for our study. We also wish to thank Dr Aimilia-Eleni Nikolopoulou (Agricultural University of Athens) for collecting specimens of *Petrorhagia laconica* with mature capsules.

References

- Ball P. W. & Heywood V. H. 1964: A revision of the genus *Petrorhagia*. – Bull. Brit. Mus. (Nat. Hist.), Bot. 3: 119–172.
- Bittrich V. 1993: Caryophyllaceae. Pp. 206–236 in: Kubitzki K., Rohwer J. G. & Bittrich V. (ed.), Families and genera of vascular plants 2. – Berlin, Heidelberg & New York: Springer Verlag.
- Coode M. J. E. & Cullen J. 1967: *Petrorhagia* (Ser.) Link. – Pp. 131–135 in: Davis P. H. (ed.), Flora of Turkey and the East Aegean Islands 2. – Edinburgh: University Press.
- Dillenberger M. S. & Kadereit J. W. 2014: Maximum polyphyly: multiple origins and delimitation with plesiomorphic characters require a new circumscription of *Minuartia (Caryophyllaceae)*. Taxon **63**: 64–88.
- Dimopoulos P., Raus Th., Bergmeier E., Constantinidis Th., Iatrou G., Kokkini S., Strid A. & Tzanoudakis D. 2013: Vascular plants of Greece: an annotated checklist. – Berlin: Botanic Garden and Botanical Museum Berlin-Dahlem; Athens: Hellenic Botanical Society. [Englera 31].
- Fior S., Karis P. O., Casazza G., Minuto L. & Sala F. 2006: Molecular phylogeny of the *Caryophyllaceae* (*Caryophyllales*) inferred from chloroplast *matK* and nuclear rDNA ITS sequences. Am. J. Bot. **93**: 399–411.
- Georgiou O. (1997) *Petrorhagia* (Ser.) Link. Pp. 333–343 in: Strid A. & Tan K. (ed.) Flora hellenica
 1. Königstein: Koeltz Scientific Books.
- Greenberg A. K. & Donoghue M. J. 2011: Molecular systematics and character evolution in *Caryophyllaceae*. – Taxon **60**: 1637–1652.
- Greuter W. 2012: Results of the Seventh "Iter Mediterraneum" in the Peloponnese, Greece, May to June 1995. (Occasional Papers from the Herbarium Greuter $- N^{\circ}$ 1). - Bocconea **25:** 5–127.
- Harbaugh D. T., Nepokroeff M., Rabeler R. K., McNeill J., Zimmer E. A. & Wagner W. L. 2010: A new lineage-based tribal classification of the family *Caryophyllaceae*. – Int. J. Plant Sci. **171**: 185–198.
- Hernández-Ledesma P., Berendsohn W. G., Borsch Th., Mering S. von, Akhani H., Arias S., Castañeda-Noa I., Eggli U., Eriksson R., Flores-Olvera H., Fuentes-Bazán S., Kadereit G., Klak C., Korotkova N., Nyffeler R., Ocampo G., Ochoterena H., Oxelman B., Rabeler R. K., Sanchez A., Schlumpberger B. O. & Uotila P. 2015: A taxonomic backbone for the global synthesis of species diversity in the angiosperm order *Caryophyllales.* – Willdenowia 45: 281–383.

- Hilooğlu M., Poyraz I. E., Poyraz I., Ataşlar E. & Sözen E. 2016: Genetic relationships among some Turkish *Petrorhagia* (Ser.) Link (*Caryophyllaceae*) taxa using ISSR markers. – Phytotaxa 272: 165–172.
- Huberty C. J. & Olejnik S. 2006: Applied MANOVA and Discriminant Analysis, 2nd ed. Hoboken: Wiley and Sons.
- Iatrou G. 1985: Petrorhagia grandiflora sp. nov. (Caryophyllaceae) from Greece. – Nord. J. Bot. 5: 441–445.
- Kalpoutzakis E. & Constantinidis T. 2005: New data on the distribution of endemic and rare taxa in the flora of east Peloponnisos, Greece. – Bot. Chron. 18(2): 115–136.
- Kalpoutzakis E. & Constantinidis T. 2006: Additions and annotations to the flora of Peloponnisos (Greece). – Willdenowia 36 (Special Issue): 271–284.
- Kalpoutzakis E., Trigas P. & Constantinidis T. 2012: Allium orestis sp. nov. (Amaryllidaceae) from Parnon and Taigetos mountains, south Peloponnisos, Greece. – Nord. J. Bot. **30**: 195–200.
- Kougioumoutzis K., Kalpoutzakis E. & Constantinidis T. 2017: Multivariate morphometric analysis and taxa delimitation in two narrow Greek endemics: Astragalus maniaticus and Aethionema saxatile subsp. corinthiacum. – Pl. Biosyst. 151: 108–116.
- Marhold K. 2011: Multivariate morphometrics and its application to monography at specific and infraspecific levels. – In: Stuessy T. F. & Lack H. W. (ed.), Monographic Plant Systematics: fundamental assessment of plant biodiversity. – Ruggell: Gantner.
- Naimi B., Hamm N. A., Groen T. A., Skidmore A. K. & Toxopeus A. G. 2014: Where is positional uncertainty a problem for species distribution modelling? – Ecography 37: 191–203.
- Pirani A., Zarre S., Pfeil B. E., Bertrand Y., Assadi M. & Oxelman B. 2014: Molecular phylogeny of *Acanthophyllum (Caryophyllaceae, Caryophylleae)*, with emphasis on subgeneric classification. – Taxon 63: 592–607.
- R Core Team 2015: R: A language and environment for statistical computing. R Foundation for Statistical Computing, Vienna. – Published at https://www.Rproject.org/ [accessed 3 Jul 2017].
- Rabeler R. K. & Hartman R. L. 2005: *Caryophyllaceae*.
 Pp. 3–8 in: Flora of North America Editorial Committee (ed.), Flora of North America north of Mexico
 5. Oxford: University Press.
- Španiel S., Zozomová-Lihová J. & Marhold K. 2017: Revised taxonomic treatment of the *Alyssum monta-num – A. repens* complex in the Balkans: a multivariate morphometric analysis. – Plant Syst. Evol. **303**: 1413–1442.
- Thiers B. 2017+ [continuously updated]: Index Herbariorum: A global directory of public herbaria and associated staff. New York Botanical Garden's Virtual Herbarium. Published at http://sweetgum.nybg.org/ science/ih/ [accessed 3 Aug 2017].

- Trigas P., Iatrou G. & Karetsos G. 2007: Species diversity, endemism and conservation of the family *Caryophyllaceae* in Greece. – Biodivers. Conserv. **16**: 357–376.
- Trigas P., Tsiftsis S., Tsiripidis I. & Iatrou G. 2012: Distribution patterns and conservation perspectives of

the endemic flora of Peloponnese (Greece). – Folia Geobot. **47:** 421–439.

Wickham H. 2009: ggplot2: Elegant Graphics for Data Analysis. – New York: Springer.

Willdenowia

Open-access online edition www.bioone.org/loi/will Reference BioOne COMPLETE Online ISSN 1868-6397 · Print ISSN 0511-9618 · Impact factor 0.680 Published by the Botanic Garden and Botanical Museum Berlin, Freie Universität Berlin © 2018 The Authors · This open-access article is distributed under the CC BY 4.0 licence