

Notes on the genus Argentochiloides Błeszyński, 1961, with description of a new species (Lepidoptera: Pyraloidea, Crambidae)

Author: Bassi, Graziano

Source: Revue suisse de Zoologie, 130(2): 291-295

Published By: Muséum d'histoire naturelle, Genève

URL: https://doi.org/10.35929/RSZ.0103

The BioOne Digital Library (https://bioone.org/) provides worldwide distribution for more than 580 journals and eBooks from BioOne's community of over 150 nonprofit societies, research institutions, and university presses in the biological, ecological, and environmental sciences. The BioOne Digital Library encompasses the flagship aggregation BioOne Complete (https://bioone.org/subscribe), the BioOne Complete Archive (https://bioone.org/archive), and the BioOne eBooks program offerings ESA eBook Collection (https://bioone.org/esa-ebooks) and CSIRO Publishing BioSelect Collection (https://bioone.org/csiro-ebooks).

Your use of this PDF, the BioOne Digital Library, and all posted and associated content indicates your acceptance of BioOne's Terms of Use, available at www.bioone.org/terms-of-use.

Usage of BioOne Digital Library content is strictly limited to personal, educational, and non-commmercial use. Commercial inquiries or rights and permissions requests should be directed to the individual publisher as copyright holder.

BioOne is an innovative nonprofit that sees sustainable scholarly publishing as an inherently collaborative enterprise connecting authors, nonprofit publishers, academic institutions, research libraries, and research funders in the common goal of maximizing access to critical research.

Notes on the genus *Argentochiloides* Bleszyński, 1961, with description of a new species (Lepidoptera: Pyraloidea, Crambidae)

Graziano Bassi

Via Sant'Agostino, 51, I-10051 Avigliana (Torino), Italy; Muséum d'histoire naturelle de Genève, C.P. 6434, CH-1211 Genève 6, Suisse (Corresponding member); alphacrambus@gmail.com; https://orcid.org/0000-0002-6028-0740

Abstract: A new species of *Argentochiloides* Błeszyński is described and illustrated: *Argentochiloides mirabilis* **sp. nov.** from the Republic of South Africa. A list of the known species and an image of the holotype of each species are given. *Charltona argyrastis* Hampson is transferred from *Aurotalis* Błeszyński to *Argentochiloides*.

Keywords: Afrotropical - Ancylolomiini - Crambinae - distribution - new combination.

INTRODUCTION

The genus *Argentochiloides* was erected by Błeszyński (1961: 36, 37) for *A. xanthodorsellus* Błeszyński from Tanzania. A new South African species was added by Bassi (1999). A small series of another South African species here described allows me to update the actual composition of this genus, with the addition of *A. argyrastis* (Hampson, 1919) comb. n., originally described in *Charltona* Swinhoe, 1886 and here transferred from *Aurotalis* Błeszyński, 1970.

MATERIAL AND METHODS

Genitalia preparations were made following Robinson (1976). The terminology of the genitalia follows Błeszyński (1965), Klots (1970) and Kristensen (2003). Genitalia photographs were taken with a Canon S120 digital camera. The habitus photos were made with a Nikon D3300 digital camera. The images were enhanced with Adobe Photoshop Elements. The following abbreviations are used: 'GB' for Graziano Bassi, 'GS' for genitalia slides, 'RCGB' for Graziano Bassi research Collection, Avigliana, Italy (to be deposited in Muséum d'histoire naturelle de Genève, Switzerland), and 'TMSA' for Ditsong National Museum of Natural History (formerly the Transvaal Museum), Pretoria, Republic of South Africa.

TAXONOMY

Argentochiloides Błeszyński, 1961

Argentochiloides Błeszyński, 1961: 36, 37, type species Argentochiloides xanthodorsellus Błeszyński, 1961, by original designation.

Diagnosis: The adults differ from the other Afrotropical Crambinae in having a medium-large size (22-33 mm) and wings, especially the forewings, of a particularly reflective glossy white. The male genitalia are characterized by the heavily curved uncus and gnathos associated with strongly asymmetrical valvae and a long, sinuous phallus, characters which are not present all together in other genera of Afrotropical Crambinae. In the female genitalia the most evident feature is the distal large globular extension of the ductus bursae, associated with papillae anales with strong setae originating from short projections as in Aurotalis Błeszyński, 1970. However, in Aurotalis the ductus bursae is simple. In Ancylogastra gangraensis Bassi, Sáfián, Müller & Kravchenko (2021), the only other Afrotropical species which has a globular extension of the ductus bursae, the extension is proximal in the ductus bursae, and in both Aurotalis and Ancylogastra the male genitalia have symmetrical valvae. The Oriental genus Pseudoclasseya Błeszyński, 1964 has asymmetrical valvae and a globular extension of the ductus bursae also, but the male genitalia lack the gnathos, the asymmetry in the valvae is restricted to the costal process, the phallus is not differentiated, and the

Manuscript accepted 06.06.2023 DOI: 10.35929/RSZ.0103

This is an open access article distributed under the terms of the Creative Commons Attribution License (CC BY 4.0), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited (see https://creativecommons.org/licenses/by/4.0/).

292 G. Bassi

female genitalia (Bassi, 1989, figs 22-25) have papillae anales of the *Calamotropha-Pseudocatharylla* type, the globular extension has its own short ductus, and a signum is present in the bursa copulatrix.

Remarks: Błeszyński (1961: 36) placed the genus next to *Chilo* Zincken in the original description, but in later years he realized that this association was incorrect and included the genus in the *Pseudocatharylla* complex (Błeszyński, 1964: 755), despite having doubts on its position. In the modified uncus and gnathos in the male genitalia, the papillae anales and general structure of the female genitalia and the tympanal organs (Fig. 9) *Argentochiloides* is a member of the Ancylolomiini Ragonot (1889), and of the *Ancylolomia* complex of genera (Bassi, 2013), close to *Aurotalis* and *Ancylogastra* Bassi & Poltavsky (in Bassi *et al.*, 2021). Unfortunately, all specimens are too old to allow for successful standard DNA extraction; the live specimen photographed in 2010 (Figs 5, 6) was not collected.

Argentochiloides mirabilis sp. nov. Figs 1, 5, 6, 7, 9

Holotype: female, [South Africa, Free State, 31°56'S 22°08'E] Modderpoort, 18.xii. [19]25, A.J.T. Janse, TMSA.

Paratypes: 2 females, with same data as holotype, GS 3788 GB, TMSA and CB; 2 females, Buffelspoort, 15.xii. [19]24, A.J.T. Janse, TMSA and RCGB.

Additional material: 1 male, Gauteng, Pretoria, Tshwane Metropolitan Municipality, 1340 m, 25°34'31"S, 28°34'40"E, 9.xii.2020, specimen only photographed by J.-M. Van Tonder.

Etymology: Derived from *mirabilis* (Latin) = wonderful, referring to the striking habitus of the species.

Diagnosis: No other species of Afrotropical Crambinae have a forewing pattern in any way similar to that of this species in the silvery white forewings with three large olive-green stripes. In female genitalia *A. mirabilis* (Fig. 7) is most similar to *A. argyrastis* (Fig. 8) but its apophyses posteriores are longer, narrower and sinuous instead of straight, and the ductus bursae has two sclerotised patches basally and distally, absent in *A. argyrastis*.

Description: (Figs 1, 5, 6). Wingspan 33 mm. Labial palpi 3.5 X eye diameter, yellow sprinkled with brown, with inner side white. Maxillary palpi basally olive brown, then yellow. Antenna simple, brown with costa olive green. Frons yellow sprinkled with brown and bordered white. Ocelli and chaetosemata poorly developed. Vertex white. Patagium yellow bordered white. Tegulae olive green bordered white. Thorax

white sprinkled with pale brown. Forewing with pointed apex and termen oblique; ground colour silvery white with three olive green stripes, the medial one enlarging distally; costa basally brown, then silvery white; fringes silvery white; underside with large white border and medially brown. Hindwing silvery white with costa and medial fascia pale bronze brown; fringes pure white; underside white with costa yellow-brown and veins M2-Cu2 brown; frenulum triple. Fore and mid legs bronze brown with inner side pale yellow; hind leg paler with tibial spurs short, inner spurs as long as outer. Abdomen olive green with first tergite white.

Female genitalia (Fig. 7). Papillae anales stout, ventrally rounded; apophyses posteriores 1.6 times as long as apophyses anteriores, curved inward; abdominal segment VIII large, ventrally membranous; ostium bursae asymmetrical; ductus bursae with two more sclerotised patches basally and distally; globular extension strongly wrinkled; corpus bursae 1.5 longer than ductus bursae, suboval

Male. Judging from the photographs (Figs 5, 6), similar to female, with antenna slightly serrate.

Biology: Unknown. Specimens were caught in December.

Distribution: South Africa: Free State, Gauteng and North West Provinces.


Remarks: All of the specimens on which the species of the genus are based are very old and the suspicion that the species were extinct was strong, when, surprisingly, a live male of *A. mirabilis* was photographed in December 2020 in the surroundings of Pretoria. This specimen was not collected, which prevents insights into the taxonomy and systematics of the species and genus, but perhaps it is for the best and this specimen demonstrates how much life can survive the toughest trials.

Argentochiloides argyrastis (Hampson, 1919) comb. n. Figs 2, 8

Charltona argyrastis Hampson, 1919: 306, 307. Aurotalis argyrastis (Hampson, 1919). – Bassi, 2016: 14.


Type locality: Kenya

Remarks: The female holotype from western Kenya and an additional specimen from Nigeria were both collected early in the past century. They remain the only two known specimens and the Nigerian one is without abdomen. The holotype is figured on Figs 2, 8.



Figs 1-6. Argentochiloides spp., habiti. (1) A. mirabilis sp. n., holotype, wingspan 32 mm. (2) A. argyrastis (Hampson), holotype and its labels, photo A. Giusti, © Courtesy of the Trustees of the Natural History Museum, London, U.K. (3) A. xanthodorsellus Błeszyński, holotype and its labels, wingspan 22 mm, photo: H. Bruckner, Lepidoptera Image Collection, Natural History Museum Vienna, Austria. (4) A. meridionalis Bassi, holotype, Johannesburg, Liniord, wingspan 28 mm. (5, 6) A. mirabilis sp. n., male, Pretoria, Tshwane (photo Jaunne-Marelize Van Tonder, Dec. 9, 2020).

294 G. Bassi



Figs 7-9. Argentochiloides spp., female genitalia and tympanal organs. (7) A. mirabilis sp. n., holotype. (8) A. argyrastis, holotype. (9) A. mirabilis sp. n., tympanal organs.

Argentochiloides xanthodorsellus Błeszyński, 1961 Fig. 3

Type locality: Tanzania.

Remarks: Described from four males collected in the 1930s, the species is only known from the Matengo Highlands in southern Tanzania. No other specimens are known beyond the typical series. The holotype is figured on Fig. 3.

Argentochiloides meridionalis Bassi, 1999 Fig. 4

Type locality: South Africa.

Remarks: Described from two males collected early in the past century in areas now integrated into the city of Johannesburg. No other specimens are known beyond the type series. The holotype is figured on Fig. 4.

ACKNOWLEDGMENTS

I thank J.-M. Van Tonder, City of Tshwane Metropolitan Municipality (Pretoria), South Africa, for providing the pictures of a live adult of *Argentochiloides mirabilis*, S. Gaal-Haszler and H. Bruckner, Naturhistorisches Museum, Wien, Austria, for providing a photo of the holotype of *A. xanthodorsellus*, and A. Giusti and K. Tuck, Natural History Museum, London, U. K., for sending images of the type of *A. argyrastis*. A warm memory goes to the late M. Krüger for his great help during my study of the TMSA material.

REFERENCES

- Bassi G. 1989. Contributions to the study of the Crambinae (Lepidoptera, Pyralidae), III: Notes on the genus *Pseudoclasseya* Bleszynski. *Bollettino del Museo Regionale di Scienze Naturali Torino* 7(2): 373-382.
- Bassi G. 1999. Contributo allo studio delle Crambinae: X. Nuove specie africane (Lepidoptera: Pyraloidea, Crambinae). Shilap. Revista de Lepidopterologia 27(105): 57-67.
- Bassi G. 2013. Revisione delle specie afrotropicali del genere *Ancylolomia* Hübner, [1825] (Lepidoptera: Pyralidae, Crambinae). I: i gruppi *indica* e *chrysargyria*. *Shilap*. *Revista de Lepidopterologia* 41(164): 517-529.

- Bassi G. 2016. Studies on Afrotropical Crambinae (Lepidoptera, Pyraloidea, Crambidae): Notes on the genus Aurotalis Błeszyński, 1970. Revue suisse de Zoologie 123 (1): 11-20. DOI: 10.5281/zenodo.46283
- Bassi G., Sáfián S., Léger T., Müller G., Kravchenko V., Poltavsky A. 2021. Ancylogastra, a new genus of Afrotropical Crambinae, with descriptions of seven new species (Lepidoptera, Pyraloidea, Crambidae). Zootaxa 5052 (1): 042-060. DOI: 10.11646/zootaxa.5052.1.2.
- Błeszyński S. 1961. Studien über die Crambidae (Lepidoptera).
 Teil XXXI. Bemerkungen über einige Arten mit Beschreibungen von zwei neuen Gattungen und einer neuen Art.
 Zeitschrift der Wiener entomologischen Gesellschaft 46:
 33-38.
- Błeszyński S. 1964. Revision of the world species of the family Crambidae (Lepidoptera). Part 2. Genera: *Pseudocatha-rylla* Blesz., *Classeya* Blesz., *Pseudoclasseya* Blesz. and *Argentochiloides* Blesz. *Acta Zoologica Cracoviensia* 9(11): 683-760.
- Błeszyński S 1965. Crambinae. In: Amsel H.G., Gregor F., Reisser H. (Eds). Microlepidoptera Palaearctica 1(1-2). Georg Fromme & Co., Wien, 553 pp., 133 pls.
- Błeszyński S. 1970. New genera and species of tropical Crambinae (Studies on the Crambinae, Lepidoptera, Pyralidae, Part 48). *Tijdschrift voor Entomologie* 113: 1-26.
- Hampson G.F. 1919. Descriptions of new Pyralidae of the subfamilies Crambinae and Siginae. *Annals and Magazine of Natural History, including Zoology, Botany and Geology, London* (ser. 9) 4: 53-68, 137-154, 305-326.
- Klots A.B. 1970. Lepidoptera (pp. 115-130). In: Tuxen S. L. (Ed.). Taxonomist's glossary of genitalia in insects. 2nd Revised and Enlarged Edition. *Munksgaard, Copenhagen*, 359 pp.
- Kristensen N.P. 2003. Skeleton and muscles: adults (pp. 39-131). In: Kristensen N.P. (Ed.). Lepidoptera, moths and butterflies. Vol. 2. Morphology, physiology, and development. *Handbook of Zoology IV (36)*. Walter de Gruyter, Berlin and New York, XII + 564 pp.
- Ragonot E.L. 1889. In: de Joannis J., Ragonot E.L. Descriptions de genres nouveaux et espèces nouvelles de Lépidoptères. Annales de la Société Entomologique de France, Paris (ser. 6) 8[3 (1888)]: 271-284.
- Robinson G.S. 1976. The preparation of slides of Lepidoptera genitalia with special reference to the Microlepidoptera. *Entomologist's Gazette* 27: 127-132.
- Swinhoe C. 1886 On the Lepidoptera of Bombay and the Deccan. Part IV. Heterocera (continued). Proceedings of the General Meetings for Scientific Business of the Zoological Society of London 1885 (4): 852-886, pls 56-57.