

## **Characterization of Microsatellite Markers for the Vulnerable Grassland Forb *Senecio macrocarpus* (Asteraceae)**

Authors: Ahrens, Collin W., and James, Elizabeth A.

Source: *Applications in Plant Sciences*, 1(11)

Published By: Botanical Society of America

URL: <https://doi.org/10.3732/apps.1300041>

---

BioOne Complete ([complete.BioOne.org](https://complete.BioOne.org)) is a full-text database of 200 subscribed and open-access titles in the biological, ecological, and environmental sciences published by nonprofit societies, associations, museums, institutions, and presses.

Your use of this PDF, the BioOne Complete website, and all posted and associated content indicates your acceptance of BioOne's Terms of Use, available at [www.bioone.org/terms-of-use](https://www.bioone.org/terms-of-use).

Usage of BioOne Complete content is strictly limited to personal, educational, and non - commercial use. Commercial inquiries or rights and permissions requests should be directed to the individual publisher as copyright holder.

---

BioOne sees sustainable scholarly publishing as an inherently collaborative enterprise connecting authors, nonprofit publishers, academic institutions, research libraries, and research funders in the common goal of maximizing access to critical research.

## CHARACTERIZATION OF MICROSATELLITE MARKERS FOR THE VULNERABLE GRASSLAND FORB *SENECIO MACROCARPUS* (ASTERACEAE)<sup>1</sup>

COLLIN W. AHRENS<sup>2,3</sup> AND ELIZABETH A. JAMES<sup>2</sup>

<sup>2</sup>Royal Botanic Gardens Melbourne, National Herbarium of Victoria, Private Bag 2000, Birdwood Avenue, South Yarra, Victoria 3141, Australia

- *Premise of the study:* Development of microsatellite markers for the vulnerable forb *Senecio macrocarpus* was performed to begin an assessment of its population structure and breeding method to aid in the conservation of the species in Victoria, Australia.
- *Methods and Results:* Fifteen microsatellite markers were developed for *S. macrocarpus* from 454 pyrosequencing. The markers were tested on 104 individuals from four populations. The markers produced between two and seven alleles per locus while the expected heterozygosity ranged from 0.20 to 0.67 and the observed heterozygosity ranged from 0.00 to 1.00. The observed heterozygosity is suggestive that the populations may be apomictic.
- *Conclusions:* The microsatellite markers developed for *S. macrocarpus* are intended to be used on future studies that aim to assess the population genetics and local breeding dynamics of the species with an emphasis on conservation.

**Key words:** 454 sequencing; apomixis; Asteraceae; conservation; microsatellites; polyploidy; *Senecio macrocarpus*.

The Natural Temperate Grassland of the Victorian Volcanic Plain currently occupies less than 1% of its former range (McDougall and Kirkpatrick, 1994). Consequently, many of the plant species that are endemic to the Volcanic Plain are in decline due to habitat loss. *Senecio macrocarpus* F. Muell. ex Belcher (Asteraceae) is no exception. It is a perennial forb that is found in southeastern Australia. It is listed as vulnerable under the Australian *Environment Protection and Biodiversity Conservation Act 1999* (EPBC Act). Currently, *S. macrocarpus* populations are highly fragmented due to agricultural and industrial development and occupy areas that can be categorized as urban fringe, roadside patches, and native remnant vegetation. It has been used with some success for revegetating native grasslands in Victoria (McDougall and Morgan, 2005). To aid grassland revegetation efforts and secure the conservation of *S. macrocarpus*, understanding the genetic variability on the population level is crucial to create viable populations that will persist. Genomic microsatellite markers would be valuable for uncovering the variation across the entire population. Microsatellite markers are available for related species, including *S. madagascariensis* Poir. (Le Roux and Wiczorek, 2006) and *S. inaequidens* DC. (Justy and Maurice, 2012). In Australia, the genus *Senecio* L. consists of a diverse group of native species with both self-compatible and self-incompatible species that are thought to be wind dispersed (Belcher, 1994); the genus also includes differing ploidy levels, with the majority of

species being tetraploid or hexaploid (Lawrence, 1985). We characterize microsatellite markers for *S. macrocarpus* to identify the genetic variation within and among populations and to understand how Victorian species are related. We included two other Victorian species to test for possible cross-amplification. The developed markers will be used to identify proper conservation strategies by assessing the level of genetic variability population wide.

### METHODS AND RESULTS

Voucher specimens from each population were deposited in the herbarium at the Royal Botanic Gardens, Melbourne (voucher no.: Messent Conservation Park-23705226A, Deep Lead Flora Reserve-2370528A, Bulban Road-2370530A, Derrimut Grassland Reserve-2370532A). Leaf samples were dried and kept in silica gel until DNA extraction could be performed. Genomic DNA was extracted at the Australian Genome Research Facility (AGRF, Adelaide, South Australia, Australia) from 20 mg of plant material using the cetyltrimethylammonium bromide (CTAB) method. The extracted DNA was quantified using the QuantiFluor system (Promega Corporation, Madison, Wisconsin, USA).

Isolated DNA from one *S. macrocarpus* individual was sequenced by a Roche GS FLX sequencing platform (454 Life Sciences, a Roche Company, Branford, Connecticut, USA) for 1/8 coverage. The 454 results were processed in the program QDD (Megléczy et al., 2009) for selection and design of microsatellite primers. The QDD output isolated a total of 22,909 possible microsatellite primer pairs. Paring down the number of possible markers was done with selection criteria that consisted of penalty score (lowest score indicates 'best' primer pair at the locus), size of PCR product (100–400 bp), primer designs (ranked by primer length, melting temperature, percentage of G and C nucleotides, and mononucleotide repeats), and motif repeat size. The 96 selected loci were manufactured (Integrated DNA Technologies, Coralville, Iowa, USA) and screened for amplification and variability in seven *S. macrocarpus* individuals using the three-primer approach described in Blacket et al. (2012). This method enabled efficient amplification of loci in multiplex PCRs using all four fluorochromes (FAM, VIC, NED, PET). Each fluorochrome was attached to

<sup>1</sup>Manuscript received 16 May 2013; revision accepted 1 July 2013.

The authors acknowledge the Royal Botanic Gardens Melbourne; the study was funded by Hansons Construction Materials Pty Ltd.

<sup>3</sup>Author for correspondence: collin.ahrens@rbg.vic.gov.au

doi:10.3732/apps.1300041

TABLE 1. Characteristics of 15 microsatellite primers developed for *Senecio macrocarpus* ( $2n = 60$ ).

Locus	GenBank accession no.	Primer sequences (5'–3')	Repeat motif	$T_a$ (°C)	Size range (bp)	Label
sm003*	KF018218	F: TGAACCACTCTCAACCAC R: GTGAGCTTGGACTCATGGGA	(ACC) <sub>6</sub>	60	106–128	NED
sm008*	KF018219	F: GTGAGATAAGAGCATAACGTACCA R: GGAAGGAATTGATTGCGAGTT	(AAAC) <sub>5</sub>	60	136–140	PET
sm010*	KF018220	F: TTGGACGAGTTGGTGGAGTC R: TGATCTAAGGAAGTCCGGCA	(ACC) <sub>7</sub>	60	119–148	VIC
sm015*	KF018221	F: GTCTAGGCGGTTCCATGTTG R: CACTGCGGAGCTTACCTTGT	(ACC) <sub>5</sub>	60	149–158	NED
sm020*	KF018222	F: GGGAGCTAAATTCTGCGT R: TCCCGGTCTATATTGGCCTT	(AAC) <sub>8</sub>	60	131–160	PET
sm023*	KF018223	F: CTGCCCAATTGTTGACCTCT R: GCATCGATTTGAAAAGTTGCC	(AC) <sub>10</sub>	60	148–160	FAM
sm026*	KF018224	F: CCTTCAACCACCGAGGATAA R: CACCATATCAATCGCATCCA	(AC) <sub>6</sub>	60	174–180	PET
sm036*	KF018225	F: TTCTTCGAATCGGAAACAGC R: CCAACCGTACACATCTCCGT	(ACT) <sub>5</sub>	60	183–191	FAM
sm040*	KF018226	F: TCATCCTCAGACTCGTCCGGT R: GAGCGTTCGGATCAGAGTA	(AGT) <sub>5</sub>	60	195–260	VIC
sm043*	KF018227	F: ATGCAAGGTTCTTCGTTGGA R: TGTTTGCATCAACATTTGGG	(AAG) <sub>5</sub>	60	199–211	FAM
sm049*	KF018228	F: ATGCAAAATTCACCATGGACC R: CCTCACGAAATGTCTCTTTG	(ACC) <sub>5</sub>	60	196–228	VIC
sm051*	KF018229	F: CGCTCCTTGAATTTAGCAA R: CAGGCCCAATAATGACCAA	(AT) <sub>7</sub>	60	221–268	PET
sm052*	KF018230	F: AAGCTTGCTTACGAGGCTCA R: CACAGCAACTGAACACCCAA	(AG) <sub>8</sub>	60	216–244	NED
sm057*	KF018231	F: GGGCTGGAACAACAAGAGAA R: TGGCTGGACTTGAAGGCTAT	(AC) <sub>7</sub>	60	236–246	VIC
sm064*	KF018232	F: ACCTCACTTGGTGGGATTTG R: CCCAACGGAATTAACCTGGAC	(AAC) <sub>8</sub>	60	252–264	FAM
sm002		F: TACCGCAACCTCTCACTTCC R: GGTTTGAGGATGGTGGATTG	(AGT) <sub>5</sub>	60	113	VIC
sm005		F: AAGTCAAGGCGATGAAGCAG R: AAGTGGATTCCCTTTCTGGG	(AG) <sub>7</sub>	60	119	FAM
sm011		F: GGCAAAATCATTTACAGGCCA R: GGACTTGAACCCCTTGCTTGA	(AAC) <sub>5</sub>	60	135	NED
sm012		F: ATGCCACCGTTAGAAGCAGA R: CAATGTTGACAGCCATTTCC	(AAG) <sub>5</sub>	60	138	PET
sm034		F: ACGTGAAGAATGCAGCCAT R: TCTCTGCATCACGAAACGAA	(ACT) <sub>5</sub>	60	183	PET
sm039		F: CCATGTGGAGAGGTACCAGC R: CATGGAGCTCAATCCTCCTC	(AAG) <sub>5</sub>	60	190	FAM
sm046		F: GCAGAATCAGCAAGGGAAA R: GGAACATGGACATTGCTTTGA	(ACT) <sub>12</sub>	60	189	PET
sm047		F: GGAATAGTTGCCTATATTATGTGTTCCG R: AATCCAATGGGTTGGCTTCT	(AAT) <sub>8</sub>	60	211	FAM
sm053		F: CACAATCCAGACCAACCCAT R: GGTACATACGGCAGCGAGAG	(AATC) <sub>5</sub>	60	222	VIC
sm061		F: TTCCAGTCAATTCAGTTCCA R: TGGATTGCTGATGATCTTGG	(AGT) <sub>6</sub>	60	242	VIC
sm068		F: TTTCACCTTCTCCTTGACCA R: ATCGCTCATTGCATCTTTGC	(AGT) <sub>6</sub>	60	244	NED
sm069		F: TTGGCCACTTTGCTAGAGA R: TGTTTCATGCATGCTGGAGT	(AAC) <sub>5</sub>	60	262	VIC
sm070		F: ATCCACCTGCCATCTGAGAA R: ACGCTGGGTCGTATGCTAAA	(AGT) <sub>5</sub>	60	277	FAM
sm071		F: CAATGTTTGGCGGATAGAT R: CGTGTGATATAAATGAAACCGAA	(AAG) <sub>5</sub>	60	281	PET
sm077		F: CCAAACATTAATTGGGGCTG R: TGTTAATTGAGTGCCCGGTT	(AT) <sub>5</sub>	60	300	PET
sm082		F: TCGGCACCTCGTATGGAGTTT R: GCTAAACAGGAGGGTTGAGAA	(AG) <sub>5</sub>	60	311	NED
sm083		F: GCGTAGCCAACCTTTGAGAG R: GGAGTGGTTAACGGACATGC	(AG) <sub>5</sub>	60	313, 317	VIC
sm088		F: TTCATCTACAACGGGATGGG R: GGAGATCACTCCATTCTGTGA	(AT) <sub>5</sub>	60	334	FAM
sm090		F: GTTACCTTGAGGGAAGCGGT R: TGAAGCGCGTAAGGTCTAGG	(AGG) <sub>5</sub>	60	354	NED
sm091		F: TCTGCTTGAACCAACCCAGT R: CAACACTAAGACTGAGCAAATGGA	(AT) <sub>5</sub>	60	349	VIC

TABLE 1. Continued.

Locus	GenBank accession no.	Primer sequences (5′–3′)	Repeat motif	$T_a$ (°C)	Size range (bp)	Label
sm095		F: GACCAGGCCAACCAAGAGAT R: TCTGTCCATTTCCTTCCTCCA	(AG) <sub>5</sub>	60	374	VIC

Note:  $T_a$  = annealing temperature.

\*Polymorphic and variable locus; primer pairs for these loci were used in this study, while primer pairs for the other loci presented in the table were not.

one of four unique universal “tails” (Blackett et al., 2012). Forward locus-specific primers were allocated to one of the four tails, which then dictated the fluorochrome labeling locus-specific PCR products. PCRs consisted of 10- $\mu$ L reactions with approximately 20 ng of DNA, 5  $\mu$ L Master Mix Type-it Kit (QIAGEN, Hilden, Germany), 0.1  $\mu$ L of forward primers (10  $\mu$ M), 0.2  $\mu$ L of reverse primers (10  $\mu$ M), and 0.1  $\mu$ L of each dye (10  $\mu$ M; FAM, VIC, NED, PET). All PCRs were carried out under the same conditions: 5 min denaturation at 95°C, followed by 32 cycles of 95°C for 30 s, annealing at 60°C for 1 min 30 s and 72°C for 30 s, and a final extension at 60°C for 30 min. Each reaction was multiplexed with four different primer pairs. The initial primer screening left 15 primer pairs that were amplifiable and variable among the seven individuals (Table 1). These primer pairs included five dinucleotide, nine trinucleotide, and one tetranucleotide repeat motifs. The 15 primer pairs were subsequently used on 104 individuals among four *S. macrocarpus* populations: Messent Conservation Park (36.10156°S, 139.74366°E), Deep Lead Flora Reserve (37.01269°S, 142.72166°E), Bulban Road (37.92488°S, 144.59474°E), and Derrimut Grassland Reserve (37.80962°S, 144.71980°E) with the same protocol used in the amplification described above. Two different Victorian species were used to test cross-species amplification. The more common *S. quadridentatus* Labill. (GPS coordinates: 37.62569°S, 144.96216°E; unvouchered specimen determined by Neville Walsh) is closely related to *S. macrocarpus* and is widespread in southern Australia, whereas *S. behrianus* Sond. & F. Muell. ex Sond. (voucher no.: MEL 2296795A) is a rare Victorian endemic known to reproduce both clonally and by seed. Fragment analysis of the PCR product was performed via capillary electrophoresis on an ABI 3130xl (Applied Biosystems, Foster City, California, USA) by Macrogen (Seoul, South Korea). Alleles were sized against the internal size standard GeneScan 500 LIZ (Applied Biosystems) and scored using Geneious Software version 5.6.2 (Biomatters Ltd., Auckland, New Zealand). All 15 microsatellite markers showed a clear consistent amplification pattern; locus sm043 had one or two alleles (three individuals had three alleles), all individuals had three alleles for locus sm052, and all other loci had one or two alleles. Chromosome counts show that *S. macrocarpus* is hexaploid ( $2n = 60$ ; data not shown), while *S. quadridentatus* is tetraploid ( $2n = 40$ ; Lawrence, 1985) and *S. behrianus*’ ploidy is unknown. The allele numbers

per individual are unexpectedly low for a hexaploid outcrossing species. However, these patterns could be attributed to apomixis and *S. macrocarpus*’ evolutionary history.

The number of alleles, observed heterozygosity, and expected heterozygosity were calculated in GenoDive version 2.0b24 (Meirmans and Van Tienderen, 2004) for each locus within and among the four populations (Table 2). The deviations from Hardy–Weinberg equilibrium were extreme and not typical of predominantly outcrossing species. The results show that some alleles are fixed (as homozygotes and heterozygotes) within but not among populations. Apomixis could be contributing to these patterns as it occurs naturally in Asteraceae in at least 22 genera and seven tribes (reviewed in Noyes, 2007). Indeed, our observation of low pollen to ovule ratio and lack of pollen exertion, in addition to the population structure found in *S. macrocarpus*, further support this idea.

In terms of cross-amplification, only two markers failed to amplify in the other two species. Five markers amplified and were polymorphic for *S. behrianus* while a different set of five markers amplified and were polymorphic for *S. quadridentatus*. Three markers were monomorphic for *S. behrianus* while eight markers were monomorphic for *S. quadridentatus* (Table 3).

## CONCLUSIONS

The 15 microsatellite loci characterized herein for the vulnerable forb *S. macrocarpus* are being used for research to document the population structure, gene flow, and breeding system, and to assess population viability of current and future populations. They will also be used to investigate interspecific relationships between Australian species. The results of these studies will aid in the conservation of *S. macrocarpus* in a fragmented Australian landscape.

TABLE 2. Results for primer screening of all samples of 15 microsatellite loci from four populations of *Senecio macrocarpus* ( $2n = 60$ ).<sup>a</sup>

Locus	Messent CP ( $N = 54$ )			Derrimut Grassland ( $N = 20$ )			Bulban Road ( $N = 15$ )			Deep Lead FR ( $N = 15$ )			Total ( $N = 104$ )		
	A	$H_o$	$H_e$	A	$H_o$	$H_e$	A	$H_o$	$H_e$	A	$H_o$	$H_e$	A	$H_o$	$H_e$
sm003	3	0.32	0.28	1	0.00	0.00	2	1.00	0.52	2	1.00	0.52	3	0.58	0.43
sm008	1	0.00	0.00	1	0.00	0.00	1	0.00	0.00	1	0.00	0.00	2	0.00	0.50
sm010	7	0.31	0.55	0	—	—	1	0.00	0.00	1	0.00	0.00	7	0.10	0.49
sm015	2	0.94	0.50	2	1.00	0.51	1	0.00	0.00	1	0.00	0.00	2	0.49	0.37
sm020	3	0.08	0.56	1	0.00	0.00	2	0.07	0.07	2	0.07	0.07	5	0.05	0.67
sm023	3	0.00	0.52	1	0.00	0.00	1	0.00	0.00	1	0.00	0.00	3	0.00	0.47
sm026	2	0.00	0.14	1	0.00	0.00	1	0.00	0.00	1	0.00	0.00	2	0.00	0.50
sm036	2	0.87	0.49	1	0.00	0.00	1	0.00	0.00	1	0.00	0.00	2	0.22	0.20
sm040	2	0.00	0.07	2	0.05	0.05	1	0.00	0.00	1	0.00	0.00	2	0.01	0.50
sm043	5	1.00	0.63	2	1.00	0.51	3	1.00	0.58	2	1.00	0.52	5	1.00	0.61
sm049	2	1.00	0.51	1	0.00	0.00	2	1.00	0.52	2	0.93	0.52	2	0.73	0.48
sm051	0	—	—	1	0.00	0.00	2	0.00	0.40	2	0.07	0.07	3	0.02	0.56
sm052	4	1.00	0.68	3	1.00	0.68	3	1.00	0.68	3	1.00	0.68	4	1.00	0.71
sm057	2	0.04	0.10	2	0.10	0.10	2	1.00	0.52	2	1.00	0.52	3	0.53	0.41
sm064	3	0.07	0.62	1	0.00	0.00	1	0.00	0.00	1	0.00	0.00	3	0.02	0.58

Note: A = number of alleles per locus;  $H_e$  = expected heterozygosity;  $H_o$  = observed heterozygosity.

<sup>a</sup>Population locality information: Messent Conservation Park (36.10156°S, 139.74366°E), Derrimut Grassland Reserve (37.80962°S, 144.71980°E), Bulban Road (37.92488°S, 144.59474°E), Deep Lead Flora Reserve (37.01269°S, 142.72166°E).

TABLE 3. Cross-amplification of 15 microsatellite markers for two Victorian species related to *Senecio macrocarpus* ( $2n = 60$ ): *S. behrianus* ( $2n = \text{unknown}$ ) and *S. quadridentatus* ( $2n = 40$ ).<sup>a</sup>

Species	N	sm003	sm008	sm010	sm015	sm020	sm023	sm026	sm036	sm040	sm043	sm049	sm051	sm052	sm057	sm064
<i>S. behrianus</i>	4	P (3)	—	—	NV (2)	—	P (3)	P (3)	M (1)	—	P (3)	—	NV (2)	P (2)	—	—
<i>S. quadridentatus</i>	5	M (1)	—	M (1)	NV (2)	—	P (4)	M (1)	M (1)	M (1)	P (3)	NV (2)	P (2)	NV (2)	P (3)	P (2)

Note: — = failed amplification; M = monomorphic; NV = not variable; P = polymorphic.

<sup>a</sup>Numbers in parentheses indicate the number of alleles amplified at the locus.

### LITERATURE CITED

- BELCHER, R. O. 1994. *Senecio*. In Flora of Australia, vol. 49. Australian Government Publishing Service, Canberra, Australia.
- BLACKET, M. J., C. ROBIN, R. T. GOOD, S. F. LEE, AND A. D. MILLER. 2012. Universal primers for fluorescent labelling of PCR fragments: An efficient and cost-effective approach to genotyping by fluorescence. *Molecular Ecology Resources* 12: 456–463.
- JUSTY, F., AND S. MAURICE. 2012. Isolation and characterization of polymorphic microsatellite markers for the invasive *Senecio inaequidens* DC. (Asteraceae). *Conservation Genetics Resources* 4: 703–705.
- LAWRENCE, M. E. 1985. *Senecio* L. (Asteraceae) in Australia: Rcombination [sic] systems of a polyploid series. *Australian Journal of Botany* 33: 209–219.
- LE ROUX, J. J., AND A. M. WIECZOREK. 2006. Isolation and characterization of polymorphic microsatellite markers from fireweed, *Senecio madagascariensis* Poir. (Asteraceae). *Molecular Ecology Notes* 7: 327–329.
- MCDUGALL, K. L., AND J. B. KIRKPATRICK. 1994. Conservation of lowland native grasslands in south-eastern Australia. World Wide Fund for Nature, Sydney, New South Wales, Australia.
- MCDUGALL, K. L., AND J. W. MORGAN. 2005. Establishment of native grassland vegetation at Organ Pipes National Park near Melbourne, Victoria: Vegetation changes from 1989 to 2003. *Ecological Management & Restoration* 6: 34–42.
- MEGLÉCZ, E., C. COSTEDOAT, V. DUBUT, A. GILLES, T. MALAUSA, N. PECH, AND J.-F. MARTIN. 2009. QDD: A user-friendly program to select microsatellite markers and design primers from large sequencing projects. *Bioinformatics (Oxford, England)* 26: 403–404.
- MEIRMANS, P. G., AND P. H. VAN TIENDEREN. 2004. GENOTYPE and GENODIVE: Two programs for the analysis of genetic diversity of asexual organisms. *Molecular Ecology Notes* 4: 792–794.
- NOYES, R. D. 2007. Apomixis in the Asteraceae: Diamonds in the rough. *Functional Plant Science and Biotechnology* 1: 207–222.