

Microsatellite Markers for *Senna spectabilis* var. *excelsa* (Caesalpinioideae, Fabaceae)

Authors: López-Roberts, M. Cristina, Barbosa, Ariane R., Queiroz, Luciano Paganucci de, and Berg, Cássio van den

Source: Applications in Plant Sciences, 4(1)

Published By: Botanical Society of America

URL: <https://doi.org/10.3732/apps.1500062>

BioOne Complete (complete.BioOne.org) is a full-text database of 200 subscribed and open-access titles in the biological, ecological, and environmental sciences published by nonprofit societies, associations, museums, institutions, and presses.

Your use of this PDF, the BioOne Complete website, and all posted and associated content indicates your acceptance of BioOne's Terms of Use, available at www.bioone.org/terms-of-use.

Usage of BioOne Complete content is strictly limited to personal, educational, and non - commercial use. Commercial inquiries or rights and permissions requests should be directed to the individual publisher as copyright holder.

BioOne sees sustainable scholarly publishing as an inherently collaborative enterprise connecting authors, nonprofit publishers, academic institutions, research libraries, and research funders in the common goal of maximizing access to critical research.

MICROSATELLITE MARKERS FOR *SENNA SPECTABILIS* VAR. *EXCELSA* (CAESALPINIOIDEAE, FABACEAE)¹

M. CRISTINA LÓPEZ-ROBERTS^{2,4}, ARIANE R. BARBOSA², LUCIANO PAGANUCCI DE QUEIROZ³,
AND CÁSSIO VAN DEN BERG³

²Pós-graduação em Botânica, Universidade Estadual de Feira de Santana (UEFS), Av. Transnordestina s/n, 44036-900 Feira de Santana, Bahia, Brazil; and ³Departamento Ciências Biológicas, Universidade Estadual de Feira de Santana (UEFS), Av. Transnordestina s/n, 44036-900 Feira de Santana, Bahia, Brazil

- **Premise of the study:** *Senna spectabilis* var. *excelsa* (Fabaceae) is a South and Central American tree of great ecological importance and one of the most common species in several sites of seasonally dry forests. Our goal was to develop microsatellite markers to assess the genetic diversity and structure of this species.
- **Methods and Results:** We designed and assessed 53 loci obtained from a microsatellite-enriched library and an intersimple sequence repeat library. Fourteen loci were polymorphic, and they presented a total of 39 alleles in a sample of 61 individuals from six populations. The mean values of observed and expected heterozygosities were 0.355 and 0.479, respectively. Polymorphism information content was 0.390 and the Shannon index was 0.778.
- **Conclusions:** Polymorphism information content and Shannon index indicate that at least nine of the 14 microsatellite loci developed are moderate to highly informative, and potentially useful for population genetic studies in this species.

Key words: Caatinga; Fabaceae; genetic variation; microsatellite library; polymorphic loci; *Senna spectabilis*.

Senna Mill. is a large, diverse, and widespread genus of Caesalpinioideae (Fabaceae), with about 350 species. Most species are American, and some species also occur in Africa, Madagascar, Australia, and Asia (Marazzi et al., 2006). *Senna spectabilis* (DC.) H. S. Irwin & Barneby is a deciduous tree or shrub ranging from southern Mexico to the Caribbean and northern Argentina and including two varieties: *S. spectabilis* var. *spectabilis* and *S. spectabilis* var. *excelsa* (Schrad.) H. S. Irwin & Barneby (Irwin and Barneby, 1982). This taxon has been introduced to many countries (including the United States, Malaysia, and Africa) as an ornamental street tree, where it is often regarded as an invasive plant (Richardson and Rejmánek, 2011).

Senna spectabilis var. *excelsa* is widespread in the Caatinga Phytogeographical Domain of northeastern Brazil (Queiroz, 2009). Due to its long flowering period (December to April), it is an important source of pollen for “buzzing bees” (*Xylocopa*, *Bombus*, and *Centris* species), which constitute its main pollinators

(Manente-Balestieri and Machado, 1999). This tree has been used by local people as a source of fuel wood and timber, as well as fodder for sheep and goats (Santos et al., 2013). Extracts of this plant also have several applications in folk medicine, and recent studies confirm sedative, anticonvulsive, antimalarial, antimicrobial, cytotoxic, and leishmanicidal activities (de Albuquerque Melo et al., 2014). Despite this, there are no studies that assess the levels of genetic diversity in populations of *S. spectabilis*. The only study in this species assessed the genetic diversity in germplasm collections using RAPD markers (Santos et al., 2013). In the current study, our goal was to develop microsatellite markers to assess the genetic variation and population structure of *S. spectabilis* var. *excelsa*, generating useful tools for future studies and management of this species.

METHODS AND RESULTS

A leaf sample from a single individual of *S. spectabilis* var. *excelsa* cultivated on the campus of Universidade Estadual de Feira de Santana was collected and deposited in the herbarium of Universidade Estadual de Feira de Santana (HUEFS; *H. Huaylla* & *M. C. L. Roberts* 3737, see Appendix 1). Genomic DNA was extracted using the cetyltrimethylammonium bromide (CTAB) 2x protocol (Doyle and Doyle, 1990) adapted for Eppendorf microtubes (Axygen Scientific, Union City, California, USA). Two types of microsatellite libraries were constructed: a microsatellite-enriched library, following the protocol of Billotte et al. (1999), and an intersimple sequence repeat (ISSR) library (Provan and Wilson, 2007). Fragments obtained by both strategies were linked into the pGEM-T Easy Vector System (Promega Corporation, São Paulo, Brazil) and subsequently transformed and cloned into TOP10 competent cells (Invitrogen, Life Technologies, São Paulo, Brazil). The positive clones were amplified by PCR using the T7 (5'-TAATACGACTACTATAGGG-3') and SP6 (5'-ATTTAGGTGACTATAGAA-3') primers with the following program: premelt of 94°C for 5 min, 35x (denaturation at 94°C for 15 s, annealing at

¹Manuscript received 1 June 2015; revision accepted 4 August 2015.

The authors thank Daniel Nascimento de Oliveira for helping with the construction of the SSR libraries; Ricardo Vilas-Bôas and Élvia R. Souza for collaboration in the laboratory; and Hibert Hually, Cosme Correia dos Santos, and Wallace São Matheus for support in the field. This work is part of Núcleo de Excelência em Sistemática e Variabilidade Molecular de Plantas e Fungos (Fundação de Amparo à Pesquisa do Estado da Bahia [FAPESB] PNX0014/2009) and Patterns of Leguminosae diversity in Brazilian biomes (Conselho Nacional de Desenvolvimento Científico e Tecnológico [CNPq] 563084/2010-3).

⁴Author for correspondence: macrissroberts@gmail.com

doi:10.3732/apps.1500062

55°C for 35 s, and extension at 72°C for 90 s), followed by a final extension at 72°C for 7 min. After quantification performed in agarose gel electrophoresis with a 100-bp ladder (Ludwig Biotech, Alvorada, Rio Grande do Sul, Brazil), fragments larger than 500 bp were sequenced using the Big Dye 3.1 Terminator Cycle Sequencing Kit and an ABI 3130XL automated sequencer (Applied Biosystems, São Paulo, Brazil). Identification of simple sequence repeats was performed with the software SSRIT (Simple Sequence Repeat Identification Tool; <http://archive.gramene.org/db/markers/ssrtool>) and Imperfect Microsatellite Extractor (Mudunuri and Nagarajaram, 2007). Primer design was done using the software Primer3Plus (Untergasser et al., 2007) with the following criteria: product size range 100–300 bp, primer melting temperature (T_m) 50–60°C (with a maximum difference of 2°C), and GC content 40–60%.

Genotyping reactions were performed in 10- μ L reactions with Top Taq Master Mix (QIAGEN, São Paulo, Brazil), adding 0.15 μ M of the forward primer (tailed at the 5' with the M13 universal sequence: 5'-CACGACGTTG-TAAAACGAC-3'), 0.30 μ M of the reverse primer, 0.060 μ M of the M13-labeled tail (with 6-FAM, VIC, NED, or PET fluorescent dyes; Applied Biosystems) and approximately 5 ng of individual DNA. Optimal annealing temperature for each primer pair was determined using touchdown (60–54°C, 58–48°C) and gradient (from 62–45°C) PCR programs. Amplicons were visualized by agarose gel electrophoresis (1.5%) with a 100-bp ladder. Microsatellite profiles were analyzed with GeneMapper 4.0 (Applied Biosystems). For polymorphism assessment, we sampled 61 individuals from six populations of *S. spectabilis* var. *excelsa* (8–21 individuals per population; Appendix 1). Cross-amplification was tested in *S. spectabilis* var. *spectabilis* using a foliar sample from one individual deposited at HUEFS (Queiroz L. P. de 13481, see Appendix 1). The estimated number of alleles, observed heterozygosity (H_o), expected heterozygosity (H_e), Shannon index, and Hardy–Weinberg equilibrium (HWE) deviations were calculated using GenAlEx 6.5 (Peakall and Smouse, 2006). We also performed a linkage disequilibrium test using GENEPOP (Raymond and Rousset, 1995) to check if genotypes at one locus are independent from genotypes at the other loci. The polymorphism information content (PIC) was calculated with the online tools of the Centre for Genomic Research at the University of Liverpool (<http://www.genomics.liv.ac.uk/animal/Protocol1.html>).

We found 146 perfect and imperfect microsatellite loci in 79 fragments of the enriched library, with motifs of four or more repeats. In the ISSR library, we

found 54 microsatellites in only 13 sequences. The proportions of perfect and imperfect-motif microsatellites obtained are shown in Appendix S1. From both sets of loci, we designed a total of 53 primer pairs, of which 34 were designed from the enriched library.

A total of 35 microsatellite loci were successfully amplified, from which 14 were polymorphic. Variation was observed in both perfect and imperfect motifs, particularly in di- and trinucleotide repeats (Table 1). The linkage disequilibrium test detected significant values for the following pairwise comparisons: SS_5/SS_51 ($P = 0.0131$) and SS_51/SS_52 ($P = 0.0097$), which should be used carefully. The allele number per locus ranged from two to five, with a total of 39 alleles identified (Table 2). H_o and H_e values varied from 0.037 to 0.727 and 0.036 to 0.743, respectively. Significant deviations from expected HWE proportions ($P < 0.01$) were detected in seven loci. Values of PIC ranged from 0.037 to 0.697 with a mean of 0.390. The Shannon index varied from 0.092 to 1.423 (mean 0.778). Nine loci presented values of PIC > 0.3 and Shannon index > 0.6, indicating that these are moderately to highly informative microsatellites. Successful cross-amplification was observed in the single tested individual of *S. spectabilis* var. *spectabilis* (Table 2).

CONCLUSIONS

We successfully developed and amplified a set of polymorphic microsatellite markers in *S. spectabilis* var. *excelsa*, which could prove very useful in population genetic studies. Due to the successful cross-amplification of these microsatellite loci in *S. spectabilis* var. *spectabilis*, we expect these markers could also be used to assess genetic variation of germplasm collections and to support plant breeding programs in this species. *Senna spectabilis* var. *excelsa* is one of the few tree species from the caatinga for which microsatellites have been developed, and their use should foster the understanding of patterns of diversity in this fragile ecosystem with high levels of endemic biodiversity.

TABLE 1. Characterization of 14 polymorphic microsatellite markers developed for *Senna spectabilis* var. *excelsa*.

Locus	Primer sequences (5'–3')	Repeat motif	Allele size range (bp) ^a	M13 5' PIG-tail	T_a (°C)	GenBank accession no.
SS_1	F: TTCTGAAGCAGACGGAAGGT R: CTCACCTTCCAAATTCAAAGC	(AGAGA) ₄	132–142	PET	58	KJ398419
SS_5	F: GTCCTTTTCACCGATTGA R: CTCCTGTCAAACACTCACAAA	(AG) ₉	136–148	6-FAM	59	KJ398420
SS_10	F: CACAGCCAAGAGGGAGAAAG R: CCTTCTCAATGGCCTTCTCTT	(AGA) ₅	187–196	VIC	58	KJ398421
SS_14	F: GAGTATGGGGAATACGAATC R: TGGGAAGTCAGATACCAT	(TC) ₃ C(GT) ₃	305–307	NED	56	KJ398422
SS_15	F: GGTATCTGACTTCCCAA R: AGATGTCCGGATTGCTAC	(ATTT) ₄	244–248	PET	56	KJ398422
SS_18	F: GCGATGCTCAAGTTTCTCTT R: CAAACTAAACCAAAGGACGA	(ATC) ₃ (TTC) ₃ (TC) ₆	147–149	NED	56	KJ398423
SS_24	F: GCCTGTGATCTGAAGGTGGA R: GGAGCAGTTTAAGCCAGCTTATT	(AG) ₉	115–125	NED	56	KJ398424
SS_26	F: GCGCCTCACTCATACATTGA R: GGTCCACCTCCGGGTAGTAT	(CT) ₄	124–132	VIC	56	KJ398425
SS_27	F: GAACTGGGAAGGCAGAAAAA R: GATTCTGGGCAGGCTCCTAT	(TC) ₄	314–316	PET	TD 60–54	KJ398426
SS_34	F: GAACTGGGAAGGCAGAAAAA R: ACTTTTCCCAACCTCCGTCT	(AT) ₄	153–157	VIC	TD 60–54	KJ398427
SS_41	F: CCAGAACACGACACAAATGG R: CGTGGACTAGCCTTCCCAATC	(AGAA) ₄	200–216	6-FAM	56	KJ398428
SS_48	F: TTGTCTTGGTGATCTCTTCTC R: GCACGTCAGTCAATATGCAA	(GTT) ₃	215–248	6-FAM	60	KJ398429
SS_51	F: TGGCAGTACCACCGTACAGA R: TCCCAATGGCTCACAGATG	(ATA)T(ATA) ₃ n ₉₀ (TCA) ₃	308–327	VIC	60	KJ398430
SS_52	F: ACTTCTTGATTTGGTGGCTGA R: CCTGCCAACCCAGATAGAAA	(AGT) ₃	118–127	6-FAM	57	KJ398431

Note: T_a = annealing temperature; TD = touchdown program.

^aExpected product size in base pairs (bp) including 5' M13 universal sequence.

TABLE 2. Population genetic parameters for 14 polymorphic microsatellite loci in *Senna spectabilis* var. *excelsa* ($n = 61$).

Locus	A	H_o	H_e	HWE ^a	I_s ^b	PIC ^c	Cross-amplification in <i>S. spectabilis</i> var. <i>spectabilis</i> ^d
SS_1	4	0.385	0.385	0.994 ^{ns}	0.672	0.308	Yes
SS_5	5	0.393	0.696	0.000 ^{***}	1.349	0.649	Yes
SS_10	2	0.200	0.455	0.012 [*]	0.647	0.352	No
SS_14	3	0.111	0.290	0.000 ^{***}	0.557	0.269	Yes
SS_15	2	0.455	0.351	0.168 ^{ns}	0.536	0.289	Yes
SS_18	2	0.200	0.295	0.107 ^{ns}	0.471	0.252	Yes
SS_24	5	0.727	0.743	0.805 ^{ns}	1.423	0.697	Yes
SS_26	4	0.545	0.479	0.000 ^{***}	0.879	0.425	Yes
SS_27	2	0.037	0.036	0.922 ^{ns}	0.092	0.037	Yes
SS_34	3	0.273	0.549	0.003 ^{**}	0.934	0.490	Yes
SS_41	4	0.500	0.696	0.198 ^{ns}	1.291	0.649	Yes
SS_48	3	0.048	0.534	0.000 ^{***}	0.841	0.435	Yes
SS_51	3	0.409	0.429	0.916 ^{ns}	0.687	0.355	Yes
SS_52	3	0.333	0.285	0.840 ^{ns}	0.519	0.256	Yes
Mean	3.214	0.355	0.479		0.778	0.390	

Note: A = number of alleles; H_e = expected heterozygosity; H_o = observed heterozygosity; HWE = probability of deviation from Hardy–Weinberg equilibrium; I_s = Shannon index; n = sample size; PIC = polymorphism information content.

^aSignificant deviation from Hardy–Weinberg equilibrium: * $P < 0.05$, ** $P < 0.01$, *** $P < 0.001$; ns = not significant.

^b $I_s > 0.6$ indicates a moderately to highly informative value.

^cPIC > 0.3 indicates a moderately to highly informative value.

^dCross-amplification was tested in a single individual of *Senna spectabilis* var. *spectabilis* (Appendix 1).

LITERATURE CITED

- BILLOTTE, N. P., P. J. L. LAGODA, A. M. RISTERUCCI, AND F. C. BAURENS. 1999. Microsatellite enriched libraries: Applied methodology for the development of SSR markers in tropical crops. *Fruits* 54: 277–288.
- DE ALBUQUERQUE MELO, G. M., M. C. SILVA, T. P. GUIMARÃES, K. M. PINHEIRO, C. B. DA MATTA, A. C. DE QUEIROZ, M. PIVATTO, ET AL. 2014. Leishmanicidal activity of the crude extract, fractions and major piperidine alkaloids from the flowers of *Senna spectabilis*. *Phytomedicine* 21: 277–281.
- DOYLE, J. J., AND J. S. DOYLE. 1990. Isolation of plant DNA from fresh tissue. *Focus (San Francisco, Calif.)* 12: 13–15.
- IRWIN, H. S., AND R. BARNEBY. 1982. The American Cassiniae. A synoptical revision of Leguminosae tribe Cassieae subtribe Cassiniae in the New World. *Memoirs of the New York Botanical Garden*, vol. 35. New York Botanical Garden Press, Bronx, New York, USA.
- MANENTE-BALESTIERI, F. C. D. L., AND V. L. L. MACHADO. 1999. Flowering entomofauna in *Cassia spectabilis* (L.) DC. (Leguminosae). *Anais da Sociedade Entomológica do Brasil* 28: 429–437.
- MARAZZI, B., P. K. ENDRESS, L. P. QUEIROZ, AND E. CONTI. 2006. Phylogenetic relationships within *Senna* (Leguminosae, Cassiinae) based on three chloroplast DNA regions: Patterns in the evolution of floral symmetry and extrafloral nectaries. *American Journal of Botany* 93: 288–303.
- MUDUNURI, S., AND H. NAGARAJARAM. 2007. IMEX: Imperfect microsatellite extractor. *Bioinformatics* 23: 1181–1187.
- PEAKALL, R., AND P. E. SMOUSE. 2006. GenAIEx 6: Genetic analysis in Excel: Population genetic software for teaching and research. *Molecular Ecology Notes* 6: 288–295.
- PROVAN, J., AND P. J. WILSON. 2007. Development of microsatellites for the peat moss *Sphagnum capillifolium* using ISSR cloning. *Molecular Ecology Notes* 7: 254–256.
- QUEIROZ, L. P. 2009. Leguminosas da Caatinga. Universidade Estadual de Feira de Santana, Feira de Santana, Brazil.
- RAYMOND, M., AND F. ROUSSET. 1995. GENEPOP (version 1.2): Population genetics software for exact tests and ecumenicism. *Journal of Heredity* 86: 248–249.
- RICHARDSON, D. M., AND M. REJMÁNEK. 2011. Trees and shrubs as invasive alien species: A global review. *Diversity & Distributions* 17: 788–809.
- SANTOS, M. F., R. B. ARAÚJO NETO, M. P. S. B. C. NASCIMENTO, AND P. S. C. LIMA. 2013. Determination of the genetic diversity among accessions of *Senna spectabilis* (canafístula) by using RAPD markers. *Genetics and Molecular Research* 12: 6149–6155.
- UNTERGASSER, A., H. NIJVEEN, X. RAO, T. BISSELING, R. GEURTS, AND J. A. M. LEUNISSEN. 2007. Primer3Plus, an enhanced web interface to Primer3. *Nucleic Acids Research* 35: W71–W74.

APPENDIX 1. Voucher and location information for populations of *Senna spectabilis* used in this study for microsatellite development, polymorphism testing, and cross-amplification. Voucher specimens are deposited at the herbarium of the Universidade Estadual de Feira de Santana (HUEFS), Feira de Santana, Bahia, Brazil.

Species	Population no. (n)	Collection locality	Geographic coordinates	Voucher no.
<i>Senna spectabilis</i> var. <i>excelsa</i> (Schrad.) H. S. Irwin & Barneby	Cultivated individual	Feira de Santana, Bahia, Brazil	12°11'58"S, 38°58'08"W	H. Huaylla & M. C. L. Roberts 3737
	Population 1 (21)	Morro de Chapéu, Bahia, Brazil	11°19'23.7"S, 41°00'51.4"W	C. C. Santos 828
	Population 2 (8)	Xique Xique, Bahia, Brazil	10°51'31.9"S, 42°37'09.6"W	C. C. Santos 833
	Population 3 (8)	Dirceu Arcoverde, Piauí, Brazil	09°19'53.6"S, 42°26'47.8"W	C. C. Santos 860
	Population 4 (8)	Milagres, Ceará, Brazil	7°15'56.7"S, 38°50'45.7"W	C. C. Santos 917
	Population 5 (8)	Pena Forte, Pernambuco, Brazil	7°49'08.8"S, 39°05'08.4"W	C. C. Santos 920
	Population 6 (8)	Jeremoabo, Bahia, Brazil	9°50'55"S, 38°15'51.2"W	C. C. Santos 934
<i>Senna spectabilis</i> (DC.) H. S. Irwin & Barneby var. <i>spectabilis</i>	Individual	Salvador de Jujuy, Mendieta, Jujuy, Argentina	24°19'00"S, 64°58'00"W	Queiroz, L. P. de 13481

Note: n = sample size.