

## **Development and Characterization of EST-SSR Markers in *Stipa breviflora* (Poaceae)**

Authors: Ren, Jing, Su, Zhi-Zhen, Dang, Zhen-Hua, Ding, Yu, Wang, Pei-Xuan, et al.

Source: Applications in Plant Sciences, 5(4)

Published By: Botanical Society of America

URL: <https://doi.org/10.3732/apps.1600157>

---

BioOne Complete ([complete.BioOne.org](https://complete.BioOne.org)) is a full-text database of 200 subscribed and open-access titles in the biological, ecological, and environmental sciences published by nonprofit societies, associations, museums, institutions, and presses.

Your use of this PDF, the BioOne Complete website, and all posted and associated content indicates your acceptance of BioOne's Terms of Use, available at [www.bioone.org/terms-of-use](https://www.bioone.org/terms-of-use).

Usage of BioOne Complete content is strictly limited to personal, educational, and non - commercial use. Commercial inquiries or rights and permissions requests should be directed to the individual publisher as copyright holder.

---

BioOne sees sustainable scholarly publishing as an inherently collaborative enterprise connecting authors, nonprofit publishers, academic institutions, research libraries, and research funders in the common goal of maximizing access to critical research.

## DEVELOPMENT AND CHARACTERIZATION OF EST-SSR MARKERS IN *STIPA BREVIFLORA* (POACEAE)<sup>1</sup>

JING REN<sup>2</sup>, ZHI-ZHEN SU<sup>2</sup>, ZHEN-HUA DANG<sup>2,4</sup>, YU DING<sup>2</sup>, PEI-XUAN WANG<sup>2</sup>, AND  
JIAN-MING NIU<sup>2,3,4</sup>

<sup>2</sup>School of Ecology and Environmental Sciences, Inner Mongolia University, Hohhot, People's Republic of China; and <sup>3</sup>Sino-US Center for Conservation, Energy, and Sustainability Science, Inner Mongolia University, Hohhot, People's Republic of China

- **Premise of the study:** *Stipa breviflora* (Poaceae) is one of the dominant species of the desert steppe in the eastern Eurasian grasslands. Simple sequence repeat (SSR) markers were developed for use in genetic diversity studies of this species.
- **Methods and Results:** A total of 1954 potentially polymorphic loci were obtained by comparing transcriptome data of eight different *S. breviflora* individuals. We selected 81 loci to verify polymorphism and 63 loci amplified, of which 21 loci exhibited polymorphism. The number of alleles per locus varied from two to 24, the observed heterozygosity ranged from 0.083 to 0.958, and the expected heterozygosity ranged from 0.396 to 0.738.
- **Conclusions:** These newly identified SSR loci can be used for population genetic and landscape genetic studies of *S. breviflora*. In addition, 14 loci also amplified in six related *Stipa* species (*S. grandis*, *S. krylovii*, *S. bungeana*, *S. aliena*, *S. gobica*, and *S. purpurea*).

**Key words:** microsatellite markers; Poaceae; polymorphism; *Stipa breviflora*; transcriptome.

*Stipa breviflora* Griseb. (Poaceae) is one of the dominant species covering the desert steppe region of the eastern Eurasian grassland. In China, it is widely distributed in the southern Mongolia Plateau, western Ordos Plateau, northwestern Loess Plateau, Tibet Plateau, and Xinjiang Province (Zhang et al., 2009). *Stipa breviflora*, which is characterized by drought resistance, grazing tolerance, fine palatability, and early spring growth, serves as an indispensable forage resource for herbivores in dryland areas. Moreover, it plays an important role in both conserving soil and water and preventing desertification (Zhang et al., 2010). *Stipa breviflora* grassland, however, has been severely degraded by global warming and anthropogenic disturbances. Despite the ecological significance of *S. breviflora*, we know little about its population genetics and evolutionary biology using current technologies. The only study that has evaluated the genetic diversity of this species used RAPD markers (Zhang et al., 2012). Microsatellite (simple sequence repeat [SSR]) markers, which can be developed using genomes or transcriptomes, are powerful tools for examining population genetic diversity. SSRs are often presumed to be neutral, but can be subject to both positive and negative selection for a variety of reasons. Although we reported preliminary results on SSR development of *S. breviflora* (Zhao et al., 2016), a systematic and improved study is also required.

<sup>1</sup>Manuscript received 17 December 2016; revision accepted 8 March 2017.

This study is supported by the National Natural Science Foundation of China (NSFC; no. 5141116).

<sup>4</sup>Authors for correspondence: jmn2005@163.com (J.M.N.), Zhdang\_1982@aliyun.com (Z.H.D.)

doi:10.3732/apps.1600157

## METHODS AND RESULTS

Transcriptome sequencing of eight *S. breviflora* individuals, collected from a wide geographic range, was conducted using Illumina HiSeq 4000 (Illumina, San Diego, California, USA). Approximately six million 150-bp reads of each individual were obtained and de novo assembled into 178,901 unigenes (>300 bp) using Trinity (Grabherr et al., 2011), with mean length of 1235 bp. SSRs were detected using MicroSATellite Identification Tool (MISA; Thiel et al., 2003), with the criteria of 12, six, five, five, four, and four repeat units for mono-, di-, tri-, tetra-, penta-, and hexanucleotide motifs, respectively. The criteria were determined by the settings for minimum number of repeats according to the MISA software instructions. We adjusted the repeat numbers of pentanucleotide and hexanucleotide motifs on the basis of our search results for a higher number of SSR loci. A total of 29,817 SSRs were identified, with trinucleotide repeats (16,785, 62.76%) being the most common, followed by dinucleotide (7489, 28.00%), mononucleotide (3071, 10.30%), hexanucleotide (958, 3.58%), pentanucleotide (923, 3.45%), and tetranucleotide (591, 2.21%) repeats.

By comparing the transcriptome sequences, we obtained 1954 unigenes likely to contain polymorphic SSR loci. Then 81 loci demonstrating significant length variation among data for eight transcriptomes were selected. Primer3 (Untergasser et al., 2012) was used to design primer pairs with lengths of 18–21 bp amplifying product sizes ranging from 90–250 bp. We initially screened 81 primer pairs using 16 *S. breviflora* individuals; 63 of these loci were successfully amplified after PCR optimization. The polymorphism of these loci was tested using 24 samples from eight populations of *S. breviflora* (three individuals per population). Finally, we obtained 21 polymorphic expressed sequence tag–SSR (EST-SSR) markers, which were deposited into GenBank (Table 1). Polymorphism of these loci was assessed using 96 individuals from eight populations (12 individuals per population). *Stipa grandis* P. A. Smirn., *S. krylovii* Roshev., *S. bungeana* Trin., *S. aliena* Keng, *S. gobica* Roshev., and *S. purpurea* Griseb. (five individuals of each population) were used to test the cross-species amplification of polymorphic markers in *S. breviflora* (Appendix 1). Owing to lack of plant specimens, voucher specimens of these species could not be provided.

Genomic DNA was extracted from frozen leaf tissues using the Plant Genomic DNA Extraction Kit (Tiangen Biotech, Beijing, China). PCR amplification was performed in a reaction mixture (25  $\mu$ L) containing 1  $\mu$ L of template

*Applications in Plant Sciences* 2017 5(4): 1600157; <http://www.bioone.org/loi/apps> © 2017 Ren et al. Published by the Botanical Society of America.

This is an open access article distributed under the terms of the Creative Commons Attribution License (CC-BY-NC-SA 4.0), which permits unrestricted noncommercial use and redistribution provided that the original author and source are credited and the new work is distributed under the same license as the original.

TABLE 1. Characteristics of 21 microsatellite primers developed for *Stipa breviflora*.

Locus	Primer sequences (5'-3')	Repeat motif	Allele size (bp)	T <sub>a</sub> (°C)	Fluorescent dye	GenBank accession no.	Putative function [Organism]	E-value
SB13	F: CTTCCTGCGAGTACAGCGATTT R: CAAACAGAGCTCAACATCACAAA	(TC) <sub>8</sub>	137	57	HEX	KY355614	PREDICTED: plasminogen activator inhibitor 1 RNA-binding protein-like [ <i>Oryza brachyantha</i> ]	4.00E-171
SB16	F: CAGTGGTTTTTGTAAACAGCAG R: GCCCGTACCAFAAATTTCTTTTT	(CAA) <sub>6</sub>	148	51	TAMRA	KY355615	Predicted protein [ <i>Hordeum vulgare</i> subsp. <i>vulgare</i> ]	1.00E-83
SB21	F: ATGTACTTGGAAAGAAACGAAGCA R: TGCTGTGTGATCTACAGGTTTG	(CGG) <sub>6</sub>	94	57	FAM	KY355616	Predicted protein [ <i>Hordeum vulgare</i> subsp. <i>vulgare</i> ]	6.00E-13
SB32	F: CAATGTAGAGGGTAAACAACGA R: AGTCAGTGTGCTGTCTGTCAAATA	(TTC) <sub>5</sub>	153	57	TAMRA	KY355617	Predicted protein [ <i>Hordeum vulgare</i> subsp. <i>vulgare</i> ]	3.00E-102
SB35	F: CTACTGACATCCACGTTATGAA R: GAGATCAGGTTTACGAACCCC	(G) <sub>16</sub>	153	57	TAMRA	KY355618	Uncharacterized protein LOC100845604 [ <i>Brachypodium distachyon</i> ]	2.00E-34
SB40	F: GATCGCATTTGGTAGTATGTAAA R: TTCCCTTCATCCCTCCACTTTG	(AG) <sub>9</sub>	160	57	FAM	KY355619	—	—
SB42	F: TCCCTCAGAAAAAATCAAACA R: ATCATCCTGTACACCGTCTGCTTA	(CT) <sub>7</sub>	138	56	HEX	KY355620	Uncharacterized protein LOC100826478 [ <i>Brachypodium distachyon</i> ]	3.00E-24
SB43	F: AAATCCTTCCTCGGCTC R: CTCATCGATCTCCCTCGCTCT	(ACC) <sub>5</sub>	152	60	TAMRA	KY355621	Uncharacterized protein LOC101756208 [ <i>Setaria italic-d</i> ]	9.00E-100
SB45	F: CCGACACACAAAGATGAGC R: GCTGFGCAGGACCTCCC	(GCG) <sub>5</sub>	111	51	FAM	KY355622	Uncharacterized protein LOC100823613 [ <i>Brachypodium distachyon</i> ]	3.00E-53
SB46	F: TCCTCTCTGTATATAAAGCCG R: ATGCAFTTGGCTGGAATGTT	(CT) <sub>8</sub>	136	56	HEX	KY355623	PREDICTED: NAC domain-containing protein 78 [ <i>Brachypodium distachyon</i> ]	0
SB49	F: ACTCTCTGCAACTCTGTGAAAG R: TAATGGAAGCATTTGGCTAFACA	(GA) <sub>9</sub>	144	56	TAMRA	KY355624	—	—
SB50	F: AGAGCATCATCTTTGTCTC R: ACCGACTTATCTCTCTTTCTTT	(GAG) <sub>6</sub>	134	57	HEX	KY355625	PREDICTED: paramyosin-like isoform X1 [ <i>Brachypodium distachyon</i> ]	2.00E-122
SB52	F: AGAAGAAGGAAGAAGACCCA R: AGATCCACCGCTCTTCCTAGT	(GGC) <sub>6</sub>	146	57	TAMRA	KY355626	Uncharacterized protein LOC101770650 [ <i>Setaria italic-d</i> ]	5.00E-78
SB53	F: GCAAAGGAACCTACGCTCTCC R: GAGAGGCTCATATGGCTGAAC	(GGC) <sub>6</sub>	133	59	HEX	KY355627	PREDICTED: cysteine-rich receptor-like protein kinase 10 [ <i>Brachypodium distachyon</i> ]	2.00E-147
SB54	F: CACAAGGTACCGAAAAGGAAG R: ACCAACCCACTCTCTCTCTCT	(GA) <sub>6</sub>	90	59	FAM	KY355628	Hypothetical protein EUGRSUZ_C01942, partial [ <i>Eucalyptus grandis</i> ]	5.00E-11
SB55	F: AAATCTGCTCAGGTGGAATC R: AAATCAATCGCACTCGCAAT	(TGCGAT) <sub>4</sub>	131	54	HEX	KY355629	PREDICTED: heat shock protein 82 [ <i>Brachypodium distachyon</i> ]	2.00E-13
SB57	F: AACTTGTGAAGTTTGCAATGTC R: AACCCAGTCACTCTGACAACTA	(AT) <sub>10</sub>	148	57	TAMRA	KY355630	Predicted protein [ <i>Hordeum vulgare</i> subsp. <i>vulgare</i> ]	1.00E-37
SB77	F: ACTTTATCCGCAATGCTA R: TTCGTTCTTTTGTCTGTG	(AAG) <sub>5</sub>	125	56	FAM	KY355631	Predicted protein [ <i>Hordeum vulgare</i> subsp. <i>vulgare</i> ]	0
SB78	F: TCACCATACCCATTCGTTTCCT R: TCATCTTCGGATCTCTCTCC	(GCC) <sub>5</sub> TTG(GA) <sub>10</sub>	191	60	FAM	KY355632	PREDICTED: ATP synthase subunit d, mitochondrial-like [ <i>Setaria italic-d</i> ]	6.00E-88
SB79	F: GATGTFCCACTCATCCAGGCTGT R: GTGCGTGAAGAAAGAACGGTCC	(TC) <sub>9</sub>	235	54	FAM	KY355633	RecName: Full = Thioredoxin H-type; Short = Trx-H; AltName: Full = TrxTa [ <i>Triticum aestivum</i> ]	6.00E-54
SB81	F: CGCTCCACTACCTTTTCGTATCAC R: GGAATGAATGCCCTTGAGTGAGTC	(CT) <sub>11</sub> (GT) <sub>8</sub>	194	60	FAM	KY355634	Uncharacterized protein LOC100845363 [ <i>Brachypodium distachyon</i> ]	9.00E-137

Note: T<sub>a</sub> = annealing temperature.

TABLE 2. Results of initial screening of 21 polymorphic loci identified in eight populations of *Stipa breviflora*.<sup>a</sup>

Locus	Wulate (N = 12)			Junggar (N = 12)			Gaolian (N = 12)			Hainan (N = 12)			Alxa (N = 12)			Chifeng (N = 12)			Putian (N = 12)			Hejing (N = 12)			Total		
	A	H <sub>o</sub>	H <sub>e</sub> <sup>b</sup>	A	H <sub>o</sub>	H <sub>e</sub> <sup>b</sup>	A	H <sub>o</sub>	H <sub>e</sub> <sup>b</sup>	A	H <sub>o</sub>	H <sub>e</sub> <sup>b</sup>	A	H <sub>o</sub>	H <sub>e</sub> <sup>b</sup>	A	H <sub>o</sub>	H <sub>e</sub> <sup>b</sup>	A	H <sub>o</sub>	H <sub>e</sub> <sup>b</sup>	A	H <sub>o</sub>	H <sub>e</sub> <sup>b</sup>	A	H <sub>o</sub>	H <sub>e</sub> <sup>b</sup>
SB13	4	0.08	0.57***	3	0.08	0.54**	4	0.17	0.68***	3	0.00	0.50***	4	0.167	0.413*	2	0.083	0.080	1	0.000	0.000 <sup>M</sup>	3	0.083	0.434**	7	0.083	0.401
SB16	5	0.83	0.68	5	0.73	0.75	6	0.67	0.74	6	0.83	0.78	5	0.583	0.774	3	0.500	0.403	3	1.000	0.622**	2	0.917	0.497*	5	0.758	0.655
SB21	3	0.75	0.66***	4	0.67	0.69***	4	0.75	0.69*	4	0.55	0.67*	4	0.667	0.705**	5	0.833	0.694**	2	1.000	0.500**	3	1.000	0.594**	2	0.777	0.649
SB32	7	0.67	0.75	6	0.17	0.70***	8	0.75	0.75	11	0.67	0.81	5	0.333	0.722**	8	0.583	0.757*	3	0.182	0.169	3	0.455	0.517	22	0.475	0.647
SB35	4	0.17	0.66***	10	0.08	0.87***	8	0.33	0.81***	11	0.42	0.85***	9	0.455	0.855***	6	0.167	0.656***	3	0.167	0.156	5	0.083	0.698***	18	0.234	0.695
SB40	7	0.33	0.72***	8	0.25	0.78***	4	0.25	0.52**	4	0.00	0.65***	6	0.083	0.726***	6	0.273	0.653***	3	0.091	0.288	5	0.083	0.649***	13	0.180	0.623
SB42	7	0.25	0.77***	6	0.00	0.78***	7	0.25	0.78***	5	0.17	0.72***	5	0.000	0.736***	5	0.417	0.476	2	0.091	0.087	5	0.333	0.747***	15	0.188	0.636
SB43	3	0.25	0.34	6	0.33	0.48**	6	0.83	0.70	7	0.75	0.62	3	0.167	0.156	2	0.333	0.278	2	0.083	0.080	6	0.583	0.569	12	0.417	0.403
SB45	2	0.91	0.50*	2	0.75	0.47	2	1.00	0.50**	2	0.42	0.33	2	0.583	0.413	2	0.750	0.469	2	1.000	0.500**	2	0.083	0.080	11	0.687	0.407
SB46	11	0.67	0.88*	9	0.50	0.80***	9	0.75	0.71	11	1.00	0.87	7	0.667	0.781	4	0.750	0.538	5	0.917	0.625	4	0.917	0.628	10	0.771	0.729
SB49	8	0.33	0.82***	6	0.25	0.65***	5	0.36	0.74*	8	0.42	0.84***	8	0.250	0.844***	4	0.250	0.538**	3	0.250	0.226	3	0.167	0.635***	6	0.285	0.661
SB50	7	0.73	0.79	7	0.92	0.69	5	0.92	0.76	6	0.75	0.73*	9	0.750	0.712	5	0.833	0.642	3	0.833	0.542	4	1.000	0.677	8	0.841	0.692
SB52	8	0.67	0.73	7	0.83	0.79	7	0.92	0.80	9	0.75	0.77	6	0.917	0.792	7	0.667	0.701	4	0.583	0.451	5	0.833	0.632	17	0.771	0.708
SB53	7	0.92	0.81*	4	0.83	0.64	5	0.92	0.74	6	0.82	0.76**	6	0.818	0.814	6	0.917	0.819**	3	0.333	0.288	7	0.917	0.799**	20	0.809	0.709
SB54	8	0.83	0.74	5	1.00	0.67	6	1.00	0.81***	6	1.00	0.82*	8	0.833	0.774	7	1.000	0.722***	5	1.000	0.628*	2	1.000	0.500**	16	0.958	0.708
SB55	5	0.92	0.70	4	1.00	0.73	5	0.83	0.67	6	0.58	0.69	5	0.833	0.667	5	1.000	0.704	1	0.000	0.000 <sup>M</sup>	3	1.000	0.611**	12	0.771	0.596
SB57	9	0.75	0.82	7	0.42	0.48	9	0.67	0.84*	9	0.73	0.85*	8	1.000	0.726	8	0.583	0.698*	4	0.333	0.295	7	0.500	0.663*	11	0.622	0.671
SB77	5	0.58	0.63	5	0.73	0.73	5	0.75	0.60	6	0.92	0.78	6	0.833	0.688	4	0.917	0.705	3	0.833	0.517	3	0.917	0.538*	19	0.810	0.647
SB78	4	0.27	0.68**	5	0.08	0.57***	2	0.17	0.15	6	0.08	0.72***	4	0.000	0.514	4	0.083	0.295**	2	0.083	0.080	2	0.000	0.153*	11	0.097	0.396
SB79	8	0.67	0.83	4	0.92	0.74	7	0.92	0.80	13	1.00	0.85	10	0.917	0.837	8	0.917	0.753*	3	0.833	0.517	3	1.000	0.580*	18	0.896	0.738
SB81	6	0.42	0.64*	8	0.33	0.82***	8	0.67	0.67	14	0.92	0.90	7	0.583	0.778	7	0.583	0.705	7	0.545	0.459	6	0.417	0.476	24	0.558	0.681

Note: A = total number of alleles; H<sub>e</sub> = expected heterozygosity; H<sub>o</sub> = observed heterozygosity; N = total number of samples analyzed.

<sup>a</sup> Locality information is provided in Appendix 1.

<sup>b</sup> Asterisks indicate significant deviation from Hardy–Weinberg equilibrium (\*P < 0.05, \*\*P < 0.01, \*\*\*P < 0.001); M = monomorphic.

TABLE 3. Cross-species amplification results of 21 polymorphic EST-SSR loci developed for *Stipa breviflora* in six other *Stipa* species.<sup>a</sup>

Locus	<i>S. grandis</i>	<i>S. aliena</i>	<i>S. bungeana</i>	<i>S. gobica</i>	<i>S. krylovii</i>	<i>S. purpurea</i>
SB13	1	1	1	1	1	1
SB16	1	1	1	1	1	1
SB21	1	1	1	1	1	1
SB32	1	1	1	1	1	1
SB35	0	0	0	1	0	0
SB40	1	1	1	1	1	1
SB42	1	1	1	1	1	1
SB43	1	0	1	1	1	0
SB45	0	0	1	0	0	1
SB46	1	1	1	1	1	1
SB49	1	1	1	1	1	1
SB50	0	1	1	1	1	1
SB52	0	0	1	0	0	0
SB53	1	1	1	1	1	1
SB54	1	1	1	1	1	1
SB55	0	0	1	1	0	0
SB57	1	1	1	1	0	0
SB77	1	1	1	1	1	1
SB78	1	1	1	1	1	1
SB79	1	1	1	1	1	1
SB81	1	1	1	1	1	1

Note: 1 = successful amplification; 0 = failed amplification.

<sup>a</sup>Locality information is provided in Appendix 1.

DNA (30–40 ng/μL), 0.5 μL (10 pM) of each primer, 12.5 μL of Premix *Taq* (TaKaRa Biotechnology Co., Dalian, Liaoning Province, China), and 10.5 μL of ddH<sub>2</sub>O. Conditions for PCR amplification were as follows: 4 min at 94°C; 35 cycles of 30 s at 94°C, 30 s at a primer-specific annealing temperature (Table 1), 30 s at 72°C; and a final extension step at 72°C for 10 min. PCR products were first detected by 1.5% agarose gel electrophoresis to check for successful amplification. Forward primers for the 21 successfully amplified polymorphic loci were labeled with one of three different fluorescent dyes (FAM, HEX, or TAMRA) and used for amplifications with the same protocol. The labeled PCR products were analyzed on an ABI 3730 DNA Analyzer with a GeneScan 500 LIZ Size Standard (Applied Biosystems, Beijing, China). Allele sizes were called using GeneMarker version 2.6.0 (SoftGenetics, State College, Pennsylvania, USA). Number of alleles per locus, observed heterozygosity, and expected heterozygosity were calculated using GenAlEx version 6.5 (Peakall and Smouse, 2012). GENEPOP version 4.42 (Rousset, 2008) was used to measure the departure from Hardy–Weinberg equilibrium and linkage disequilibrium, and MICRO-CHECKER version 2.2.3 (van Oosterhout et al., 2004) was used to check the possibility of null alleles.

Among these 21 polymorphic loci, the total number of alleles per locus ranged from two to 24, observed heterozygosity ranged from 0.083 to 0.958, and expected heterozygosity ranged from 0.396 to 0.738. Only locus SB21 with null alleles consistently departed from Hardy–Weinberg equilibrium in all populations, and we did not detect linkage disequilibrium between any loci (Table 2). Cross-species amplification of the 21 polymorphic markers was tested in six related species, namely *S. grandis*, *S. krylovii*, *S. bungeana*, *S. aliena*, *S. gobica*, and *S. purpurea* (five individuals per population). Fourteen (66.67%) of the 21 loci successfully amplified in all tested species; the remaining loci were amplified in some species (Table 3).

## CONCLUSIONS

This is the first known report of 21 polymorphic EST-SSRs for *S. breviflora*. These SSRs will be used to evaluate impacts of isolation by distance and recent habitat fragmentation on the genetic diversity and structure of *S. breviflora* populations. These

SSRs may also be used in investigations of genetic diversity of other *Stipa* L. species. Hodel et al. (2016) indicate that microsatellites generated from transcriptomes could likely be found in translated regions of the genome. The majority of loci favored in translated regions are trinucleotide repeats (Hodel et al., 2016). Markers occurring in or near coding regions are prone to selective pressures (Morgante et al., 2002), which casts doubt on the application of microsatellites in genetic diversity analysis. However, one study about *Glycine* Willd. and *Oenothera* L. demonstrates that many trinucleotide repeats linked closely to translated regions are not themselves within a translated region of a gene (Hodel et al., 2016); therefore, these loci are potentially useful. In our study, 21 SSRs were derived from transcriptome data, and a significant number of these loci (11/21) were or contained trinucleotide or hexanucleotide repeats. Therefore, we suggest that all 21 loci could be used to examine genetic diversity while neutrality is tested. Researchers investigating selection should use the 11 loci with trinucleotide repeats.

## LITERATURE CITED

- GRABHERR, M. G., B. J. HAAS, M. YASSOUR, J. Z. LEVIN, D. A. THOMPSON, I. AMIT, X. ADICONIS, ET AL. 2011. Full-length transcriptome assembly from RNA-Seq data without a reference genome. *Nature Biotechnology* 29: 644–652.
- HODEL, R. G. J., M. A. GITZENDANNER, C. C. GERMAIN-AUBREY, X. LIU, A. A. CROWL, M. SUN, J. B. LANDIS, ET AL. 2016. A new resource for the development of SSR markers: Millions of loci from a thousand plant transcriptomes. *Applications in Plant Sciences* 4: 1600024.
- MORGANTE, M., M. HANAFEY, AND W. POWELL. 2002. Microsatellites are preferentially associated with nonrepetitive DNA in plant genomes. *Nature Genetics* 30: 194–200.
- PEAKALL, R., AND P. E. SMOUSE. 2012. GenAlEx 6.5: Genetic analysis in Excel. Population genetic software for teaching and research—An update. *Bioinformatics (Oxford, England)* 28: 2537–2539.
- ROUSSET, F. 2008. GENEPOP'007: A complete re-implementation of the GENEPOP software for Windows and Linux. *Molecular Ecology Resources* 8: 103–106.
- THIEL, T., W. MICHALEK, R. K. VARSHNEY, AND A. GRANER. 2003. Exploiting EST databases for the development and characterization of gene-derived SSR-markers in barley (*Hordeum vulgare* L.). *Theoretical and Applied Genetics* 106: 411–422.
- UNTERGASSER, A., I. CUTCUTACHE, T. KORESSAAR, J. YE, B. C. FAIRCLOTH, M. REMM, AND S. G. ROZEN. 2012. Primer3—New capabilities and interfaces. *Nucleic Acids Research* 40: e115.
- VAN OOSTERHOUT, C., W. F. HUTCHINSON, D. P. M. WILLS, AND P. SHIPLEY. 2004. MICRO-CHECKER: Software for identifying and correcting genotyping errors in microsatellite data. *Molecular Ecology Notes* 4: 535–538.
- ZHANG, Q., J. M. NIU, J. J. DONG, Y. DING, F. L. WANG, J. GUO, AND L. Q. ZHAO. 2009. Floristic study of spermatophyte on *Stipa breviflora* steppe distributed in Inner Mongolia. *Journal of Desert Research* 29: 451–456.
- ZHANG, Q., J. M. NIU, Y. DING, S. KANG, AND J. J. DONG. 2010. Current research advances and future prospects of biology and ecology of *Stipa breviflora*. *Chinese Journal of Grassland* 32: 93–101.
- ZHANG, Q., J. M. NIU, S. B. WU, AND J. J. DONG. 2012. Impact of climatic factors on genetic diversity of *Stipa breviflora* populations in Inner Mongolia. *Genetics and Molecular Research* 11: 2081–2093.
- ZHAO, L., Z. H. DANG, Y. N. ZHAO, Y. ZHANG, J. M. NIU, J. REN, AND J. M. ZHOU. 2016. Optimization of SSR-PCR reaction system and primers screening in *Stipa breviflora*. *Journal of Inner Mongolia University* 47: 58–64.

APPENDIX 1. Location information for populations of *Stipa breviflora* and six other *Stipa* species used in this study.<sup>a</sup>

Species	Collection locality	Altitude (m)	Geographic coordinates	<i>N</i>
<i>Stipa breviflora</i> Griseb.	Wulate Middle Banner, Inner Mongolia	1321	41°29'28.63"N, 108°57'17.64"E	12
<i>S. breviflora</i>	Jungar, Inner Mongolia	1160	39°46'8.43"N, 110°57'25.91"E	12
<i>S. breviflora</i>	Gaolan, Gansu	1784	36°15'55.94"N, 103°48'44.18"E	12
<i>S. breviflora</i>	Hainan, Qinghai	2956	36°18'59.32"N, 100°33'43.37"E	12
<i>S. breviflora</i>	Alxa, Inner Mongolia	1461	39°50'39.65"N, 105°03'34.30"E	12
<i>S. breviflora</i>	Chifeng, Inner Mongolia	576	42°38'22.48"N, 119°12'14.24"E	12
<i>S. breviflora</i>	Pulan, Tibet	3863	30°16'33.28"N, 81°10'11.19"E	12
<i>S. breviflora</i>	Hejing, Xinjiang	2141	42°54'05.90"N, 86°17'56.83"E	12
<i>S. aliena</i> Keng	Haibei, Qinghai	3201	101°19'31.3"N, 37°36'38.9"E	5
<i>S. bungeana</i> Trin.	Hohhot, Inner Mongolia	1040	40°47'24.11"N, 111°28'7.28"E	5
<i>S. gobica</i> Roshev.	Wuhai, Inner Mongolia	1599	39°37'51.50"N, 106°53'43.77"E	5
<i>S. grandis</i> P. A. Smirn.	Manzhouli, Inner Mongolia	659	49°33'01"N, 117°33'59"E	5
<i>S. krylovii</i> Roshev.	Hulunbuir, Inner Mongolia	930	47°51'58"N, 115°46'48"E	5
<i>S. purpurea</i> Griseb.	Tianjun, Qinghai	3195	37°12'40.2"N, 98.55'31.4"E	5

Note: *N* = numbers of individuals sampled.

<sup>a</sup>Voucher specimens were not collected at the time of the study.