

## **Morphological Variability of the Paratethyan Oligocene-Miocene Small Reticulofenestrid Coccolites and Its Paleocological and Paleogeographical Implications**

Author: Holcová, Katarína

Source: Acta Palaeontologica Polonica, 58(3) : 651-668

Published By: Institute of Paleobiology, Polish Academy of Sciences

URL: <https://doi.org/10.4202/app.2009.0006>

---

BioOne Complete ([complete.BioOne.org](https://complete.BioOne.org)) is a full-text database of 200 subscribed and open-access titles in the biological, ecological, and environmental sciences published by nonprofit societies, associations, museums, institutions, and presses.

Your use of this PDF, the BioOne Complete website, and all posted and associated content indicates your acceptance of BioOne's Terms of Use, available at [www.bioone.org/terms-of-use](https://www.bioone.org/terms-of-use).

Usage of BioOne Complete content is strictly limited to personal, educational, and non-commercial use. Commercial inquiries or rights and permissions requests should be directed to the individual publisher as copyright holder.

---

BioOne sees sustainable scholarly publishing as an inherently collaborative enterprise connecting authors, nonprofit publishers, academic institutions, research libraries, and research funders in the common goal of maximizing access to critical research.

# Morphological variability of the Paratethyan Oligocene–Miocene small reticulofenestrid coccolites and its paleoecological and paleogeographical implications

KATARÍNA HOLCOVÁ



Holcová, K. 2013. Morphological variability of the Paratethyan small reticulofenestrid coccolites and its paleoecological and paleogeographical implications. *Acta Palaeontologica Polonica* 58 (3): 651–668.

The analysis of size changes (length of placoliths, their width, length of central opening and its width) in elliptical reticulofenestrids from the NP25–NN5 zonal interval of the Central Paratethys allowed to discriminate two size categories of placoliths: (i) small *Reticulofenestra minuta* (< 3.5  $\mu\text{m}$ ); (ii) *Reticulofenestra haqii–pseudoumbilicus* group (4–10.0  $\mu\text{m}$ ). The latter group appeared for the first time (FO, first occurrence) in the upper Egerian (size 4–7  $\mu\text{m}$ ) with the size of placoliths in this plexus increasing gradually. The FO of *R. pseudoumbilicus* > 8.0  $\mu\text{m}$  has been established in the Zone NN2 while its FCO (first common occurrence) in the Zone NN5. This study shows that the FOs of size-defined morphotypes of the *R. haqii–pseudoumbilicus* group differ in the Central Paratethys and oceanic realm. Blooms of *R. minuta* at the Oligocene–Miocene boundary and in the Early Middle Miocene may be correlated with the incoming of warm water into the higher latitude Central Paratethys basins during connection with the Mediterranean Sea. Transgression favored the expansion of near-shore areas associated probably by some short-time oscillations of salinity. The FO of *R. haqii–pseudoumbilicus* group and the FO of *R. pseudoumbilicus* > 8.0  $\mu\text{m}$  can be correlated with the opening of new pathways between the Mediterranean and the Central Paratethys. Gradual size changes in the *R. haqii–pseudoumbilicus* group probably reflect climatic changes: the decrease of coccolith size in the late Egerian may reflect cooling (Mi1 event) while the increase in coccolith size in the interval from the FO of *Helicosphaera ampliapertura* to the FO of *Sphenolithus heteromorphus* occurred due to warming. Two size categories of placoliths in the *R. haqii–pseudoumbilicus* group (3.5–6  $\mu\text{m}$  and 6–8  $\mu\text{m}$ ) recorded in the interval from Zone NN1 to the lower part of the Zone NN2 may represent seasonal populations.

Key words: Calcareous nannofossils, size changes, paleoecology, Oligocene, Miocene, Central Paratethys.

Katarína Holcová [holcova@natur.cuni.cz], Department of Geology and Paleontology, Charles University in Prague, Albertov 6, 128 43 Praha 2, Czech Republic.

Revised version received 6 September 2011, accepted 31 December 2011, available online 12 January 2012.

Copyright © 2013 K. Holcová. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

## Introduction

The collision of the African/Apulia/Arabian plates and Eurasian continent began during the Eocene, and resulted in the uplift and emergence of the Alpine chains and the break-up of the Tethyan Realm into the Mediterranean and Paratethyan domains (Popov et al. 2004). This resulted in the differentiation of the Paratethys as a biogeographic entity. The first endemic Paratethyan assemblages of mollusks, calcareous nanoplankton and foraminifera appeared in the Oligocene (Báldi 1986). The subsequent history of the Paratethys involved periods of isolation from the adjacent Mediterranean Sea and Indian Ocean during which endemic biota evolved, and periods of oceanic communication during exchanges occurred with the faunas of the Mediterranean and the Indo-Pacific provinces. Additionally episodic paleogeographic

changes, global climatic variations, local oscillations of salinity, oxygen content and globally and locally affected sea-level changes also influenced the evolution of Paratethyan ecosystems (Rögl 1998; Kováč 2000; Popov et al. 2004; Piller et al. 2007; etc.). Interactions of the local geodynamic and the global climatic factors lead to major problems in their biostratigraphical, paleoecological and paleogeographical interpretation. As a consequence of periodical paleogeographic changes affecting them, the morphological changes (size) of Central Paratethys populations may be local, differing those from contemporaneous oceanic populations.

Coccolithophorids were widespread autotrophs in marine Paratethyan ecosystems and are of great biostratigraphical importance. Besides *Coccolithus pelagicus*, reticulofenestrids are the most common components of the assemblages and their detailed study can provide important data for reconstruct-




Fig. 1. A. Sketch of the main Middle European geological units (after Kováč 2000). B. Moravian part of the Carpathian Foredeep with location of borehole PY-1 near village Přemyslovice; and section Kralice (KRA), geological situation according Chlupáč et al. (2002). C. Eastern part of the South Slovak Basin with location of boreholes LKS-1, BE-2, FV-1, EH-1, and EH-2 drilled by State Geological Institute of Dionýz Štúr; boreholes LR-9 and LR-10 drilled by “Geologický prieskum n.p.” regional centrum Rožňava and section Lipovany (LIP). Geological situation after Halásová et al. (1996).

ing the factors that have induced morphologic changes in calcareous nannoplankton. *Reticulofenestra bisecta*, *R. abisecta* and *Cyclicargolithus floridanus* are common in the late Oligocene while in the lower and early Middle Miocene, reticulofenestrids are represented by *Reticulofenestra minuta*, *R. haqii*, and *R. pseudoumbilicus* together with endemic *R. excavata* (Lehotayová 1975, 1984; Marunteanu 1992; Andreyeva-Grigorovich et al. 1997; Holcová 2005). The aim of this study is to show the relationship between size changes in reticulofenestrids and paleogeographic and paleoecologic evolution in an intracontinental basin.

**Abbreviations.**—FCO, first common occurrence; FO, first occurrence; LCO, last common occurrence; LO, last occurrence.

## Geological setting

The area of the Central Paratethys studied here includes the Pannonian Basin system and the Carpathian Foreland basins (Seneš 1961).

The majority of samples were taken from the South Slovak Basin (Fig. 1). Sediments filling of this geomorphological unit were deposited in the northern part of the Buda Basin during the Late Oligocene (Kiscellian Stage) to the Chattian and lower Aquitanian (Egerian Stage), in the Filakovo/Péteřvářa Basin during the upper Aquitanian and lower Burdigalian (Eggenburgian Stage) and in the Novohrad/Nógrád basin in the upper Burdigalian and Langhian (Ottngian, Karpatian, and early Badenian; Vass 1996). These three basins form the northern part of the Pannonian Basin system. Biostratigraphy, lithostratigraphy and sedimentology of this area were summarized in Vass et al. (1979, 1989, 2007). Lithostratigraphic units have been defined by Vass and Elečko (1982) and correlation with standard nannoplankton zones (Martini 1971) has been established by Lehotayová (1982). The paleogeographical maps for every Central Paratethys stage have been produced by Vass et al. (1979, 1989, 2007). Vass et al. (1993), Márton et al. (1995), and Vass (1996) delineated the most important tectonic events and local sea-level changes; the latter by have been correlated with the global sea level curve of Haq et al. (1988) by Vass (1995).

The Middle Miocene samples have been collected from sections in the Carpathian Foredeep. These sections are described in detail by Zágoršek et al. (2008, 2009) and Zágoršek and Holcová (2009; Fig. 1).

The study interval from the Oligocene/Miocene boundary interval to the lower Middle Miocene (zones NP25–NN5) corresponds to the Egerian, Eggenburgian, Ottngian, Karpatian, and Lower Badenian stages of the Central Paratethys (Popov et al. 2004; Harzhauser and Piller 2007; Hohenegger et al. 2009).




Fig. 2. Lithology, stratigraphical ranges, and sampled interval of studied sections. Location of boreholes LR-10, LR-9, LR-2, C-2, FV-1, EH-1, EH-2, D-19, LKŠ-1, BE-2, PY-1, and sections Lipovany and Kralice II is illustrated on Fig. 1.



## Material and methods

A total of 49 samples taken from 13 sections have been studied. The set of samples were chosen from about 500 of samples analysed in the study area in previous years (e.g., Holcová 2001, 2005). Nannofossils in selected samples were common to abundant (10–30 specimens in visual field of microscope) and well preserved without apparent diagenetic changes (dissolution, recrystallization). Relative abundances of reticulofenestrids in studied samples varied from 10 to 30% with exception of samples from the zones NN1 and NN5 with more than 50% of reticulofenestrids. Besides Langhian samples (Zone NN5) were dominated by *Coccolithus pelagicus* in the studied assemblages. The location of the sections, their lithology and stratigraphical ranges are summarized in Figs. 1 and 2.

Calcareous nannofossils were examined and photographed using a light microscope and 1000× magnification with both bright field and crossed polarized light. Smear slides were prepared from about 1 cc of rock sample (claystone, siltstone, and fine sandstone) using the method of Švábenická (2002). The first 50 specimens of reticulofenestrids were measured in every sample. Placoliths of *Reticulofenestra* spp. were measured from the digital microphotographs. In total, 2447 specimens were analyzed. The following characters were measured for each placolith: length (the largest diameter of placoliths), width (diameter of placolith perpendicular to the length of the placolith), length of central opening (along the length of the placolith) and its width (across the width of the placolith) (Fig. 3).

The measurements were analyzed by simple statistical methods using STATISTICA software. Frequency histograms were used for the first graphical evaluation of the distribution type of a simple biometric parameter. XY-plots and the Pearson correlation coefficient were used for evaluation of correlations between two parameters.

The size-dependent classification of *Reticulofenestra minuta*–*haqii*–*pseudoumbilicus* group (Young 1999) has been used as an initial taxonomic concept (*Reticulofenestra minuta*: < 3 µm, *R. haqii*: 3–5 µm, and *R. pseudoumbilicus* > 5 µm). This classification is broadly accepted also in the Central Para-




Fig. 3. Dimensions measured in placoliths. Abbreviations: lo, length of central opening; lp, length of placoliths; wo, width of central opening; wp, width of placoliths.

tethys (Ćorić and Švábenická 2004; Tomanová Petrová and Švábenická 2007; Ćorić and Hohenegger 2008). However, the small-sized placoliths identified like *Reticulofenestra minuta* may represent two or more species which cannot be distinguished under the light microscope (Haq 1980).

To analyze size changes in reticulofenestrids over time, the studied interval was divided into seven intervals on the basis of six calcareous nannoplankton events. The age of these bioevents in the Central Paratethys may differ from their age in the world ocean (Berggren et al. 1995; Lourens et al. 2004) because new taxa were only able to migrate into the Central Paratethys when communication existed between them.

(i) The FO of *Helicosphaera carteri* was recorded within Zone NP 25 in the study area (Holcová 2005).

(ii) The LCO of *Reticulofenestra bisecta*. The extinction of this species is often used to approximate the Oligocene/Miocene boundary (Berggren et al. 1995; Young 1999). However, in the Buda Basin, rare specimens of *R. bisecta* occur in zones NN1 and NN2 (Holcová 2005). These are probably reworked specimens and therefore only the LCO of *R. bisecta* was used to characterize the NP25/NN1 zonal boundary.

(iii) The FO of *Helicosphaera scissura* and the FO of *Discoaster druggii* are isochronous events in the Central Paratethys (Holcová 2005). *Helicosphaera scissura* is, however, more abundant and its FO is more easily determined.

(iv) The FO of *H. ampliaptera* was recorded in Zone NN2 in the Central Paratethys (Marunteanu 1992; Holcová 2002, 2005).

(v) The FO of *S. belemnus* was established to lie at the NN2/NN3 boundary in the Central Paratethys (Marunteanu 1992, Andreyeva-Grigorovich et al. 1997).

(vi) The FO of *S. heteromorphus*/LO of *S. belemnus*: In the Central Paratethys both species briefly co-occur.

(vii) For comparison reticulofenestrids from Zone NN5 were studied. This zone is characterized by the absence of *H. ampliaptera* and the presence of *S. heteromorphus* (Martini 1971).

Seven samples were intended to analyze from every interval. However, from the LO of *Helicosphaera ampliaptera* to the FO of *Sphenolithus heteromorphus* (upper part of the Zone NN2 and Zone NN3), terrigenous sedimentation dominated in the study area. Therefore only limited number of samples with well preserved and abundant nannofossils was found in this interval. On the other hand, longer intervals with variegated marine facies (zones NP25 and NN4) were sampled to more detail.

## Results

### Morphologic variability of placoliths

**Size and length/width-ratio of placoliths and size of central opening.**—The length of the placoliths varies from 1.6 to 10 µm, their width from 1 to 9 µm. The length-width scatter plot and the correlation coefficient (0.95) between the length




Fig. 4. Scatter plots and frequency histograms visualizing correlations between biometric characteristics of the placoliths. **A.** Length/width-ratio of placoliths. **B.** Length/width-ratio of central opening. **C.** Length of placolith/length of central opening.

<http://dx.doi.org/10.4202/app.2009.0006>

and width of the placoliths indicate a very high correlation between these two parameters (Fig. 4A). The distribution of these characters is polymodal and shows three groups that are also distinct on the width-length plot (Fig. 4A): (i) small placoliths (1.6–3  $\mu\text{m}$ ); (ii) medium-size placoliths (3.5–8  $\mu\text{m}$ ), and (iii) large placoliths (8–10  $\mu\text{m}$ ). The small placoliths constitute a distinct size group, but transitional size categories occur between the medium-size and large placoliths.

#### Relative size and length/width-ratio of central opening.—

The size of the central opening varied from 0.35 to 3.5  $\mu\text{m}$ . The correlation between the length and width of the central opening is high (0.83) with two groups in which the length are (i) 0.35–1  $\mu\text{m}$  and (ii) 1–3.5  $\mu\text{m}$  (Fig. 4B).

The relative size of the central opening was expressed as a ratio between the size of the placolith and the size of the central opening (Fig. 4C). The ratio varies from 0.15 to 0.48, and shows a polymodal distribution with three groups: (i) placoliths with narrow central opening (size of placolith/size of central opening varies from 0.15 to 0.24), the placoliths with narrow central opening are small and reach maximum diameter 7.5  $\mu\text{m}$ ; (ii) placoliths with medium central opening (size of placolith/size of central opening from 0.25 to 0.40) were recorded in all size categories of placoliths; (iii) placoliths with wide central opening (size of placolith/size of central opening from 0.40 to 0.48) were not recorded among the smallest placoliths (1.6–3  $\mu\text{m}$ ).

#### Size changes over time

Correlations between the parameters analyzed here for placoliths and the age of samples were investigated on the basis of correlation coefficient. A negative correlation was recorded between the length of the placoliths ( $p = 0.01$ ) and their age. The shift in the total size of placoliths and that of their central opening over time is illustrated in Figs. 5–8 for individual samples. A summary of the data is showed in Fig. 9. From these data we may conclude the following:

(i) Small placoliths (1.6–3  $\mu\text{m}$ ) constitute a well separated group in all assemblages, independent of age. However, they are abundant in two intervals: (1) in the lowest Miocene (Zone NN1 and lower part of Zone NN2 below the FO of *H. ampliapertura*), in these intervals they account for 30–70% of the nannoplankton assemblages; (2) in the lower Middle Miocene (Zone NN5), where they account for 50–95% of the assemblages.

(ii) The size of the oldest placoliths in the *R. haqii*–*pseudoumbilicus* group (upper Zone NP25) varies from 4.0–7.0  $\mu\text{m}$ . Than the size of placoliths in the group slightly increased though this trend may differ in some individual samples (Figs. 5, 6). In the Zone NN3, the size varies from 4.0–8.0  $\mu\text{m}$ .

(iii) Bimodal size distribution was recorded firstly during Zone NN1; it dominates in the lower part of the NN2 Zone (Figs. 9, 10).

(iv) Wide variations in the relative size of the central opening characterize interval from zones NP25 to lower part of NN2; generally, the relative size of the central open-

ing increases (Fig. 9). At the level of the FO of *H. ampliapertura*, larger *R. haqii* (6–7  $\mu\text{m}$ ) with small central opening (1.7–2.0  $\mu\text{m}$ ) appeared, and relative size of central opening at this level decreased (Fig. 9).

(v) Broad variability of size of placolith as well as of the central opening characterizes the assemblages in Zone NN4 (Fig. 9).

(vi) Large placoliths (> 8  $\mu\text{m}$ ) occur first at the Zone NN5 (Fig. 9).

## Discussion

### Taxonomic implications

The size of placoliths is commonly used for classification of *Reticulofenestra* species. Using principal component analysis, Backman (1980) showed that the majority of morphological variation in *Reticulofenestra* can be explained by variation in the size of placoliths. However, the degree of closure of the central area can also be used for classification of *Reticulofenestra* (Backman 1978, 1980; Pujos 1985). Young (1990) suggested that this latter character is an unstable ecophenotypic character, and based his classification of the Late Miocene–Pliocene *Reticulofenestra* mainly on the length of the placoliths.

Young (1999) also differentiated the Early Miocene *R. minuta*–*haqii*–*pseudoumbilicus* group on the basis of overall size of the placoliths. The taxonomic difference between *R. minuta* and *R. haqii* was based on a superior limit size of 3  $\mu\text{m}$  for the former. This criterion is well applicable to the reticulofenestrids of the Central Paratethys among which 3.1 to 3.7  $\mu\text{m}$  long placoliths have not been observed (Fig. 4A).

Two different size limits have been used to separate *R. haqii* from *R. pseudoumbilicus*: Backman (1980) proposed an upper size limit of 5  $\mu\text{m}$  for *R. haqii*, whereas Rio et al. (1990) and Raffi et al. (1995) proposed 7  $\mu\text{m}$ . In any case, *R. haqii* and *R. pseudoumbilicus* represent a continuous plexus in world ocean and therefore Young (1990) classified them as one species, *R. pseudoumbilicus*, with two varieties: *R. p. pseudoumbilicus* and *R. p. haqii*. Nevertheless, separation of two species at 5  $\mu\text{m}$  has prevailed.

As in the oceanic realm, the length of the placoliths in the *R. haqii*–*pseudoumbilicus* group changes continuously in the Central Paratethys. Limit of 5  $\mu\text{m}$  has also been used for separation of *R. pseudoumbilicus* and *R. haqii* (Ćorić and Švábenická 2004; Tomanová Petrová and Švábenická 2007; Ćorić and Hohenegger 2008). In addition, Ćorić and Hohenegger (2008) have distinguished two size groups among *R. pseudoumbilicus*, one between 5–7  $\mu\text{m}$ , the other > 7  $\mu\text{m}$ . In studied material, boundary 8  $\mu\text{m}$  seems to be pronounced. It appears that separation of group > 8  $\mu\text{m}$  is specific only for studied set of samples because qualitative examination of placoliths from the other Central Paratethys sections of Zone NN5 and NN6 showed gradual changes in size of *R. pseudoumbilicus* from 5 to 10  $\mu\text{m}$  (Jamrich and Halásová 2010).




Fig. 5. Variability of the placolith size for the individual assemblages, Egerian to Eggenburgian. Vertical axis, number of specimens; horizontal axis, placolith size (in  $\mu\text{m}$ ). Location of the boreholes LR10, LR9, LR2, C2, FV1, EH1, EH2, and section Lipovany (LIP) is illustrated on Fig. 1, their lithology and sampled intervals on Fig. 2. Abbreviation: LO, last occurrence.

<http://dx.doi.org/10.4202/app.2009.0006>






Fig. 6. Variability of the placolith size for the individual assemblages, Ottnangian to Early Badenian. Vertical axis, number of specimens; horizontal axis, placolith size (in  $\mu\text{m}$ ). Location of the boreholes D19, LKŠ1, BE2, PY1, and sections Lipovany (LIP) and Kralice (KRAS) is illustrated on Fig. 1, their lithology and sampled intervals on Fig. 2.




Fig. 7. Variability of the central opening size for the individual assemblages, Egerian to Eggenburgian. Vertical axis, number of specimens; horizontal axis, central opening size (in µm). Location of the boreholes LR10, LR9, LR2, C2, FV1, EH1, EH2, and section Lipovany (LIP) is illustrated on Fig. 1, their lithology and sampled intervals on Fig. 2. Abbreviation: LO, last occurrence.




Fig. 8. Variability of the central opening size for the individual assemblages, Ottomanian to Early Badenian. Vertical axis, number of specimens; horizontal axis, central opening size (in µm). Location of the boreholes D19, LKS1, BE2, PY1, and sections Lipovany (LIP) and Kralice (KRAS) is illustrated on Fig. 1, their lithology and sampled intervals on Fig. 2.




Fig. 9. Morphometric changes of placoliths of *Reticulofenestra minuta* and *Reticulofenestra haqii-pseudoumbilicus* group from the Egerian to the Early Badenian (Burdigalian–Langhian) in the Central Paratethys. Cooling Mi events according Zachos et al. 2001. MMCO = the Middle Miocene Climatic Optimum (Zachos et al. 2001). Abbreviation: LO, last occurrence.



In conclusion, a satisfactory size has not been found upon which to distinguish species in the *R. haqii*–*pseudumbilicus* group and its taxonomy is simply a matter of convention.

### Size changes in the Late Oligocene–Early Miocene *Reticulofenestra* and paleoecologic and paleogeographic events in the Central Paratethys

**Blooms of *Reticulofenestra minuta*.**—Two blooms of the small *R. minuta* were observed in the studied material (Figs. 9, 11): (i) the oldest one in the earliest Miocene (Biochron NN1 and early part of Biochron NN2), is described for the first time. It is probably a local event in the northern part of the Pannonian Basin; (ii) the early Middle Miocene (NN5) event was already documented; it occurred throughout the Central Paratethys (Ćorić and Švábenická 2004; Ćorić and Hohenegger 2008, Spezzaferri et al. 2009). In addition to these two widespread blooms, sporadic blooms of *R. minuta* have also been described in the Central Paratethys, having occurred during the Eggenburgian (Holcová 2002, 2005) and the Karpatian (Spezzaferri and Ćorić 2001; Švábenická et al. 2003).

Besides the Central Paratethys, blooms of *R. minuta* have been described from several stratigraphical levels in the Miocene from the middle to high latitudes (e.g., Cita et al. 1978; Backman 1980). These blooms are heterochronous with the Central Paratethys ones and reflect local paleoenvironmental events (e.g., Messinian salinity crisis; Cita et al. 1978; Wade and Bown 2006).

Interpretation of the causes of the blooms is ambiguous due to inconsistent data about the ecological requirements of small *Reticulofenestra*. Generally, the species is opportunistic and characterizes assemblages from near-shore environment (Haq 1980). Their blooms are connected with environmental stress characterized by rapid environmental changes (Wade and Bown 2006).

Gartner et al. (1983) suggested that variations in the relative abundance of *R. minuta* correlate with changes in nutrient dynamics. Wells and Okada (1997), Flores et al. (1997), Bollmann et al. (1998) and Kameo (2002) regard small *Reticulofenestra* spp. as eutrophic species while Hallock (1987), Bafort and Aubry (1992), Ćorić and Rögl (2004) suggested that blooms of small *Reticulofenestra* indicate oligotrophic warm water. Wade and Bown (2006) showed that *R. minuta* may have tolerated the brackish to hypersaline, high productivity environments, which prevailed immediately before and after the evaporite deposition during the Messinian salinity crisis.

The two conspicuous blooms of small *R. minuta* in the Central Paratethys occurred during transgression events when the Slovenian corridor permitted incursions of warm water into the Paratethys (Rögl 1998, 1999; Popov et al. 2004). These events are marked by horizons with larger Foraminifera—*Miogypsina* and *Lepidocyclina* in the lowermost Miocene (Váňová 1975), *Amphistegina* and *Planostegina* in the Lower Badenian—and replacement of small *Globigerina* fauna by diversified assemblages with large-sized warm-




Fig. 10. Two morphotypes of *Reticulofenestra haqii*–*pseudumbilicus* group in the lower part of the Zone NN2. Interval FO *Helicosphaera scissural* FO *Discoaster druggii*–FO *Helicosphaera ampliapertura* lower Zone NN2 (22.8–20.0 Ma); n = 243 (samples LR2/28, C2/10, FV1/240, FV1/40, EH2/155, EH1/85, EH1/105).

water species—*Globigerinoides* in the Lower Miocene and *Globigerinoides*, *Praeorbulina*, and *Orbulina* in the Middle Miocene). At the beginning of the blooms, *R. minuta* replaces *Coccolithus pelagicus*, which is an indicator of cold and nutrient-rich waters (McIntyre and Be 1967; Rahman and Roth 1990). Among the benthic foraminifera, the abundance of agglutinated foraminifera decreases (Vass et al. 1983; Holcová 2001; Spezzaferri et al. 2009). The paleoenvironmental turnover correlatable with the Middle Miocene bloom of *R. minuta* has been well explained: Upwelling regime with cooler surface waters and partly dysoxic bottom conditions of the latest Early Miocene were gradually changed to the warm-water condition in the early Middle Miocene and antiestuarine circulation with evaporation prevailing over fresh-water input (Brzobohatý 1987; Ćorić and Rögl 2004; Báldi 2006; Spezzaferri et al. 2009). Comparable circulation turnover due to mixing of in situ cold and incoming warm water masses and paleogeographical changes may be expected also in the earliest Miocene in the South Slovak Basin (Halášová et al. 1996).

Although the incoming of warm-water elements in the marine realm is specific of the two transgressive events, the Early and Middle Miocene conditions in the Central Paratethys substantially differed from the global paleoclimatic situation. The Early Miocene event was marked with the appearance of arctotertiary elements among palynomorphs (Vass et al. 1988; Planderová 1990) indicative of a global cooling correlated with the Mi1 and Mi1a events. In contrast, the Middle Miocene event can be correlated with the Middle Miocene Climatic Optimum (Gonera et al. 2000; Bicchi et al. 2003; Böhme 2003). Short-term oscillations in the abundance of *R. minuta* during the two blooms were observed (Ćorić and Hohenegger 2008; own unpublished data). These may indicate unstable




Fig. 11. Correlations between reticulofenestrid biometric events, local and global sea-level changes and paleogeographical events in the Egerian–Early Badenian (Burdigalian–Langhian) of the Central Paratethys. Abbreviations: FCO, first common occurrence; FO, first occurrence; LCO, last common occurrence; LO, last occurrence; MMCO, the Middle Miocene Climatic Optimum; *D.*, *Discoaster*; *H.*, *Helicosphaera*; *P.*, *Praeorbulina*; *R.*, *Reticulofenestra*; *S.*, *Sphenolithus*.

conditions with short-term oscillations of paleoenvironmental parameters mainly salinity and nutrients. These oscillations have been interpreted in the Biochron NN5 by Hohenegger et al. (2008): warm intervals characterized by high terrigenous input and higher seasonality were alternated with cold periods with lower terrigenous input and lower seasonality. *Reticule-*

*fenestram minuta* strongly dominated in the near-shore facies during warm intervals (more than 80%) where high terrigenous input during wet seasons may cause seasonal oscillations of salinity in the upper layer of water column (own unpublished data).

The expansion of near-shore facies as a result of transgres-




Fig. 12. Elliptical reticulofenestrids from the Central Paratethys Oligocene and Early Miocene. **A–K.** *Reticulofenestra minuta* Roth, 1970, < 3.5  $\mu\text{m}$ , zones NN1 and NN2. **A.** Borehole LR-10/12 m, Zone NN2. **B–D.** Borehole LR-2/48 m, Zone NN1. **E, F.** Borehole LR-2/28 m, Zone NN2. **G.** Borehole FV-1/440 m, Zone NN1. **H.** Borehole EH-1/155 m, Zone NN2. **I, J.** Borehole EH-1/85 m, Zone NN2. **K.** Borehole EH-2/155 m, Zone NN2. **L–W.** *Reticulofenestra minuta* Roth, 1970, < 3.5  $\mu\text{m}$ , Zone NN5. **L–N.** Section Kralice, sample 4. **O, P** Section Kralice, sample 7. **Q, R.** Section Kralice, sample 10. **S–U.** Borehole PY-1/2.2 m. **V, W.** Borehole PY-1/1.5 m. **X–AY.** *Reticulofenestra haqii–pseudumbilicus* group (4–5  $\mu\text{m}$ ), zones NP25–NN4. **X, Y.** Borehole LR-10/350 m, Zone NP25.  $\rightarrow$



sion, the changes in water circulation due to the mixing of incoming warmer and in situ colder water masses and, probably, unstable conditions as a result of short term (?seasonal) oscillations of ecological parameters (such as salinity in the upper layers of water column) may have produced conditions favourable to the blooms of small-sized *Reticulofenestra*.

**Size changes of the *Reticulofenestra haqii*–*pseudoumbilicus* group.**—Discrepancies between the FOs of individual size categories in the *R. haqii*–*pseudoumbilicus* group (see chapter: Size changes over time) in the Central Paratethys and in the oceanic realm is notable for interpretation of local vs. global character of size changes in the group. The following discrepancies have been recorded:

(i) The FO of *R. haqii* ( $> 4 \mu\text{m}$ ) was observed in the studied area in Zone NP25 (Holcová 2005) whereas in the world ocean it lies at the NN1/NN2 boundary (Young 1999).

(ii) The FO of *R. pseudoumbilicus* ( $> 5 \mu\text{m}$ ) is given in Zone NN4 in the world ocean (Young 1999). In the Central Paratethys, Marunteanu (1999) and Chira (2004) reported this species from near the NN1/NN2 boundary, Andreyeva-Grigorievich et al. (2008) from Zone NN2 and Molčíková and Straník (1987) from Zone NN1. Holcová (2005, this paper) describe the FO of specimens  $> 5 \mu\text{m}$  together with FO of *R. haqii*  $> 4 \mu\text{m}$  in the uppermost part of the NP25 Zone (Figs. 5, 6, 9).

(iii) The FO of *R. pseudoumbilicus*  $> 7 \mu\text{m}$  is given most often being a Middle Miocene event. In the Mediterranean, the FO has been described from Zone NN6 (Fornaciari and Rio 1996), and the FCO of *R. pseudoumbilicus*  $> 7.0 \mu\text{m}$  to define the zones MNN6a and MNN6b (Fornaciari et al. 1996); in the Eastern equatorial Pacific the event was recorded near the NN5/NN6 boundary (Raffi and Flores 1995). However, Howe and Sblendorio-Levy (1998) described the FO of *R. pseudoumbilicus*  $> 7 \mu\text{m}$  from the upper part of Zone NN 2 in the Atlantic Ocean near Madeira. This correlates with its FO in the Central Paratethys in Zone NN2 (Molčíková and Straník 1987; this paper). The FCO of *R. pseudoumbilicus*  $> 7 \mu\text{m}$  in the Central Paratethys has been recorded from the NN4 Zone. (Molčíková 1983; Švábenická et al. 2003; Eva Halášová, personal communication 2009; this paper).

The Central Paratethys FOs of all three size categories in *R. haqii*–*pseudoumbilicus* group occurred earlier in the global ocean (including the Mediterranean realm) and cannot be used for interregional biostratigraphic correlations.

The evolution of size in the *R. haqii*–*pseudoumbilicus* group in the Central Paratethys would suggest an earlier ap-

pearance of larger specimens owing to specific conditions in an intracontinental basin. However, the most important morphological changes in the *R. haqii*–*pseudoumbilicus* group can be correlated with the opening of new pathways between the Central Paratethys and the Mediterranean and the immigration of incoming species in the Central Paratethys (Fig. 11). First, the FO of *R. haqii* is associated with the reopening of sea-ways after the Kiscellian anoxia, which is marked by the immigration of foraminifera (Holcová 2008). Second at the upper part of Zone NN2 marked by the FO of *H. ampliapertura*, larger specimens of *R. haqii* ( $6\text{--}7 \mu\text{m}$ ) with small central opening ( $1.5\text{--}1.7 \mu\text{m}$ ) were recorded. The FO of *H. ampliapertura* can be correlated with transgression from the Mediterranean via the Pre-Alpine passage (Rögl 1998; Popov et al. 2004). The transgression is characterized by substantial faunal turnover marked by numerous first occurrences of species of western Mediterranean origin (Kroh and Harzhauser 1999; Vávra 1979; Mandić and Steininger 2003; Piller et al. 2007). Third, the FCO of *R. pseudoumbilicus*  $> 7 \mu\text{m}$  at the base of Zone NN4 correlates with the transgression and paleogeographic reorganization and the establishment of a new broad connection via the “Trans-Tethyan trench corridor”. The connection enabled faunal exchange between the Mediterranean and Paratethys (Rögl 1998; Rögl et al. 2003; Popov et al. 2004). Fourth, the FO of *R. pseudoumbilicus*  $> 8 \mu\text{m}$  is associated with the reopening of the Mediterranean–Indo-Pacific seaway (Rögl and Steininger 1983; Rögl 1998, 1999; Popov et al. 2004). This large transgression caused rapid immigration of the marine fauna (e.g., Harzhauser and Piller 2007; Holcová 2008) termed the EBBE, or Early Badenian-Build-up-Event (Harzhauser and Piller 2007).

Slight gradual size changes in the *R. haqii*–*pseudoumbilicus* group were recorded during periods without paleogeographical changes (Fig. 11). The decrease in the placoliths size in the late Egerian (upper NP25 to NN1 zonal interval) may correspond with cooling interpreted from occurrence of arcto-tertiary elements among palynomorphs (Planderová 1990). Also Kürschner et al. (2008) hypothesized a cooling during this interval correlatable with the Mi1/1a event.

The size of placoliths slightly increased in the Eggenburgian, although this increase is not general (Figs. 5, 6, 9). This size increase can be correlated with warming (Kürschner et al. 2008).

Bimodal size distribution in the *R. haqii*–*pseudoumbilicus* group was recorded firstly during the Zone NN1 and in the lower part of the Zone NN2 (Fig. 9, 10). The smaller

Z, AA. Borehole LR-9/350 m, Zone NP25. AB. Borehole LR-2/98 m, Zone NP25. AC–AE. Borehole C-2/500 m, Zone NP25. AF, AG. Borehole EH2/160 m, Zone NN1. AH, AI. Borehole EH-1/105 m, Zone NN2. AJ, AK. Borehole EH-1/85 m, Zone NN2. AL. Borehole EH-2/75 m, Zone NN2. AM, AN. Borehole LKŠ-1/340 m, Zone NN3. AO–Q. Borehole LKŠ-1/181 m, Zone NN4. AR, AS. Borehole LKŠ-1/105 m, Zone NN4. AT–BE *Reticulofenestra haqii*–*pseudoumbilicus* group ( $5\text{--}7 \mu\text{m}$ ), zones NP25–NN4. AT, AU. Borehole LR-10/350 m, Zone NP25. AV. Borehole LR-9/350 m, Zone NP25. AW, AX. Borehole C-2/500 m, Zone NP25. AY. Borehole EH2/160 m, Zone NN1. AZ, BA. Borehole EH-1/105 m, Zone NN2. BB, BC. Borehole EH-1/85 m, Zone NN2. BD, BE. Borehole EH-2/155 m, Zone NN2. BF–BK. *Reticulofenestra haqii*–*pseudoumbilicus* group ( $6\text{--}7 \mu\text{m}$ ) narrow central opening, Zone NN2. BF, BG. Borehole EH-1/155 m, Zone NN2. BH, BI. Borehole EH-1/105 m, Zone NN2. BJ, BK. Borehole EH-2/85m, Zone NN2. BL–BQ. *Reticulofenestra haqii*–*pseudoumbilicus* group ( $6\text{--}8 \mu\text{m}$ ), Zone NN4. BL, BM. Borehole LKŠ-1/105 m, Zone NN4. BN, BO. Borehole LKŠ-1/26 m, Zone NN4. BP, BQ Borehole BE-2/114 m, Zone NN4. BR, BU. *Reticulofenestra haqii*–*pseudoumbilicus* group ( $> 8 \mu\text{m}$ ), Zone NN5. BR. Borehole PY-1/2.2 m. BS. Borehole PY-1/1.5 m. BT, BU. Section Kralice, sample 4.



morphotype (< 6 µm) is characterized by a small central opening 0.5–1.5 µm. Larger morphotype (> 6 µm) have larger central openings (Fig. 10).

Backman (1980) and Beaufort and Aubry (1992) described a dependence of the morphometric variability of *R. pseudoumbilicus* on seasonal or latitudinal temperature differences during the Miocene and Pliocene. According to Backman (1980) low latitude individuals are characterized by a larger central opening and can be slightly larger. High latitude individuals are distinguished by a smaller central opening. Beaufort and Aubry (1992) have proposed alternations of winter and summer morphotypes in the Lower Miocene of high-latitude populations. The occurrence of two size categories (3.5–6 µm and 6–8 µm) in the late Egerian–early Eggenburgian of analysed samples may also represent seasonal morphotypes: the smaller morphotype could represent the winter morphotype while the larger morphotype with a larger central opening could represent the summer morphotype (Fig. 10). This assumed seasonality has been corroborated by palaeoclimate estimates based on plant macrofossils preserved in the Břešťany Clay from the Most Basin (Teodoridis and Kvaček 2006; Kvaček and Teodoridis 2007). These studies have shown a significant difference between a Coldest Month Mean Temperature (CMMT) and Warmest Month Mean Temperature (WMMT), which is 15–20°C (Teodoridis 2007; Teodoridis et al. 2011).

## Conclusions

Based on biometric analysis, two size groups of elliptical reticulofenestrids from the NP25–NN5 zonal interval were distinguished. One group corresponds to the small *R. minuta* (< 3 µm); the other is the *R. haqii*–*pseudoumbilicus* group (3.5–10 µm). A clear criterion upon which to define species in this group has not been found, and species taxonomy (*R. haqii* vs. *R. pseudoumbilicus*) can only be conventional (Fig. 12).

Two blooms of small *R. minuta* were observed. The earliest Miocene bloom (NN1 and lower part of Zone NN2) is probably a local event for the northern part of the Pannonian Basin. The early Middle Miocene bloom (Zone NN5) was already known and has a global occurrence in the Central Paratethys basins. These blooms can be correlated with the expansion of near-shore facies associated with transgression, the incursions of warm water into the Paratethys, which may cause changes in water circulation due to the mixing of warmer and colder water masses. Unstable conditions with short-term oscillations of ecological parameters (probable salinity in the upper layers of water column) can be also expected.

The FOs of all three size categories in *R. haqii*–*pseudoumbilicus* group (3–5 µm, 5–7 µm, > 7 µm) occurred earlier in the Central Paratethys than in the global ocean and the Mediterranean region. They cannot be used for interregional biostratigraphical correlations.

The slight gradual changes in size in the *R. haqii*–*pseudoumbilicus* group can be correlated with paleoclimatic trends.

The decrease in size of the placoliths in the late Egerian (upper Zone NP25 and Zone NN1) may result from cooling during the Mi1/1a event.

The size of placoliths slightly increased from the Eggenburgian to the Badenian; the interval can be generally characterized by warming.

The two morphotypes occurring in the late Egerian–Early Eggenburgian may represent seasonal morphotypes. The smaller morphotype could represent the winter morphotype while the larger morphotype with a slightly larger central opening could represent the summer morphotype.

## Acknowledgements

I would like to acknowledge the helpful discussion on Oligocene and Miocene paleoclimate by Zlatko Kvaček and Vasilis Teodoridis (both Charles University in Prague, Czech Republic). I thank the Marie-Pierre Aubry (Rutgers University, New Jersey, USA), and Jörg Mutterlose (Ruhr-Universität Bochum, Germany) for their comments and corrections, which substantially improved the quality of this manuscript. This research was supported by grants no. MSM0021620855 and GAČR 205/09/0103.

## References

- Andreyeva-Grigorovich, A.S., Kulchytsky, Y.O., Gruzman, A.D., Lozonyak, P.Y., Petrashkevich, M.I., Portnyagina, L.O., Ivanina, A.V., Smirnov, S.E., Trofimovich, N.A., and Savitskaya, N.A. 1997. Regional stratigraphic scheme of Neogene formations of the Central Paratethys of Ukraine. *Geologica Carpathica* 48: 123–136.
- Andreyeva-Grigorovich, A.S., Oszczyppo, N., Slaczka, A., Oszczyppo-Clowes, M., Savitskaya, N.A., and Trofimovich, N.A. 2008. New data on the stratigraphy of the folded Miocene Zone at the front of the Ukrainian Outer Carpathians. *Acta Geologica Polonica* 58: 325–354.
- Backman, J. 1978. Late Miocene–Early Pliocene nannofossil biochronology and biogeography in the Vera Basin, SE Spain. *Stockholm Contributions in Geology* 32: 93–114.
- Backman, J. 1980. Miocene–Pliocene nannofossils and sedimentation rates in the Hatton-Rockall Basin, NE Atlantic Ocean. *Stockholm Contributions in Geology* 36: 1–91.
- Báldi, K. 2006. Paleooceanography and climate of the Badenian (Middle Miocene, 16.4–13.0 Ma) in the Central Paratethys based on foraminifera and stable isotope ( $\delta^{18}\text{O}$  and  $\delta^{13}\text{C}$ ) evidence. *International Journal of Earth Science (Geologische Rundschau)* 95: 119–142.
- Báldi T. 1986. *Mid-tertiary Stratigraphy and Paleogeographic Evolution of Hungary*. 201 pp. Akademiai Kiado, Budapest.
- Beaufort, L. and Aubry, M.-P. 1992. Palaeoceanographic implications of a 17 m.y. long record of high-latitude Miocene calcareous nannoplankton fluctuations. *Proceeding of Ocean Drilling Program, Scientific Results* 120: 530–549.
- Berggren, W.A., Kent, D.V., Swisher III., C.C., and Aubry, M.-P. 1995. A revised cenozoic geochronology and chronostratigraphy. In: W.A. Berggren, D.V. Kent, and J. Hardenbol (eds.), *Geochronology, Time Scale and Global Stratigraphic Correlations: A Unified Temporal Framework for an Historical Geology*. *Society of Economic Paleontologists and Mineralogists, Special Publication* 54: 129–212.
- Bicchi, E., Ferrero, E., and Gonera, M. 2003. Palaeoclimatic interpretation based on Middle Miocene planktonic Foraminifera: the Silesia Basin (Paratethys) and Monferrato (Tethys) records. *Palaeogeography, Palaeoclimatology, Palaeoecology* 196: 265–303.
- Böhme, M. 2003. The miocene climatic optimum: evidence from ectothermic

- vertebrates of Central Europe. *Palaeogeography, Palaeoclimatology, Palaeoecology* 195: 389–401.
- Bollmann, J., Baumann, K.H., and Thierstein, H.R. 1998. Global dominance of *Gephyrocapsa* coccoliths in the late Pleistocene: Selective dissolution, evolution, or global environmental change? *Paleoceanography* 13: 517–529.
- Brzobohatý, R. 1987. Contribution to paleogeography of the Miocene basins of the Central Paratethys from otolith fauna. *Miscellanea Micropalaeontologica II/2, Knižovnička zemního plynu a nafty* 6b: 101–111.
- Chira, C. 2004. Early Miocene calcareous nannofossil assemblages from Transylvania. *Acta Paleontologica Romaniaae* 4: 81–88.
- Chlupáč, I., Brzobohatý, R., Kovanda, J., and Stráník, Z. 2002. *Geologická minulost České republiky*. 436 pp. Academia, Praha.
- Cita, M.B., Wright, R.C., Ryan, W.B.F., and Longinelli, A. 1978. Messinian paleoenvironments, *Initial Reports of the Deep Sea Drilling Project* 42A: 1003–1035.
- Ćorić, S. and Hohenegger, J. 2008. Quantitative analyses of calcareous nannoplankton assemblages from the Baden-Sooss section (Middle Miocene of Vienna Basin, Austria). *Geologica Carpathica* 59: 447–460.
- Ćorić, S. and Rögl, F. 2004. Roggendorf-1 borehole, a key-section for Lower Badenian transgressions and the stratigraphic position of the Grund Formation (Molasse Basin, Lower Austria). *Geologica Carpathica* 55: 165–178.
- Ćorić, S. and Švábenická, L. 2004. Calcareous nannofossil biostratigraphy of the Grund Formation (Molasse Basin, Lower Austria). *Geologica Carpathica* 55: 147–155.
- Flores, J.A., Sierro, F.S., Francés, G., Vasquez, A., and Zamarreno, I. 1997. The last 100,000 years in the western Mediterranean: Sea surface water and frontal dynamics as revealed by coccolithophores. *Marine Micropaleontology* 29: 351–366.
- Fornaciari, E. and Rio, D. 1996. Latest Oligocene to early middle Miocene quantitative calcareous nannofossil biostratigraphy in the Mediterranean region. *Micropaleontology* 42: 1–36.
- Fornaciari, E., Di Stefano, A., Rio, D., and Negri, A. 1996. Middle Miocene quantitative calcareous nannofossil biostratigraphy in the Mediterranean region. *Micropaleontology* 42: 37–63.
- Gartner, S., Chen, M.P., and Stanton, R.J. 1983. Late Neogene nanofossil biostratigraphy and paleoceanography of the northeastern Gulf of Mexico and adjacent areas. *Marine Micropaleontology* 8: 17–50.
- Gonera, M., Peryt, T.M., and Durakiewicz, T. 2000. Biostratigraphical and paleoenvironmental implications of isotopic studies ( $^{18}\text{O}$ ,  $^{13}\text{C}$ ) of Middle Miocene (Badenian) foraminifers in the Central Paratethys. *Terra Nova* 12: 231–238.
- Halássová, E., Hudáčková, N., Holcová, K., Vass, D., Elečko, M., and Pěrszlényi, M. 1996. Sea ways connecting the Filakovo-Pétervasara Basin with the Eggenburgian/Burdigalian open sea. *Slovak Geological Magazine* 2: 125–136.
- Hallock, P. 1987. Fluctuations in the trophic resource continuum: a factor in global diversity cycles? *Paleoceanography* 2: 457–471.
- Haq, B.U. 1980. Biogeographic history of the Miocene calcareous nannoplankton and paleoceanography of the Atlantic Ocean. *Micropaleontology* 26: 414–443.
- Haq, B.U., Hardenbol, J., and Vail, P.R. 1988. Mesozoic and Cenozoic chronostratigraphy and cycles of sea-level change. *Society of Economic Paleontologists and Mineralogists, Special Publication* 42: 71–108.
- Harzhauser, M. and Piller, W.E. 2007. Benchmark data of a changing sea—paleogeography, palaeobiogeography and events in the Central Paratethys during the Miocene. *Palaeogeography, Palaeoclimatology, Palaeoecology* 253: 8–31.
- Hohenegger, J., Andersen, N., Báldi, K., Ćorić, S., Pervesler, P., Rupp, C., and Wagreich, M. 2008. Paleoenvironment of the Early Badenian (Middle Miocene) in the southern Vienna Basin (Austria)—multivariate analysis of the Baden-Sooss section. *Geologica Carpathica* 59: 461–487.
- Hohenegger, J., Coric, S., Khatun, M., Pervesler, P., Rögl, F., Rupp, Ch., Selge, A., Uchman, A., and Wagreich, M. 2009. Cyclostratigraphic dating in the Lower Badenian (Middle Miocene) of the Vienna Basin (Austria)—the Baden-Sooss core. *International Journal of Earth Sciences* 98: 915–930.
- Holcová, K. 2001. New methods in foraminiferal and calcareous nannoplankton analysis and evolution of Oligocene and Miocene basins of the Southern Slovakia. *Slovak Geological Magazine* 7: 19–41.
- Holcová, K. 2002. Calcareous nannoplankton from the Eggenburgian stratotypes (Lower Miocene, Central Paratethys). *Geologica Carpathica* 53: 381–390.
- Holcová, K. 2005. Quantitative calcareous nannoplankton biostratigraphy of the Oligocene/Miocene boundary interval in the northern part of the Buda Basin (Central Paratethys). *Geological Quarterly* 49: 263–274.
- Holcová, K. 2008. Foraminiferal species diversity in the Central Paratethys—a reflection of global or local events? *Geologica Carpathica* 59: 71–85.
- Howe, R.W. and Sblendorio-Levy, J. 1998. Calcareous nannofossil biostratigraphy and sediment accumulation of turbidite sequences on the Madeira abyssal plain, sites 950–952. In: P.P.E. Weaver, H.U. Schmincke, and J.V. Firth (eds.), *Proceeding of the Ocean Drilling Program, Scientific Results* 157: 501–520.
- Jamrich, M. and Halássová, E. 2010. The evolution of the Late Badenian calcareous nannofossil assemblages as a reflexion of the paleoenvironmental changes of the Vienna Basin (Devínska Nová Ves-clay pit). *Acta Geologica Slovaca* 2: 123–140.
- Kameo, K. 2002. Late Pliocene Caribbean surface water dynamics and climatic changes based on calcareous nannofossil records. *Palaeogeography, Palaeoclimatology, Palaeoecology* 179: 211–226.
- Kováč, M. 2000. *Geodynamic, Paleogeographical and Structural Development of the Carpathian–Pannonian Region During the Miocene*. 202 pp. Veda, Bratislava.
- Kroh, A. and Harzhauser, M. 1999. An echinoderm fauna from the Lower Miocene of Austria: Paleocology and implications for Central Paratethys paleogeography. *Annalen des Naturhistorischen Museums in Wien* 101A: 145–191.
- Kürschner, W.M., Kvaček, Z., and Dilcher, D.L. 2008. The impact of Miocene atmospheric carbon dioxide fluctuations on climate and the evolution of terrestrial ecosystems. *Proceedings of the National Academy of Sciences of the United States of America* 105: 449–453.
- Kvaček, Z. and Teodoridis, V. 2007. Tertiary macrofloras of the Bohemian Massif: a review with correlations within Boreal and Central Europe. *Bulletin of Geosciences* 82: 383–408.
- Lehotayová, R. 1975. *Reticulofenestra excavata* n. sp. from the Eggenburgian to Karpatian of Central Slovakia. *Západné Karpaty, séria paleontológia* 2: 39–40.
- Lehotayová, R. 1982. Miocene nannoplankton zones in west Carpathians. *Západné Karpaty, séria paleontológia* 8: 91–110.
- Lehotayová, R. 1984. Lower Miocene calcareous nannoflora of the West Carpathians. *Západné Karpaty, séria paleontológia* 9: 99–110.
- Lourens, L., Hilgen, F., Shackleton, N.J., Laskar, J., and Wilson, D. 2004. The Neogene period. In: F. Gradstein, J. Ogg, and A. Smith (eds.), *A Geological Time Scale*, 409–440. Cambridge University Press, Cambridge.
- Mandic, O. and Steininger, F.F. 2003. Computer-based mollusc stratigraphy—a case study from the Eggenburgian (Lower Miocene) type region (NE Austria). *Palaeogeography, Palaeoclimatology, Palaeoecology* 197: 263–291.
- Martini, E. 1971. Standard Tertiary and Quaternary calcareous nannoplankton zonation. In: A. Farinacci (ed.), *Proceeding of 2nd Planktonic Conference, Roma 1970, Roma*, 739–785. Tecnoscienza, Rome.
- Márton, E., Vass, D., and Túnyi, I. 1995. The Late Tertiary rotation of the megaunit Pelső and the adjacent West Carpathians area. *Knižovnička Zemní Plyn a Nafta* 16: 97–108.
- Marunteanu, M. 1992. Distribution of the Miocene calcareous nannofossils in the Intra- and Extra- Carpathian areas of Rumania. *Knižovnička Zemní Plyn Nafta* 14b (2): 247–262.
- Marunteanu, M. 1999. Litho- and biostratigraphy (calcareous nannoplankton) of the Miocene deposits from the Outer Moldavides. *Geologica Carpathica* 50: 313–324.
- McIntyre, A. and Be, A.W.H. 1967. *Coccolithus neohelis* sp.n. a coccolith fossil type in contemporary seas. *Deep-Sea Research* 14: 369–371.
- Molčíková, V. 1983. The Karpatian and Lower Badenian Calcareous nannoflora of the Carpathian Foredeep in Moravia. *Knižovnička Zemní Plyn a Nafta* 4: 283–304.
- Molčíková, V. and Stráník, Z. 1987. Calcareous nannoplankton from the



- Ždánice-Hustopeče Formation and their relation to underlying formations. *Knihovnička Zemní Plyn a Nafta* 6b: 59–76.
- Piller, W.E., Harzhauser, M., and Mandic, O. 2007. Miocene Central Paratethys stratigraphy—current status and future directions. *Stratigraphy* 4: 151–168.
- Planderová, E. 1990. *Miocene Microflora of Slovak Central Paratethys and its Biostratigraphical Significance*. 144 pp. Geologický Ústav Dionýza Štúra, Bratislava.
- Popov, S.V., Rögl, F., Rozanov, A.Y., Steininger, F.F., Shcherba, I.G., and Kováč, M. 2004. Lithological-Paleogeographic maps of Paratethys. *Courier Forschungsinstitut Senckenberg* 250: 1–46.
- Pujos, A. 1985. Cenozoic nannofossils, central equatorial Pacific, deep sea Drilling Project Leg 85. In: L. Mayer and F. Theyer (eds.), *Initial Reports of the Deep Sea Drilling Project* 85: 581–608.
- Raffi, I. and Flores, J.A. 1995. Pleistocene through Miocene calcareous nannofossils from Eastern Equatorial Pacific Ocean (Leg 138). In: N.G. Piasias, L.A. Mayer, T.R. Janecek, A. Palmer-Julson, and T.H. van Andel (eds.), *Proceedings of the Ocean Drilling Program, Scientific Results* 138: 233–286.
- Raffi, I., Rio, D., d' Atri, A., Fornaciari, E., and Rochetti, S. 1995. Quantitative distribution patterns and biomagnetostratigraphy of middle and late Miocene calcareous nannofossils from western equatorial Indian and Pacific Oceans (Legs 115, 130, and 138). In: N.G. Piasias, L.A. Mayer, T.R. Janecek, A. Palmer-Julson, and T.H. van Andel (eds.), *Proceeding of Oceanic Drilling Project, Scientific Results* 138: 479–502.
- Rahman, A. and Roth, P.H. 1990. Late Neogene paleoceanography and paleoclimatology of the Gulf of Aden region based on calcareous nannofossils. *Paleoceanography* 5: 91–107.
- Rio, D., Fornaciari, E., and Raffi, I. 1990. Late Oligocene through early Pleistocene calcareous nannofossils from western equatorial Indian Ocean (Leg 115). *Proceedings of the Ocean Drilling Program, Scientific Results* 115: 175–221.
- Rögl, F. 1998. Paleogeographic considerations for Mediterranean and Paratethys seaways (Oligocene to Miocene). *Annalen des Naturhistorische Museum in Wien* 99A: 279–310.
- Rögl, F. 1999. Mediterranean and Paratethys. facts and hypotheses of an Oligocene to Miocene paleogeography (short overview). *Geologica Carpathica* 50: 339–349.
- Rögl, F. and Steininger, F.F. 1983. Vom Zerfall der Tethys zu Mediterran und Paratethys—Die neogene Palaeogeographie und Palinspastik des zirkummediterranen Raumes. *Annalen des Naturhistorische Museum in Wien* 85/A: 135–164.
- Rögl, F., Ćorić, S., Daxner-Höck, G., Harzhauser, M., Mandić, O., Švábenická, L., and Zorn, I. 2003. Correlation of the Karpatian Stage. In: R. Brzobohatý, I. Cicha, M. Kováč, and F. Rögl (eds.), *The Karpatian—a Lower Miocene Stage of the Central Paratethys*, 27–35. Masaryk University, Brno.
- Seneš, J. 1961. Paläogeographie des Westkarpatischen Raumes in Beziehung zur übrigen Paratethys in Miozän. *Geologické práce* 60: 160–194.
- Spezzaferri, S. and Ćorić, S. 2001. Ecology of Karpatian (Early Miocene) foraminifera and calcareous nannoplankton from Laa an der Thaya, Lower Austria: a statistical approach. *Geologica Carpathica* 52: 361–374.
- Spezzaferri, S., Ćorić, S., and Stingl, K. 2009. Palaeoenvironmental reconstruction of the Karpatian–Badenian (Late Burdigalian–Early Langhian) transition in the Central Paratethys. A case study from the Wagna Section (Austria). *Acta Geologica Polonica* 59: 523–544.
- Švábenická, L. 2002. Calcareous nannofossils of the Upper Karpatian and Lower Badenian Deposits in the Carpathian Foredeep, Moravia (Czech Republic). *Geologica Carpathica* 53: 197–210.
- Švábenická, L., Coric, S., Andreyeva-Grigorovich, A.S., Halášová, E., Marunteanu, M., Nagymarosi, A., and Oszcypko-Clowes, M. 2003. Central Paratethys Karpatian Calcareous Nannofossils. R. Brzobohatý, I. Cicha, M. Kováč, and F. Rögl (eds.), *The Karpatian—a Lower Miocene Stage of the Central Paratethys*, 151–168. Masaryk University, Brno.
- Teodoridis, V. 2007. Overview of macrofloras from the Most Basin (Czech Republic) and their phytostratigraphical correlation within Central and Boreal Europe. In: Organizing Committee in Bonn (eds.), *Terra Nostra* 2, *IPC-XII /IOPC-VIII Bonn, Germany 2008, Abstract Volume*, 280. GeoUnion Alfred-Wegener-Stiftung, Berlin.
- Teodoridis, V. and Kvaček, Z. 2006. Palaeobotanical research of the Early Miocene deposits overlying the main coal seam (Libkovice and Lom Mbs.) in the Most Basin (Czech Republic). *Bulletin of Geosciences* 81: 93–113.
- Teodoridis, V., Mazouch, P., Spicer, R.A., and Uhl, D. 2011. Refining CLAMP—Investigations towards improving the Climate Leaf Analysis Multivariate Program. *Palaeogeography, Palaeoclimatology, Palaeoecology* 299: 39–48.
- Tomanová Petrová, P. and Švábenická, L. 2007. Lower Badenian biostratigraphy and paleoecology: a case study from the Carpathian Foredeep (Czech Republic). *Geologica Carpathica* 58: 333–352.
- Váňová, M. 1975. *Lepidocyclina* and *Miogypsina* from the facistratotype localities Budikovany and Bretka (South Slovakia). In: T. Báldi and J. Seneš (eds.), *OM—Egerien, Chronostratigraphie und Neostatotypen*, 315–339. Slovak Academy of Sciences, Bratislava.
- Vass, D. 1995. Global sea level changes reflected on Northern margin of the Hungarian Paleogene the Filakovo and Novohrad (Nograd) Lower Miocene Basin (South Slovakia). *Mineralia slovacica* 27: 193–206.
- Vass, D. 1996. The origin and disappearance of Hungarian Paleogene Basins and short-term Lower Miocene Basin in Northern Hungary and Southern Slovakia. *Slovak Geological Magazine* 1: 81–95.
- Vass, D. and Elečko, M. 1982. Litostratigraphic units of the Kiscellian-Eggenburgian of the Rimava Basin and Cerová Upland. *Geologické Práce, Správy* 77: 111–124.
- Vass, D., Elečko, M., Pristaš, J., Lexa, J., Hanzel, V., Mordlitba, I., Jánová, V., Bodnár, J., Husák, L., Filo, M., Májovský, J., and Linkeš, V. 1989. *Geology of the Rimavska Kotlina Depression*. 160 pp. Geologický Ústav Dionýza Štúra, Bratislava.
- Vass, D., Hók, J., Kováč, P., and Elečko, M. 1993. Succession of the Paleogene and Neogene tectonic events in the light of tectonical analysis. *Mineralia Slovaca* 25: 79–82.
- Vass, D., Konečný, V., and Šefara, J. 1979. *Geology of the Ipeľ basin and Krupinská Planina Upland*. 227 pp. Geologický Ústav Dionýza Štúra, Bratislava.
- Vass, D., Elečko, M., Konečný, V., Krippel, M., Kubeš, P., Lexa, J., Pristaš, J., Zakovič, M., Vozár, J., Vozárová, A., Bodnár, J., Husák, L., Filo, M., Lacika, J., and Linkeš, V. 2007. *Geology of Lučenecká kotlina Depression and Cerová vrchovina Upland*. 284 pp. Geologický Ústav Dionýza Štúra, Bratislava.
- Vávra, N. 1979. Die Bryozoenfaunen des österreichischen Tertiärs. *Neues Jahrbuch für Geologie und Paläontologie, Abhandlungen* 157: 366–392.
- Wade, B.S. and Bown, P.R. 2006. Calcareous nannofossils in extreme environments: the Messinian Salinity Crisis, Polemi Basin, Cyprus. *Palaeogeography, Palaeoclimatology, Palaeoecology* 233: 271–286.
- Wells, P. and Okada, H. 1997. Response of nannoplankton to major changes in seasurface temperature and movements of hydrological fronts over site DSDP 594 (south Chatham Rise, southeastern New Zealand), during the last 130 kyr. *Marine Micropalaeontology* 32: 341–363.
- Young, Y.R. 1990. Size variations of neogene *Reticulofenestra* coccoliths from Indian DSDP cores. *Journal of Micropaleontology* 9: 71–86.
- Young, Y. 1999. Neogene. In: P.R. Bown (ed.), *Calcareous Nannofossil Biostratigraphy*, 225–265. Cambridge University Press, Cambridge.
- Zachos, J.M., Pagani, L., Sloan, E.T., and Billups, K. 2001. Trends, rhythms, and aberrations in global climate 65 Ma to present. *Science* 292: 686–693.
- Zágoršek, K., Holcová, K., and Trasoň, T. 2008. Bryozoan event from Middle Miocene (Early Badenian) lower neritic sediments from the locality Kralice nad Oslavou (Central Paratethys, Moravian part of the Carpathian Foredeep). *International Journal of Earth Sciences (Geologische Rundschau)* 97: 835–850.
- Zágoršek, K., Nehyba, S., Holcová, K., Hladíková, Š., and Kroh A. 2009. Fauna from the Middle Miocene (Early Badenian) sediments from the section Kralice nad Oslavou (Central Paratethys, Moravian part of the Carpathian Foredeep). *Bulletin of Geosciences* 84: 465–496.
- Zágoršek, K. and Holcová, K. 2009. The oldest Early Badenian Bryozoa event in the Carpathian Foredeep from the Přemyslovce (PY-1, PY-2) boreholes. *Přírodovědné Studie Muzea Prostějovska* 10–11: 53–59.