



Another new species of *Oedothorax* Bertkau, 1883 from India (Araneae, Linyphiidae)

Author: Tanasevitch, Andrei V.

Source: *Revue suisse de Zoologie*, 124(2) : 331-333

Published By: Muséum d'histoire naturelle, Genève

URL: <https://doi.org/10.5281/zenodo.893545>

BioOne Complete (complete.BioOne.org) is a full-text database of 200 subscribed and open-access titles in the biological, ecological, and environmental sciences published by nonprofit societies, associations, museums, institutions, and presses.

Your use of this PDF, the BioOne Complete website, and all posted and associated content indicates your acceptance of BioOne's Terms of Use, available at www.bioone.org/terms-of-use.

Usage of BioOne Complete content is strictly limited to personal, educational, and non-commercial use. Commercial inquiries or rights and permissions requests should be directed to the individual publisher as copyright holder.

BioOne sees sustainable scholarly publishing as an inherently collaborative enterprise connecting authors, nonprofit publishers, academic institutions, research libraries, and research funders in the common goal of maximizing access to critical research.

Another new species of *Oedothorax* Bertkau, 1883 from India (Araneae, Linyphiidae)

Andrei V. Tanasevitch

A.N. Severtsov Institute of Ecology and Evolution, Russian Academy of Sciences, Leninsky prospekt, 33, Moscow 119071, Russia. E-mail: tanasevitch@gmail.com

Abstract: A new species, *Oedothorax khasi* sp. nov., is described from the Indian state of Meghalaya on the basis of a single male. The species is similar to the Oriental *O. myanmar* Tanasevitch, 2017, different in some details of palpal structure.

Keywords: Erigoninae - Oriental Region - Meghalaya - Khasi Hills.

INTRODUCTION

At present, the spider fauna of India comprises of 14 species of *Oedothorax* Bertkau in Förster & Bertkau, 1883, which are distributed in equal proportions between the Oriental and the Palaearctic parts of the country (see Tanasevitch, 2015, 2016, 2017). An eighth *Oedothorax* from Oriental India is described below. The single male available of this new species was collected from montane subtropical forests in the Khasi Hills, a territory belonging to the Garo-Khasi range in the Indian state of Meghalaya.

MATERIAL AND METHODS

This paper is based on material kept at the Muséum d'histoire naturelle de Genève, Switzerland (MNHG). Specimens preserved in 70% ethanol were studied using a MBS-9 stereomicroscope. The sample number is given in square brackets.

Terminology of copulatory organs mainly follows that of Merrett (1963), Hormiga (2000) and Tanasevitch (2015). Chaetotaxy is given in a formula, e.g., 2.2.1.1, which refers to the number of dorsal spines on tibiae I-IV. The sequence of leg segment measurements is as follows: femur + patella + tibia + metatarsus + tarsus. All measurements are given in mm. Scale lines in the figures correspond to 0.1 mm unless indicated otherwise. Figure numbers are given above the scale lines, the alternative distance below.

Abbreviations

a.s.l.	Above sea level
C	Convector <i>sensu</i> Tanasevitch (1998) = lamella <i>sensu</i> Merrett (1963)
DAC	Distal apophysis of convector <i>sensu</i> Tanasevitch (2015)
DSA	Distal suprategular apophysis <i>sensu</i> Hormiga (2000)
E	Embolus
LEC	Lateral extension of convector <i>sensu</i> Tanasevitch (2015)
MBC	Main body of convector
MM	Median membrane <i>sensu</i> Helsdingen (1965) and Saaristo (1977)
R	Radix
TmI	Position of trichobothrium on metatarsus I


TAXONOMY

Oedothorax khasi sp. nov.

Figs 1-7

Holotype: Male; INDIA, Meghalaya, Khasi Hills, 16 km SW of Mawsynram, between Mawsynram and Balat, 1000 m a.s.l.; sifting in forest, in ravine; 27.X.1978; leg. C. Besuchet & I. Löbl [30b].

Diagnosis: The new species is characterized by an essentially unmodified carapace, by the shape of the dorsal apophysis of the palpal tibia, as well as by a relatively large, cylindrical radix and by a very short embolus.



Figs 1-7. *Oedothorax khasi* sp. nov., holotype. (1) Carapace, lateral view. (2) Distal part of right palp, retrolateral view. (3-4) Palpal tibia, dorsal and retrodorsal view, respectively. (5) Paracymbium. (6) Distal suprategular apophysis and embolic division, retrolateral view. (7) Embolic division, prolateral view.

Etymology: The specific name is a noun in apposition referring to the mountain range in which lies the type locality.

Description: *Male (holotype)*. Total length 1.80. Carapace 0.85 long, 0.60 wide, pale greyish brown. Cephalic part of carapace somewhat elevated and protruded forward as shown in Fig. 1. Eyes slightly enlarged. Chelicerae 0.33 long, unmodified. Legs pale brown. Leg I 3.46 long (0.93+0.25+0.85+0.78+0.65), IV 3.22 long (0.88+0.23+0.83+0.80+0.48). Chaetotaxy 2.2.1.1, length of spines about 1.5 times diameter of segment. All metatarsi with a trichobothrium. TmI 0.63. Palp (Figs 2-7): Tibia carrying a wide dorsal

apophysis with a small invagination apically and two small tubercles retrolaterally. Paracymbium somewhat L-shaped. Distal surface of tegulum membranous. Distal suprategular apophysis short, wide, rounded distally. Embolus very short, radix relatively large, cylindrical, with a fold on its exterior surface. Convector with a wide lateral extension and a short, pointed distal apophysis. Main body of convector long, tapering posteriorly. Abdomen 1.15 long, 0.63 wide, dorsally pale, with an indistinct grey herring-bone pattern. *Female*. Unknown.

Taxonomic remarks: *Oedothorax khasi* sp. nov. is similar to another Oriental species, *O. myanmar*,

recently described from mountains of the Chin State in Myanmar (Tanasevitch, 2017). The new species is clearly distinguished by the shape of the dorsal apophysis on the palpal tibia, as well as by the relatively large and not curved radix.

Distribution: Only known from the type locality in northern India.

ACKNOWLEDGEMENTS

I am grateful to Peter Schwendinger (MHNG) for the opportunity to work with the collections of the MHNG and for his valuable comments on the manuscript. Thanks also go to Shou-Wang Lin (University of Greifswald, Germany) for reviewing the manuscript.

REFERENCES

- Förster A., Bertkau P. 1883. Beiträge zur Kenntniss der Spinnenfauna der Rheinprovinz. *Verhandlungen des Naturhistorischen Vereins der Preussischen Rheinlande und Westfalens* 40: 205-278.
- Helsdingen P.J. van 1965. Sexual behaviour of *Lepthyphantes leprosus* (Ohlert) (Araneida, Linyphiidae), with notes on the function of the genital organs. *Zoologische Mededelingen* 41: 15-42.
- Hormiga G. 2000. Higher level phylogenetics of erigonine spiders (Araneae, Linyphiidae, Erigoninae). *Smithsonian Contributions to Zoology* 609: 1-160.
- Merrett P. 1963. The palpus of male spiders of the family Linyphiidae. *Proceedings of the Zoological Society of London* 140: 347-467.
- Saaristo M. I. 1977. Secondary genital organs in the taxonomy of Lepthyphantinae (Araneae, Linyphiidae). *Reports from the Department of Zoology, University of Turku* 5: 1-16.
- Tanasevitch A.V. 1998. *Gorbothorax* n. gen., a new linyphiid spider genus from the Nepal Himalayas (Arachnida, Araneae, Linyphiidae). *Bonner Zoologische Beiträge* 47: 421-428.
- Tanasevitch A.V. 2015. Notes on the spider genus *Oedothorax* Bertkau, 1883 with description of eleven new species from India (Linyphiidae: Erigoninae). *Revue suisse de Zoologie* 122(2): 381-398.
- Tanasevitch A.V. 2016. A case of disjunct montane linyphiid species (Araneae) in the Palaeotropics, with notes on synonymy and the description of a new species. *Revue suisse de Zoologie* 123(2): 235-240.
- Tanasevitch A.V. 2017. New genera and new species of the family Linyphiidae from Borneo, Sumatra and Java (Arachnida, Araneae). *Revue suisse de Zoologie* 124(1): 141-155.