

## **Development of 22 Microsatellite Markers for Assessing Hybridization in the Genus Gekko (Squamata: Gekkonidae)**

Authors: Okamoto, Kota, Kurita, Takaki, Nagano, Masahiro, Sato, Yukuto, Aoyama, Hiroaki, et al.

Source: Current Herpetology, 39(1) : 66-74

Published By: The Herpetological Society of Japan

URL: <https://doi.org/10.5358/hsj.39.66>

---

BioOne Complete ([complete.BioOne.org](https://complete.BioOne.org)) is a full-text database of 200 subscribed and open-access titles in the biological, ecological, and environmental sciences published by nonprofit societies, associations, museums, institutions, and presses.

Your use of this PDF, the BioOne Complete website, and all posted and associated content indicates your acceptance of BioOne's Terms of Use, available at [www.bioone.org/terms-of-use](https://www.bioone.org/terms-of-use).

Usage of BioOne Complete content is strictly limited to personal, educational, and non - commercial use. Commercial inquiries or rights and permissions requests should be directed to the individual publisher as copyright holder.

---

BioOne sees sustainable scholarly publishing as an inherently collaborative enterprise connecting authors, nonprofit publishers, academic institutions, research libraries, and research funders in the common goal of maximizing access to critical research.

## Development of 22 Microsatellite Markers for Assessing Hybridization in the Genus *Gekko* (Squamata: Gekkonidae)

KOTA OKAMOTO<sup>1\*</sup>, TAKAKI KURITA<sup>2</sup>, MASAHIRO NAGANO<sup>3</sup>,  
YUKUTO SATO<sup>4</sup>, HIROAKI AOYAMA<sup>5</sup>, SEIKOH SAITOH<sup>6</sup>,  
NAOYA SHINZATO<sup>7</sup>, AND MAMORU TODA<sup>7</sup>

<sup>1</sup>Graduate School of Engineering and Science, University of the Ryukyus, Nishihara, Okinawa 903–0213, JAPAN

<sup>2</sup>Chiba Biodiversity Center, 955–2 Aoba-cho, Chuo-ku, Chiba 260–8682, JAPAN

<sup>3</sup>Faculty of Science and Technology, Oita University, 700 Dannoharu, Oita-shi, Oita 870–1192, JAPAN

<sup>4</sup>Center for Strategic Research Project, Organization for Research Promotion, University of the Ryukyus, Nishihara, Okinawa 903–0213, JAPAN

<sup>5</sup>Research Planning Office, Organization for Research Promotion, University of the Ryukyus, Nishihara, Okinawa 903–0213, JAPAN

<sup>6</sup>College of Economics and Environmental Policy, Okinawa International University, 2–6–1 Ginowan, Ginowan-shi, Okinawa 901–2701, JAPAN

<sup>7</sup>Tropical Biosphere Research Center, University of the Ryukyus, Nishihara, Okinawa 903–0213, JAPAN

**Abstract:** *Gekko yakuensis* and *G. tawaensis*, both endemic to western Japan, are threatened by genetic introgression from *G. hokouensis* and *G. japonicus*, respectively. To know detailed situation of their hybridizations for planning relevant conservation measures, development of sensitive genetic markers is desired. We here developed microsatellite markers based on the sequences obtained from *G. hokouensis* using 454 GS Junior sequencer, and tested stability of PCR amplification and species-specificity of alleles at each locus using *G. hokouensis*, *G. yakuensis*, *G. japonicus*, and *G. tawaensis*. The results showed that 22 loci were almost constantly amplified in more than one species. We further confirmed that there were fixed or nearly fixed allelic displacement between *G. hokouensis* and *G. yakuensis*, and between *G. japonicus* and *G. tawaensis* at 14 loci. Thus, these 22 loci are considered to be useful for evaluation of hybridizations between these pairs of species.

**Key words:** *Gekko hokouensis*; Hybridization; Microsatellite; Diagnostic markers; Population genetics

### INTRODUCTION

Eight species of the genus *Gekko*, including two putative undescribed species, are known from Japan (Tokunaga, 1988; Ota, 1989;

---

\* Corresponding author.

E-mail address: k9.canislupus@gmail.com

Toda et al., 2001a, 2008). *Gekko yakuensis* and *G. tawaensis* are endemic to the southern part of Kyushu and the Osumi Island Group, and the coastal areas facing to the Seto Inland Sea and the remaining coastal area of Shikoku, respectively. By use of allozyme method, Toda et al. (2001b, 2006) detected natural hybridizations between *G. hokouensis* and *G. yakuensis*, and between *G. japonicus* and *G. tawaensis*, each in several areas where the two species involved occur sympatrically. Habitat alterations, such as coastal revetment, have been facilitating the spread of *G. hokouensis* and *G. japonicus*, both prefer open habitat, and might have induced introgressive hybridizations with *G. yakuensis* and *G. tawaensis*, respectively. For this reason, *G. yakuensis* and *G. tawaensis* are listed in the Red List of Japan as Vulnerable and Near Threatened, respectively (Ministry of the Environment, Japan, 2014). For effective conservation managements of the two species, it is necessary to know detailed situation of their hybridizations with those recently spreading species. Although the effectiveness of allozyme genotyping for detection of hybridization was elucidated in the previous studies, the method also has undesirable features in view of conservation purposes, for instance, necessities in collecting relatively large and fresh tissue samples by much damaging or even killing animals and in keeping tissues frozen with high cost. Therefore, it is desired to develop genetic markers that can detect hybrid genotypes less invasively with less cost.

Microsatellite is a popular molecular marker in population genetics and is disseminated by recent technical advances in their development and genotyping (Guichoux et al., 2011). Microsatellite markers in *Gekko* have ever been reported for *G. swinhonis* (Li and Zhou, 2007) and *G. japonicus* (Wei et al., 2015). We tested several of the markers for samples of *G. hokouensis* and *G. yakuensis*, but most of them were not amplified nor variable between these two species. Hence, we have developed another series of microsatel-

lite markers that is suitable in assessing states of hybridizations for each of the two species pairs.

## MATERIALS AND METHODS

Total genomic DNA was extracted from liver tissue of *G. hokouensis* from Yonaguni-jima Island (Zoological Collection of the Kyoto University Museum: KUZ R71161) using standard phenol–chloroform method, and was used for isolation of microsatellite loci. Microsatellite enrichment was conducted using 3' biotin-labeled oligonucleotide probes–magnetic bead complex following Glenn and Schable (2005) and Kurita et al. (2013). After recovering DNA concentration by PCR, the PCR products with length of 300–800 bp were obtained from the gel and purified by QIAquick Gel Extraction Kit (Qiagen). The amplicon was prepared with the GS Rapid Library Preparation Kit (Roche), and pyrosequencing was performed with the 454 GS Junior sequencer (Roche) to determine a series of sequences with microsatellite motifs. Procedures for sequencing with 454 GS Junior and screening of candidates of microsatellite loci were same as Kurita et al. (2014).

To develop microsatellite markers that are applicable to many of the *Gekko* species, we checked presence/absence of homologous microsatellite loci in *G. japonicus* draft genome sequences provided by Liu et al. (2015). Although both *G. hokouensis* and *G. japonicus* are assigned to the *japonicus* species group by Rösler et al. (2011), they are phylogenetically distant from each other in this species group, and most of the *Gekko* species distributed in Japan are located between the two species on phylogenetic tree (Toda, 2000). Therefore, microsatellite loci conserved in both *G. hokouensis* and *G. japonicus* are expected to be applicable to other *Gekko* species in Japan. We downloaded the draft genomic sequences of *G. japonicus* (GenBank accession no. LNDG01000000), and conducted homology search of the sequences obtained from *G.*

*hokouensis* against the *G. japonicus* genome using the NCBI BLAST+ program (Camacho et al., 2009). According to the result, we selected 47 loci that had homologs in the *G. japonicus* genome and satisfied other a few conditions (e.g., suitable repeat number of motif, composition by a single motif, enough length of flanking regions, etc.), and designed primers using Primer3Plus (Untergasser et al., 2012). We further tested the primers' specificity with the target region of the *G. japonicus* genome by using the default setting of Primer-BLAST (Ye et al., 2012). In this setting, primers with six or more mismatch bases with the target region and/or those with high sequence similarity (less than two base mismatches) with non-target region/species were judged as "less-specific primers".

The 47 potential loci were assigned to several sets of multiplex PCR system according to expected sizes of PCR products and the multiplex PCR was conducted for several representative samples using Type-it Microsatellite PCR Kit (Qiagen) and fluorescent-tagged universal primers (Blacket et al., 2012). PCR was carried out in 5  $\mu$ l reactions containing 10–100 ng of extracted DNA, 2.5  $\mu$ l of Type-it Multiplex PCR Master Mix, 0.01  $\mu$ M of each tailed forward primer, 0.2  $\mu$ M of each reverse primer, and 0.01  $\mu$ M of each fluorescent-tagged universal primer (Tail A, Tail B, Tail C, and Tail D labelled with 6-FAM, VIC, NED, and PET, respectively). The PCR conditions were as follows: initial denaturation at 95°C for 5 min; 30 cycles of denaturation at 95°C for 30 s, annealing at 60°C for 90 s, and extension at 72°C for 30 s; final extension at 60°C for 30 min. The PCR products were subjected to fragment analysis by ABI3130xl Genetic Analyzer (Applied Biosystems) and sizes of alleles at each locus were determined by Geneious software (version 10.1.3; Biomatters Ltd.) with GeneScan 600LIZ size standard (Applied Biosystems).

Applicability and allele composition at each locus were evaluated using wild samples of *G. hokouensis* (n=20; Kawanabe, Kagoshima Pref.), *G. yakuensis* (n=20; Tanegashima

Island, Kagoshima Pref.), *G. japonicus* (n=10; Usa, Saeki, and Oita, Oita Pref.), and *G. tawaensis* (n=10; Saeki, Oita Pref.), which were considered to be pure populations without any hybridizations. The samples of *G. hokouensis* and *G. yakuensis* were collected from mutually allopatric populations. Those of *G. japonicus* and *G. tawaensis* were collected from partially sympatric regions, but all of the geckos were considered to be pure individuals on the basis of morphological characters. Based on the genotypes of the above samples, we compared allele frequencies among the four species. We also calculated number of alleles per locus ( $N_A$ ), observed and expected heterozygosities ( $H_O$  and  $H_E$ , respectively) with GenAIEx 6.5 (Peakall and Smouse, 2012) for each sample. Deviations from Hardy-Weinberg equilibrium (HWE) and linkage disequilibrium (LD) were tested using GENEPOP on the web (version 4.2; Raymond and Rousset, 1995; Rousset, 2008). The significance level of the deviation from HWE and LD were adjusted using the sequential Bonferroni procedure (Rice, 1989). Presence of null alleles was tested using Micro-Checker (version 2.2.3; van Oosterhout et al., 2004).

## RESULTS AND DISCUSSION

We obtained 3,740 reads from the enriched library of *G. hokouensis* by a 454 GS Junior run. We detected 161 sequences containing single microsatellite motif and having 50 bp or more flanking regions on both sides of the microsatellite region, but many of these were still unsuitable for designing primers because of inappropriate sequences in the flanking regions. Among the designed 47 pairs of primers whose loci were expected to have homologs in the *G. japonicus* genome, the sequences of 22 primer pairs were judged as "highly specific" to the target regions of the genomic sequences of *G. japonicus*. As a result of the amplification trials using the 47 primer pairs, another set of 22 loci (two tetra-, three tri-, and 17 di-nucleotide loci)

were successfully amplified in more than one of the four species examined (Tables 1 and 2). Fifteen out of the 22 amplified loci were the ones whose primer sequences showed high specificity to the target region of the *G. japonicus* genome. BLAST searches of flanking regions (30 bps) of the 22 isolated loci against DDBJ database confirmed that none of these loci was the same as previously developed microsatellite loci, although only 5' flanking region of Gh147 showed certain similarity with those of Gs217 of *G. swinhonis* (Li and Zhou, 2007) and Hm122 of *Hemidactylus mabouia* (Short and Petren, 2008). We allocated these 22 loci to three multiplex reactions for efficient genotyping (Table 1).

In *G. hokouensis* and *G. yakuensis*, 20 and 16 (out of the 22) loci were mostly well amplified respectively. In *G. hokouensis*,  $N_A$ ,  $H_O$ , and  $H_E$  ranged from 1–8, 0–0.78, and 0–0.80, respectively (Table 2). Significant deviations from HWE were detected in Gh35, Gh71, and Gh147 loci ( $P < 0.05$ ). In *G. yakuensis*,  $N_A$ ,  $H_O$ , and  $H_E$  ranged from 1–6, 0–0.53, and 0–0.72, respectively (Table 2), and significant deviation from HWE was not detected in any locus. No significant LD was detected in any pair of loci in both species. Null alleles were suggested at Gh35, Gh71, and Gh147 in *G. hokouensis* and at Gh147 in *G. yakuensis*. Among the 16 loci that were commonly amplified in both species, allelic displacements were observed between the two species at four loci (Gh27, Gh91, Gh105, Gh143), and remarkable allele frequency differences were observed at other 10 loci (Gh35, Gh64, Gh71, Gh76, Gh82, Gh84, Gh95, Gh144, Gh147, Gh150) (Table 3).

In each of *G. japonicus* and *G. tawaensis*, 17 loci were mostly well amplified, and 16 loci were amplified in common between the two species, though a few individuals of *G. japonicus* could not be genotyped in several loci. In this species,  $N_A$ ,  $H_O$ , and  $H_E$  ranged from 1–4, 0–0.50, and 0–0.59, respectively (Table 2). A significant deviation from HWE was detected in Gh144 locus ( $P < 0.05$ ). In *G. tawaensis*,  $N_A$ ,  $H_O$ , and  $H_E$  ranged from 1–2,

0–0.20, and 0–0.18, respectively (Table 2), and no significant deviation from HWE was detected in any of the 17 loci. No significant LD was detected in any pair of loci in both species. Presence of null alleles was suggested at Gh144 in *G. japonicus*. Among the 16 commonly amplified loci, allelic displacements were observed at 12 loci (Gh27, Gh35, Gh42, Gh50, Gh53, Gh64, Gh70, Gh78, Gh95, Gh143, Gh146, Gh147), and the predominant alleles were different between the two species at other two loci (Gh28, Gh144) (Table 3).

The microsatellite markers developed here are not highly variable in *G. yakuensis* and *G. tawaensis* (Table 2), being consistent with the results of previous allozyme studies (average heterozygosity of *G. hokouensis*, *G. yakuensis*, *G. japonicus*, and *G. tawaensis* were 0.054, 0.017, 0.110, and 0.005, respectively) (Toda et al., 2001b, 2003). Nonetheless, these markers will be still useful for studies of hybridizations because the species-specificity of alleles is sufficiently high (Table 3).

In this study, 11 out of the 22 loci were mostly well amplified in the four species examined. Because all of the 11 loci had primers that were judged as highly specific to the target region of the *G. japonicus* genome, it is considered that the homolog search and specificity evaluation for the candidate primer sequences against the *G. japonicus* genome were effective to develop the loci that have high universality within the genus. Cross-amplifiable microsatellite markers were successfully developed for a variety of taxa by comparison of genome-derived sequences with multiple closely- or distantly-related species (e.g., Gotoh et al., 2013; Wang et al., 2015). Whole genome information becomes available for many non-model species in recent years, and thus it is recommended to add these procedures in developing effective genetic markers when the relevant information is available.

TABLE 1. Twenty-two microsatellite markers developed in *Gekko hokouensis*.

Locus	Repeat motif	Primer sequences (5'-3')	Multiplex	Dye	Accession No.
Gh27	(GT)12	F: Tail A-CTGGGAGAGAATCACTGCAAC R: GAGGGCATCCATGTGTACCT	III	6-FAM	LC490625
Gh28	(GT)10	F: Tail C-TGATAAGGTGACTGCCATCG R: GGAGCTAATCTGCACAAGCA	III	NED	LC490626
Gh34	(AC)6	F: Tail B-TACCCTAAGTCCACCGCAAC R: TCTCCAAAATTAAGCAGATATTCAGA	II	VIC	LC490627
Gh35	(GT)11	F: Tail D-TTTCAGCTGCCAATGTCTTG R: GATTAAGCATTTTCGCCTCCA	III	PET	LC490628
Gh42	(AC)7	F: Tail D-TGGGGTGTATGAAGCTGTGA R: AGCCCAACCTGAAGAAAACCT	I	PET	LC490629
Gh50	(AC)6	F: Tail C-GTCCACTTGGCACAAGTCCT R: TGGGGTCTTGCTAACAGTGA	II	NED	LC490630
Gh53	(AC)10	F: Tail A-GAAATGCCCAAGCTCCAATA R: AATTAAGTACAGAGCCGCATCA	III	6-FAM	LC490631
Gh64	(AC)9	F: Tail D-ACACAGCACAACGAGGAATG R: ATGTGGCTGCAATGATGTGT	II	PET	LC490632
Gh70	(AC)9	F: Tail C-TGCTTGATTTTGGGTTGAGA R: GGGACTTTGCGGAGACAGTA	I	NED	LC490633
Gh71	(AC)16	F: Tail C-GGCACTGCTTCTTTGGGTAA R: GAGAAACGGACTCGAACCCAG	III	NED	LC490634
Gh76	(AC)11	F: Tail D-TAGAGGAGCACCGGAAATGT R: TTCAGGGACCTGCTTGAGAT	III	PET	LC490635
Gh78	(GT)7	F: Tail A-AGGATGGGCGACATGATAAC R: TGTACAACCTCAAGTGCGAAGC	II	6-FAM	LC490636
Gh82	(GT)7	F: Tail A-CTCCGGGAGCTCACTTTATG R: TGCTTTTTACACAGCACTCTCTG	III	6-FAM	LC490637
Gh84	(AC)10	F: Tail A-CACCACTACTCAGCCCAGAA R: GGGAAATCACAGCCTCTTAGG	I	6-FAM	LC490638
Gh91	(GT)8	F: Tail D-GCATCCTTTGGGTTACGTGT R: CGTGTGCTGCTTACACCAAT	I	PET	LC490639
Gh95	(AC)11	F: Tail C-ACAGAGGCTGCCACTTATCC R: TGCGATTGTGCTATCCCATA	II	NED	LC490640
Gh105	(AC)14	F: Tail C-TGCTTAATCTACGTAGGGCAGA R: CCCACCCAGAGCCTCTTCTA	II	NED	LC490641
Gh143	(GTTT)12	F: Tail B-TGTGACCAGTCTGCCACTG R: AAGAGCACCGCACTTTGTTT	I	VIC	LC490642
Gh144	(ACGC)6	F: Tail A-GCAGTCCCAAGTCAATCAT R: GGGGCAAGTTTCTGTGTTGT	I	FAM	LC490643
Gh146	(GTT)7	F: Tail C-GGAGCACATTGAAGGTACGG R: GTCCACACAGCTCCAAAGT	III	NED	LC490644
Gh147	(CCT)7	F: Tail A-CCCGACACTTGTGCCATTAC R: CTGTGGCAAGGTGCTAGAAGA	II	6-FAM	LC490645
Gh150	(ATC)8	F: Tail A-TCCTGGCCACAATCAAGGAC R: AAACCTTTAGCAGTTTTTCTAAGTCT	I	6-FAM	LC490646



TABLE 2. Characteristics of 22 microsatellite loci in four *Gekko* species.

Locus	<i>G. hokouensis</i> (N=20)				<i>G. yakuensis</i> (N=20)				<i>G. japonicus</i> (N=10)				<i>G. tawaensis</i> (N=10)		$N_A$ (all)		
	$N_{\text{Amp}}$	$N_A$	$H_O$	$H_E$	$N_{\text{Amp}}$	$N_A$	$H_O$	$H_E$	$N_{\text{Amp}}$	$N_A$	$H_O$	$H_E$	$N_{\text{Amp}}$	$N_A$		$H_O$	$H_E$
Gh27	19	4	0.21	0.20	20	1	0	0	8	1	0	0	10	1	0	0	7
Gh28	20	1	0	0	—	—	—	—	10	2	0.30	0.26	10	1	0	0	3
Gh34	20	2	0.35	0.40	—	—	—	—	10	1	0	0	—	—	—	—	3
Gh35	19	7	0.26*	0.78	20	1	0	0	10	4	0.40	0.35	10	1	0	0	9
Gh42	—	—	—	—	—	—	—	—	9	2	0.11	0.28	10	1	0	0	3
Gh50	20	1	0	0	16	1	0	0	10	1	0	0	10	1	0	0	3
Gh53	20	3	0.25	0.23	—	—	—	—	6	1	0	0	10	1	0	0	5
Gh64	20	3	0.15	0.22	20	1	0	0	10	1	0	0	10	1	0	0	4
Gh70	20	1	0	0	—	—	—	—	10	1	0	0	10	1	0	0	3
Gh71	18	5	0.22*	0.66	19	2	0.05	0.15	6	1	0	0	10	1	0	0	6
Gh76	20	8	0.75	0.72	20	2	0.25	0.29	—	—	—	—	10	2	0.20	0.18	11
Gh78	—	—	—	—	—	—	—	—	10	1	0	0	10	1	0	0	2
Gh82	18	6	0.78	0.71	20	1	0	0	—	—	—	—	—	—	—	—	6
Gh84	20	7	0.75	0.80	20	1	0	0	10	1	0	0	10	1	0	0	8
Gh91	14	1	0	0	20	1	0	0	—	—	—	—	—	—	—	—	2
Gh95	20	2	0.10	0.10	20	1	0	0	10	1	0	0	10	1	0	0	4
Gh105	20	1	0	0	18	1	0	0	—	—	—	—	—	—	—	—	2
Gh143	20	1	0	0	20	3	0.15	0.14	10	1	0	0	10	1	0	0	6
Gh144	20	2	0.05	0.05	14	1	0	0	8	3	0*	0.59	10	1	0	0	5
Gh146	20	2	0.35	0.29	20	1	0	0	10	2	0.50	0.50	10	1	0	0	4
Gh147	11	6	0*	0.78	20	2	0.20	0.50	10	2	0.40	0.32	10	1	0	0	8
Gh150	20	2	0.05	0.05	19	6	0.53	0.72	—	—	—	—	—	—	—	—	7
Average		3.25	0.21	0.30		1.63	0.07	0.11		1.53	0.10	0.13		1.06	0.01	0.01	5.05

$N$ : number of individuals assayed,  $N_{\text{Amp}}$ : number of individuals successfully genotyped,  $N_A$ : number of alleles per locus,  $H_O$  and  $H_E$ : observed and expected heterozygosity. \*Significant deviation from HWE after the sequential Bonferroni correction.

TABLE 3. Allele frequencies in four *Gekko* species at 22 loci. Values for diagnostic alleles (i.e., alleles found only in one of the four species) are given in bold. The predominant alleles are indicated with asterisks.

Locus	Allele	<i>G. hokouensis</i>	<i>G. yakuensis</i>	<i>G. japonicus</i>	<i>G. tawaensis</i>	Locus	Allele	<i>G. hokouensis</i>	<i>G. yakuensis</i>	<i>G. japonicus</i>	<i>G. tawaensis</i>
Gh27	198				<b>1.000*</b>	Gh78	275	—	—		<b>1.000*</b>
	204	<b>0.026</b>					277	—	—	<b>1.000*</b>	
	206	<b>0.053</b>				Gh82	273	0.083	1.000*	—	—
	207		<b>1.000*</b>				277	<b>0.028</b>	—	—	—
	209	<b>0.895*</b>					279	<b>0.194</b>	—	—	—
	211	<b>0.026</b>					281	<b>0.028</b>	—	—	—
	214			<b>1.000*</b>			287	<b>0.444*</b>	—	—	—
Gh28	164	<b>1.000*</b>	—			291	<b>0.222</b>	—	—	—	
	166		—	<b>0.850*</b>		Gh84	241			1.000*	1.000*
168		—	0.150	1.000*	243		0.125	1.000*	—	—	
Gh34	129		—	<b>1.000*</b>	—	253	<b>0.075</b>	—	—	—	
	136	<b>0.725*</b>	—		—	258	<b>0.100</b>	—	—	—	
Gh35	140	<b>0.275</b>	—		—	262	<b>0.100</b>	—	—	—	
	188				<b>1.000*</b>	271	<b>0.325*</b>	—	—	—	
	189	<b>0.211</b>				278	<b>0.050</b>	—	—	—	
	191	0.289*		0.100		289	<b>0.225</b>	—	—	—	
	193	0.263	1.000*			Gh91	183	<b>1.000*</b>	—	—	—
	199	0.079		0.050			187		<b>1.000*</b>	—	—
	201	0.105		0.050		Gh95	152			<b>1.000*</b>	—
	203	<b>0.026</b>					154	<b>0.950*</b>	—	—	—
	205			<b>0.800*</b>			156	0.050	1.000*	—	—
	209	<b>0.026</b>					162				<b>1.000*</b>
324	—	—		<b>1.000*</b>	Gh105		225	<b>1.000*</b>	—	—	—
327	—	—	<b>0.167</b>			243		<b>1.000*</b>	—	—	
Gh50	332	—	—	<b>0.833*</b>		Gh143	193				<b>1.000*</b>
	313	1.000*	1.000*				196		<b>0.025</b>	—	—
Gh53	318			<b>1.000*</b>		197			<b>1.000*</b>	—	
	323				<b>1.000*</b>	200		<b>0.050</b>	—	—	
	125	<b>0.050</b>	—			202		<b>1.000*</b>	—	—	
	128		—		<b>1.000*</b>	204			<b>0.925*</b>	—	
	133	<b>0.875*</b>	—			Gh144	175	0.025	1.000*	—	—
	149	<b>0.075</b>	—				179	<b>0.975*</b>	—	—	—
	158		—	<b>1.000*</b>			184			<b>0.500*</b>	—
Gh64	328				<b>1.000*</b>	188			0.125	1.000*	
	341	0.875*		1.000*		192			<b>0.375</b>	—	
	343	<b>0.025</b>				Gh146	220				<b>1.000*</b>
	346	0.100	1.000*				222	<b>0.175</b>	—	—	—
Gh70	143		—	<b>1.000*</b>		229				<b>0.550*</b>	
	151		—		<b>1.000*</b>	244	0.825*	1.000*	0.450	—	
	153	<b>1.000*</b>	—			Gh147	468			<b>0.800*</b>	—
286			1.000*	1.000*	477		0.091	—	0.200	—	
304	<b>0.083</b>				480		<b>0.364*</b>	—	—	—	
310	0.111	0.921*			483		<b>0.091</b>	—	—	—	
312	0.194	0.079			484					<b>1.000*</b>	
314	<b>0.528*</b>				486		<b>0.091</b>	—	—	—	
323	<b>0.083</b>				489		0.182	0.450	—	—	
Gh76	390	0.025	0.175	—		492	0.182	0.550*	—	—	
	394		<b>0.825*</b>	—		Gh150	400	<b>0.975*</b>	—	—	—
	396	<b>0.225</b>		—			410	0.025	0.132	—	—
	398	<b>0.025</b>		—			413		<b>0.053</b>	—	—
	400	<b>0.100</b>		—			416		<b>0.395*</b>	—	—
	402	<b>0.050</b>		—			419		<b>0.053</b>	—	—
	404	<b>0.450*</b>		—			422		<b>0.316</b>	—	—
	406	<b>0.025</b>		—			425		<b>0.053</b>	—	—
	409	<b>0.100</b>		—							
	410			—	<b>0.900*</b>						
414			—	<b>0.100</b>							



## ACKNOWLEDGMENTS

We thank A. Tominaga (University of the Ryukyus) for his helpful advice and technical support on this study. We also thank two anonymous reviewers for their valuable advices/comments on our manuscript. This study was supported in part by a grant from the Japan Society for the Promotion of Science (JSPS KAKENHI Grant No. JP15K07192), the Spatio-temporal Genomics Project promoted by the University of the Ryukyus, and the Research Project Promotion Grant (Strategic Research Grant) (No. 16SP01302) by the University of the Ryukyus.

## LITERATURE CITED

- BLACKET, M. J., ROBIN, C., GOOD, R. T., LEE, S. F., AND MILLER, A. D. 2012. Universal primers for fluorescent labelling of PCR fragments—an efficient and cost-effective approach to genotyping by fluorescence. *Molecular Ecology Resources* 12: 456–463.
- CAMACHO, C., COULOURIS, G., AVAGYAN, V., MA, N., PAPADOPOULOS, J., BEALER, K., AND MADDEN, T. L. 2009. BLAST+: architecture and applications. *BMC Bioinformatics* 10: 421.
- GLENN, T. C. AND SCHABLE, N. A. 2005. Isolating microsatellite DNA loci. *Methods in Enzymology* 395: 202–222.
- GOTOH, R. O., TAMATE, S., YOKOYAMA, J., TAMATE, H. B., AND HANZAWA, N. 2013. Characterization of comparative genome-derived simple sequence repeats for acanthopterygian fishes. *Molecular Ecology Resources* 13: 461–472.
- GUICHOUX, E., LAGACHE, L., WAGNER, S., CHAUMEIL, P., LÉGER, P., LEPAIS, O., LEPOITTEVIN, C., MALAUSA, T., REVARDEL, E., SALIN, F., AND PETIT, R. J. 2011. Current trends in microsatellite genotyping. *Molecular Ecology Resources* 11: 591–611.
- KURITA, K., HIKIDA, T., AND TODA, M. 2013. Development and characterization of polymorphic microsatellite marker for East Asian species of the genus *Plestiodon*. *Conservation Genetics Resources* 5: 355–357.
- KURITA, T., AOYAMA, H., SAITOH, S., SHINZATO, N., HONDA, M., AND TODA, M. 2014. Development and characterization of 24 microsatellite markers in a eublepharid gecko, *Goniurosaurus kuroiwae*. *Conservation Genetics Resources* 6: 247–249.
- LI, J. AND ZHOU, K. 2007. Isolation and characterization of microsatellite markers in the gecko *Gekko swinhonis* and cross-species amplification in other gekkonid species. *Molecular Ecology Notes* 7: 674–677.
- LIU, Y., ZHOU, Q., WANG, Y., LUO, L., YANG, J., YANG, L., LIU, M., LI, Y., QIAN, T., ZHENG, Y., LI, M., LI, J., GU, Y., HAN, Z., XU, M., WANG, Y., ZHU, C., YU, B., YANG, Y., DING, F., JIANG, J., YANG, H., AND GU, X. 2015. *Gekko japonicus* genome reveals evolution of adhesive toe pads and tail regeneration. *Nature Communications* 6: 10033.
- MINISTRY OF THE ENVIRONMENT, JAPAN (ed.). 2014. *Red Data Book 2014—Threatened Wildlife of Japan—Volume 3, Reptilia/Amphibia*. Gyosei, Tokyo.
- OTA, H. 1989. A review of the geckos (Lacertilia: Reptilia) of the Ryukyu Archipelago and Taiwan. p. 222–261. In: M. Matsui, T. Hikida, and R. C. Goris (eds.), *Current Herpetology in East Asia*. Herpetological Society of Japan, Kyoto.
- PEAKALL, R. AND SMOUSE, P. E. 2012. GenAlEx 6.5: genetic analysis in Excel. Population genetic software for teaching and research—an update. *Bioinformatics* 28: 2537–2539.
- RAYMOND, M. AND ROUSSET, F. 1995. GENEPOP (version 1.2): population genetics software for exact test and ecumenicism. *Journal of Heredity* 86: 248–249.
- RICE, W. R. 1989. Analyzing tables of statistical tests. *Evolution* 43: 223–225.
- RÖSLER, H., BAUER, A. M., HEINICKE, M. P., GREENBAUM, E., JACKMAN, T., NGUYEN, T. Q., AND ZIEGLER, T. 2011. Phylogeny, taxonomy, and zoogeography of the genus *Gekko* Laurenti, 1768 with the revalidation of *G. reevesii* Gray, 1831 (Sauria: Gekkonidae). *Zootaxa* 2989: 1–50.
- ROUSSET, F. 2008. GENEPOP'007: a complete re-implementation of the GENEPOP software

- for Windows and Linux. *Molecular Ecology Resources* 8: 103–106.
- SHORT, K. H. AND PETREN, K. 2008. Isolation and characterization of 12 polymorphic microsatellite markers in the tropical house gecko (*Hemidactylus mabouia*). *Molecular Ecology Resources* 8: 1319–1321.
- TODA, M. 2000. Taxonomy, phylogeny, and biogeography of the genus *Gekko* (Reptilia: Squamata) in Japan: a biochemical approach. *Unpublished doctoral dissertation*. Kyoto University, Kyoto.
- TODA, M., HIKIDA, T., OKADA, S., AND OTA, H. 2003. Contrasting patterns of genetic variation in the two sympatric geckos *Gekko tawaensis* and *G. japonicus* (Reptilia: Squamata) from western Japan, as revealed by allozyme analyses. *Heredity* 90: 90–97.
- TODA, M., HIKIDA, T., AND OTA, H. 2001a. Discovery of sympatric cryptic species within *Gekko hokouensis* (Gekkonidae: Squamata) from the Okinawa Islands, Japan, by use of allozyme data. *Zoologica Scripta* 30: 1–11.
- TODA, M., OKADA, S., HIKIDA, T., AND OTA, H. 2006. Extensive natural hybridization between two geckos, *Gekko tawaensis* and *Gekko japonicus* (Reptilia: Squamata), throughout their broad sympatric area. *Biochemical Genetics* 44: 1–17.
- TODA, M., OKADA, S., OTA, H., AND HIKIDA, T. 2001b. Biochemical assessment of evolution and taxonomy of the morphologically poorly diverged geckos, *Gekko yakuensis* and *G. hokouensis* (Reptilia: Squamata) in Japan, with special reference to their occasional hybridization. *Biological Journal of the Linnean Society* 73: 153–165.
- TODA, M., SENGOKU, S., HIKIDA, T., AND OTA, H. 2008. Description of two new species of the genus *Gekko* (Squamata: Gekkonidae) from the Tokara and Amami Island Groups in the Ryukyu Archipelago, Japan. *Copeia* 2008: 452–466.
- TOKUNAGA, S. 1988. A gecko of the genus *Gekko* from Taka-shima Island, Hirado, Nagasaki, Japan (Reptilia: Lacertilia). *Japanese Journal of Herpetology* 12: 127–130.
- UNTERGASSER, A., CUTCUTACHE, I., KORESSAAR, T., YE, J., FAIRCLOTH, B. C., REMM, M., AND ROZEN, S. G. 2012. Primer3—new capabilities and interfaces. *Nucleic Acids Research* 40: e115.
- VAN OOSTERHOUT, C., HUTCHINSON, W. F., WILLS, D. P. M., AND SHIPLEY, P. 2004. MICRO-CHECKER: software for identifying and correcting genotyping errors in microsatellite data. *Molecular Ecology Notes* 4: 535–538.
- WANG, Q., FANG, L., CHEN, J., HU, Y., SI, Z., WANG, S., CHANG, L., GUO, W., AND ZHANG, T. 2015. Genome-wide mining, characterization, and development of microsatellite markers in *Gossypium* species. *Scientific Reports* 5: 10638.
- WEI, L., SHAO, W. W., ZHOU, H. B., PING, J., LI, L. M., AND ZHANG, Y. P. 2015. Rapid microsatellite development in *Gekko japonicus* using sequenced restriction-site associated DNA markers. *Genetics and Molecular Research* 14: 14119–14122.
- YE, J., COULOURIS, G., ZARETSKAYA, I., CUTCUTACHE, I., ROZEN, S., AND MADDEN, T. 2012. Primer-BLAST: a tool to design target-specific primers for polymerase chain reaction. *BMC Bioinformatics* 13: 134.

---

Accepted: 19 October 2019