



## **A New Ventrops (Diptera: Rhinophoridae) from Ethiopia**

Authors: Cerretti, Pierfilippo, Ziegler, Joachim, and Pape, Thomas

Source: African Invertebrates, 56(3) : 579-583

Published By: KwaZulu-Natal Museum

URL: <https://doi.org/10.5733/afin.056.0306>

---

BioOne Complete (complete.BioOne.org) is a full-text database of 200 subscribed and open-access titles in the biological, ecological, and environmental sciences published by nonprofit societies, associations, museums, institutions, and presses.

Your use of this PDF, the BioOne Complete website, and all posted and associated content indicates your acceptance of BioOne's Terms of Use, available at [www.bioone.org/terms-of-use](http://www.bioone.org/terms-of-use).

Usage of BioOne Complete content is strictly limited to personal, educational, and non-commercial use. Commercial inquiries or rights and permissions requests should be directed to the individual publisher as copyright holder.

---

BioOne sees sustainable scholarly publishing as an inherently collaborative enterprise connecting authors, nonprofit publishers, academic institutions, research libraries, and research funders in the common goal of maximizing access to critical research.

## A new *Ventrops* (Diptera: Rhinophoridae) from Ethiopia

Pierfilippo Cerretti<sup>1</sup>, Joachim Ziegler<sup>2</sup> and Thomas Pape<sup>3</sup>

<sup>1</sup>DAFNAE – Entomologia, Università degli Studi di Padova, Viale dell'Università 16, 35020 Legnaro (Padova), Italy; pierfilippocerretti@yahoo.it

<sup>2</sup> Museum für Naturkunde, Leibniz Institute for Evolution and Biodiversity Science, Invalidenstrasse 43, 10115 Berlin, Germany; joachim.ziegler@mfn-berlin.de

<sup>3</sup>Natural History Museum of Denmark, University of Copenhagen, Universitetsparken 15, 2100 Copenhagen, Denmark; tpape@snm.ku.dk

### ABSTRACT

A new species of the Afrotropical endemic rhinophorid genus *Ventrops* is described, illustrated and compared with congeners.

KEY WORDS: Afrotropical Region, Diptera, Rhinophoridae, *Ventrops*, new species.

### INTRODUCTION

The Afrotropical genus *Ventrops* was erected by Crosskey (1977) for the single species *V. milichioides* Crosskey on the basis of specimens from Zimbabwe and Kenya, and later Pape (1987) added three new species from South Africa and Tanzania. Recently, Cerretti and Pape (2012) revised *Ventrops*, giving cladistic arguments for a proper genus-level definition (see also Cerretti *et al.* 2014), and they described three more species from Namibia, Tanzania and Ethiopia. The present paper was prompted by the recent collection of three specimens of an undescribed species of *Ventrops* from Ethiopia, and the use of molecular sequence data from this material (Ziegler & Tóthová 2014). In this paper we describe these Ethiopian specimens as a new species in order to have a name for forthcoming studies aimed at elucidating phylogenetic relationships among rhinophorid lineages.

### MATERIAL AND METHODS

Male terminalia were dissected and prepared for examination following the methods described by Cerretti and Pape (2012). Composite focus-stacking images were produced from multiple images captured using a Nikon DS-L1 digital camera (Nikon, Tokyo, Japan) mounted on a Leica MZ12.5 stereomicroscope (head and wing) or on a Leica DMLS compound microscope (terminalia) (both microscopes: Leica, Wetzlar, Germany), and processed with CombineZM (<http://combinezm.software.informer.com/>). Environmental scanning electron microscope (ESEM) digital images were taken with a Hitachi TM1000 (Hitachi, Tokyo, Japan). Morphological terminology essentially follows McAlpine (1981) with a few exceptions as specified by Cerretti *et al.* (2014); measurements follow Cerretti (2010).

Label data of primary types are given verbatim, and the lines on each label are separated by a slash '/'.

Repositories of specimens are given using the following acronyms:

- MZUR – Zoological Museum, 'Sapienza' University of Rome, Rome, Italy;
- ZMHB – Museum für Naturkunde, Leibniz Institute for Evolution and Biodiversity Science, Berlin, Germany;

<http://africaninvertebrates.org>

urn:lsid:zoobank.org:pub:F3860A6D-3D9A-4515-B871-4994E7A50F59

ZMUM – Zoological Museum, Moscow State University, Moscow, Russia.

TAXONOMY

*Ventrops vikhrevi* sp. n.

Fig. 1

**Etymology:** This species name is a patronym for Nikita Vikhrev, who has provided invaluable specimens from his collecting for our studies, including the species newly described here.

**Diagnosis:** The new species *V. vikhrevi* sp. n. is characterised by (i) compound eye large but not enormously developed so that both gena and parafacial are distinct; (ii) posterior eye margin not indented; (iii) parafacial setose (Fig. 1A); (iv) lateral vertical seta well differentiated from postocular row; (v) basicosta yellow; (vi) wing cell  $r_{4+5}$  open (Fig. 1B); (vii) cercus not shortened,  $0.8\times$  as long as surstylus (Fig. 1E, F); and (viii) median process of ventral sclerotisation of distiphallus very long, extending beyond the acrophallus, asymmetrical and apically branching into several, long, finger-shaped sclerites (Fig. 1D).

**Description** (measurements in square brackets refer to the holotype):

*Body length.* 5.1–5.3[5.3] mm.

*Coloration.* Head shiny black covered with light brownish microtomentum only on genal dilation. Palpus blackish brown. Thorax black, covered with weak brownish grey reflecting microtomentum, legs and abdomen shiny black. Halter with yellow stalk and black knob. Upper and lower calypteres whitish. Tegula black. Basicosta yellow. Wing membrane brownish anteriorly and around veins.

*Head* (Fig. 1A). Arista thickened only proximally, covered with microtrichia which are shorter than maximum proximal diameter of arista. First and second aristomeres not elongated. Postpedicel sub-ovoid, about as long as pedicel. Frons at its narrowest point  $1.25\text{--}1.43[1.43]\times$  as wide as compound eye in dorsal view. Ocellar seta well developed, proclinate. One or 2 pairs of latero-clinate setae, behind ocellar triangle, between ocellar and postocellar setae with variable size. Medial vertical seta strong, reclinate. Lateral vertical seta well developed. Usually 3 to 4 strong proclinate orbital setae, the holotype has on the right side only 2 setae. Five to 8 frontal setae (of different thickness and length) descending anteroventrally to level of proximal edge of pedicel (or slightly above). Parafacial with a row of stout, short setulae along its whole length. Parafacial at its narrowest point (i.e. true width, not in strict lateral view),  $[0.8]0.8\text{--}1.0\times$  as wide as postpedicel. Face and lower facial margin not visible in lateral view. Vibrissa well developed. Vibrissal angle receding. Facial ridge strongly concave, with 3 decumbent setulae above vibrissa. Genal dilation well developed, covered with robust setulae. Gena in profile about  $0.25\text{--}0.35[0.30]\times$  as high as compound eye. Postocular setae short. Occiput slightly convex, covered with black setulae. Prementum stubby, not longer than wide. Palpus slightly clavate with some setulae on distal half.

*Thorax.* Prosternum and proepisternal depression bare. Ventral proepimeral seta oriented anteroventrally. Two or 3 postpronotal setae arranged in a straight line. One pair of prescutellar acrostichal setae, 2 + 3 dorsocentral setae, 0 + 2 intra-alar setae and 1 + 3 supra-alar setae (first and third postsutural supra-alar setae very short, second one




Fig. 1. *Ventrops vikhrevi* sp. n. (paratype ♂): (A) head in lateral view; (B) right wing in dorsal view; (C) left pregonite and postgonite in lateral view; (D) distiphallus in lateral view; (E) epandrial complex in posterior view; (F) epandrial complex in lateral view. Scale bars (E, F) = 200  $\mu$ m.

enormous); 1 posthumeral seta, 2 well-developed notopleural setae. Anatergite bare. Two upper katepisternal setae (posterior one clearly longer), plus 3–4 smaller additional setulae and a tuft of setulae in the lower part. One strong and 1–4 short anepimeral

setae. Katepimeron bare or with 1 small seta anteriorly. Scutellum with 2 pairs of strong marginal setae (apicals and subapicals); basal pair absent or barely distinguishable (i.e. at most  $\frac{1}{4}$  the length of subapical setae). Subapicals robust and slightly divergent; apical pair crossed and horizontal, approx.  $\frac{3}{4}$  as long as subapical setae. One pair of discal scutellar setae, in subapical position.

*Wing* (Fig. 1B). Costal spine well differentiated. Base of  $R_{4+5}$  with 2–3 relatively long setulae dorsally. Bend of  $M$  widely obtuse. Section of  $M$  between  $dm-cu$  and bend of  $M$  about  $1.5\times$  as long as section between  $dm-cu$  and  $r-m$ . Cell  $r_{4+5}$  open.

*Abdomen*. Mid-dorsal depression on syntergite 1+2 confined to anterior  $\frac{1}{3}$  of that segment. Syntergite 1+2 with 1 pair of median marginal setae; tergite 3 with 1 pair of erect median marginal setae. Tergite 4 with a complete row of marginal setae; tergite 5 with a row of marginal setae (median pair erect).

*Male terminalia* (Fig. 1C–F). Sternite 5 with a deep posterior median notch. Tergite 6 unmodified, bare. Connection between tergite 6 and syntergosternite 7+8 membranous. Sternite 6 asymmetrical, articulated with segment 7+8 on left side, and attached to it by a membrane on its right side. Cerci, in posterior view, narrow and relatively long, not fused medially. Bacilliform sclerites flattened dorso-ventrally and converging medially. Surstylus well developed, longer than cercus. Bacilliform sclerite firmly fused to base of surstylus. Articulation between surstylus and epandrium membranous. Intermedium not fused to pregonites. Pregonite lobe-like, gently bent forward, with 2 or 3 setulae along posterior margin. Postgonite subtriangular in shape, apically bent forward; submedian anterior seta of postgonite present. Postgonal apodeme present. Epiphallus strong, with simple tip (i.e. not sagittally bilobed). Extension of dorsal sclerite of distiphallus entirely fused medially. Median process of ventral sclerotisation of distiphallus very long, extending beyond acrophallus, asymmetrical and apically branching into several, long, finger-shaped sclerites (Fig. 1D). Dorsal sclerotisation of acrophallus semi-cylindrical, open ventrally. Distiphallus with long and pointed, scale-like spinulae latero-ventrally.

Holotype ♂, Paratypes 2♂: ETHIOPIA: Amhara, Zengena L. / 2530 m, 10.910N 36.965E / 02.VIII.2012 N. Vikhrev. *Holotype*: The second author added a label printed on red paper: "HOLOTYPUS / *Ventrops / vikhrevi* / Cerretti, Ziegler & Pape 2014". The holotype was deposited in ZMUM. *Paratypes*: Both specimens are labelled with "PARATYPUS / *Ventrops / vikhrevi* / Cerretti, Ziegler & Pape 2014" (printed on red paper). One specimen was deposited in MZUR, the other in ZMHB.

#### DISCUSSION

*Ventrops vikhrevi* sp. n. possesses more than two proclinate orbital setae in the male and an acrophallus with a semicylindrical dorsal sclerite, therefore sharing both autapomorphic character states supporting the monophyly of *Ventrops* as recently defined by Cerretti and Pape (2012) and Cerretti *et al.* (2014). Cerretti and Pape (2012) divided the genus into three species groups as follows:

- i) '*freidbergi*' — characterised by two homoplasious apomorphies (postpronotum with three setae;  $M$  vein gradually vanishing on wing membrane where bend of  $M$  should be).
- ii) '*milichioides*' — characterised by two homoplasious (microtrichia of the arista longer than its maximum basal diameter; male tergite 6 divided into two hemitergites) and one nonhomoplasious (posterior margin of compound eye distinctly indented) apomorphies.
- iii) '*hannemariae*' — characterised by one homoplasious (median process of ventral sclerotisation of distiphallus longitudinally divided) and one non-homoplasious



(median process of ventral sclerotisation of distiphallus ending far beyond tip of acrophallus) apomorphies.

*Ventrops vikhrevi* sp. n. does not share any of the apomorphies supporting the monophyly of the 'freidbergi' and 'milichioides' groups. Instead it has the median process of the ventral sclerotisation of distiphallus both divided longitudinally and very long, thus suggesting inclusion in the 'hannemariae' group. *Ventrops vikhrevi* sp. n. is further characterised by having the median process of the ventral sclerotisation of distiphallus strongly asymmetrical. A strongly asymmetrical ventral sclerotisation of distiphallus emerged as a non-homoplasious synapomorphy shared by *V. hannemariae* and *V. intermedius* in the analysis of Cerretti and Pape (2012), and in the absence of conflicting evidence this will support an inclusion of *V. vikhrevi* sp. n. in the 'hannemariae' group. While *V. hannemariae* and *V. intermedius* have the asymmetrical portion of the ventral sclerite harpoon-shaped (see figs in Cerretti & Pape 2012), that of *V. vikhrevi* sp. n. branches distally into several, long, finger-shaped sclerites (Fig. 1D). This is likely to be an apomorphic condition, but as it can be derived from the harpoon-shaped condition we cannot assess the relative position of the three species currently included in this group.

#### REFERENCES

- CERRETTI, P. 2010. *I tachinidi della fauna italiana* (Diptera Tachinidae), con chiave interattiva dei generi ovest-paleartici. Verona: Cierre Edizioni.
- CERRETTI, P., LO GIUDICE, G. & PAPE, T. 2014. Remarkable Rhinophoridae in a growing generic genealogy (Diptera: Calyptratae, Oestroidea). *Systematic Entomology* **39** (4): 660–690. DOI: 10.1111/syen.12080.
- CERRETTI, P. & PAPE, T. 2012. Phylogenetics and taxonomy of *Ventrops* – the largest genus of Afrotropical Rhinophoridae (Diptera). *Invertebrate Systematics* **26** (3): 274–292. DOI: 10.1071/IS12001.
- CROSSKEY, R.W. 1977. A review of the Rhinophoridae (Diptera) and a revision of the Afrotropical species. *Bulletin of the British Museum (Natural History), Entomology Series* **36**: 1–66.
- MCALPINE, J.F. 1981. Morphology and terminology — adults. In: McAlpine, J.F., Peterson, B.V., Shewell, G.E., Teskey, H.J., Vockeroth, J.R. & Wood D.M., eds, *Manual of Nearctic Diptera*, Vol. 1. Research branch, Agriculture Canada, Monograph **27**: 9–63.
- PAPE, T. 1987. Species of the Afrotropical genus *Ventrops* Crosskey (Diptera: Rhinophoridae). *Annals of the Natal Museum* **28** (2): 543–550.
- ZIEGLER, J. & TÓTHOVÁ, A. 2014. What is a Rhinophoridae? A new perspective on an old question – based on DNA sequences. In: Dorchin, N., Kotrba, M., Mengual, X. & Menzel, F., eds, 8<sup>th</sup> International Congress of Dipterology, Potsdam 2014, Abstract volume: 418.

