



OCCURRENCE OF (*Dirofilaria immitis*) IN GRAY FOX (*Urocyon cinereoargenteus*) IN ALABAMA AND GEORGIA 1 2 3

Author: SIMMONS, JEANNE M.

Source: Journal of Wildlife Diseases, 16(2) : 225-228

Published By: Wildlife Disease Association

URL: <https://doi.org/10.7589/0090-3558-16.2.225>

BioOne Complete (complete.BioOne.org) is a full-text database of 200 subscribed and open-access titles in the biological, ecological, and environmental sciences published by nonprofit societies, associations, museums, institutions, and presses.

Your use of this PDF, the BioOne Complete website, and all posted and associated content indicates your acceptance of BioOne's Terms of Use, available at www.bioone.org/terms-of-use.

Usage of BioOne Complete content is strictly limited to personal, educational, and non - commercial use. Commercial inquiries or rights and permissions requests should be directed to the individual publisher as copyright holder.

BioOne sees sustainable scholarly publishing as an inherently collaborative enterprise connecting authors, nonprofit publishers, academic institutions, research libraries, and research funders in the common goal of maximizing access to critical research.

OCCURRENCE OF (*Dirofilaria immitis*) IN GRAY FOX (*Urocyon cinereoargenteus*) IN ALABAMA AND GEORGIA [□] [□] [□]

JEANNE M. SIMMONS, [□] School of Veterinary Medicine, Auburn University, Auburn, Alabama 36830, USA.
WALLER S. NICHOLSON, EDWARD P. HILL and DARREL B. BRIGGS, Alabama Cooperative Wildlife Research Unit, Auburn, Alabama 36830, USA.

Abstract: The occurrence of *Dirofilaria immitis* in gray fox (*Urocyon cinereoargenteus*) was determined by gross examination, blood samples, and filtered tissue sediments in a sampling of 149 gray foxes taken from Alabama and Georgia during the 1977-78 trapping season. Microfilariae were not found in blood samples obtained from 24 of these gray foxes. Three of 82 male foxes (3.7%) and 1 of 67 female foxes (1.5%) were infected with heartworm. *D. immitis* rate of infection was 1 of 19 (5.3%) and 3 of 130 (2.3%), respectively, in juvenile and adult gray foxes. Single sex infections with *D. immitis* occurred in 4 of the 6 foxes, with a maximum nematode burden of approximately eight. Two other infected foxes were encountered separate from this study.

INTRODUCTION

Information obtained by a number of investigators indicates that *Dirofilaria immitis*, the canine heartworm, is found in a number of wild animals throughout North America. Although most of the work has involved wild Canidae, such as the coyote (*Canis latrans*),^{2,7,9,12,20} timber wolf (*Canis occidentalis*),¹ red fox (*Vulpes fulva*),^{2,12,17,19} and gray fox (*Urocyon cinereoargenteus*),^{2,12,14,15,18} a few reports include positive findings in the beaver (*Castor canadensis*),⁵ wolverine (*Gulo luscus*),²² muskrat (*Ondatra zibethica*),⁸ raccoon (*Procyon lotor*),^{6,10} and black bear (*Ursus americanus*).³

In general, a low rate of heartworm infection has been reported for most free-ranging wild canid populations, however

infection rates of 28% (11 of 39) were reported in red foxes in Michigan¹⁹ and eight percent (11 of 133) in Kansas and Colorado coyotes.⁹ A search of the literature revealed reports of only nine specimens of gray fox infected with *D. immitis*. Microfilariae were not found in the blood of 48 gray foxes in a study at Fort Stewart, Georgia, in 1963.²¹ In Minnesota, 26 gray fox carcasses were examined and no heartworms were found.⁴ Kazacos and Edberg¹¹ found 3 of 81 (3.7%) gray foxes in Indiana infected with *D. immitis*. Crowell *et al.*² found heartworm infection in 2 of 20 gray foxes from Louisiana. Microfilariae were found in the blood of one and adult *D. immitis* were found in the heart of the other. Stone reported heartworms and microfilariae from a gray fox in New York,¹⁸ and Miller and Harkema¹⁴ found

[□] A joint contribution of the Department of Microbiology, Auburn University School of Vet. Med. and the Alabama Cooperative Wildlife Research Unit (U.S. Fish and Wildlife Service, Wildlife Management Institute, Game and Fish Division, Alabama Department of Conservation and Natural Resources and Alabama Agricultural Experiment Station, Auburn University cooperating).

[□] Publication No. 1351 School of Veterinary Medicine, Auburn University.

[□] Funded through the Game and Fish Division, Alabama Department of Conservation and Natural Resources Pittman-Robertson Project W-44-V.

[□] Present address: P. O. Box 862, Hillsborough, North Carolina 27278, USA.

heartworms in 2 of 24 gray foxes collected in North Carolina, South Carolina and Georgia. Heartworm was found in a gray fox in Michigan during post-mortem examination (Stuht unpubl.).

The purpose of this study was to investigate the occurrence of heartworm infection in isolated populations of gray foxes in Alabama and Georgia. Gray fox carcasses were obtained from a region known to have a high prevalence of heartworm infection in dogs.

MATERIALS AND METHODS

Gray fox carcasses were obtained primarily from trappers in Alabama and Georgia from 19 November 1977 through 28 February 1978. Counties involved were Chambers, Monroe, Lee, Coffee, Elmore, Barbour, Tallapoosa, Crenshaw, Russel and Macon in Alabama; and Troup and Meriwether in Georgia. Since most of the foxes were obtained from trappers, all but six examined during this study had been skinned and then frozen. A few of the foxes had been killed on roads. Gray fox population densities in the areas sampled are believed to be similar to those (3 to 4 per ha) reported for northern Florida¹³ and found in current telemetry studies of the gray fox in eastcentral Alabama.

Blood samples from 24 locally trapped foxes were taken by cardiac puncture immediately following death. The modified Knott's¹² test was used to determine the presence of microfilariae in these samples.

Fox skulls were retained for age determination using a combination of skull suture closure and toothwear. Carcasses were classified as juvenile, less than one year of age, or adult.

Gross examination was made of internal organs with specific emphasis on the heart and major blood vessels. The atria, ventricles, and vessels were incised and examined for *D. immitis*. After gross examination the lungs and liver were incised, washed and the filtered sediment examined for parasites. The washings were filtered through a 149 μ m wire mesh sieve and the sediment examined with the aid of a dissecting microscope. All parasites found were preserved in 70% ethyl alcohol.

Nematodes suspected of being *D. immitis* were examined further after clearing in a 20% lactophenol solution for 24 h. Anterior portions of females were crushed and examined with the aid of a microscope to determine if microfilariae were present. These nematodes were considered to be *D. immitis* based on size, morphology and location.

TABLE 1. Prevalence of *Dirofilaria immitis* infection in carcasses of 149 trapped gray foxes.

Age class and sex	Number examined	Number positive (%)
Adults		
Male	75	2 (2.7)
Female	55	1 (1.8)
subtotal	130	3 (3.0)
Juveniles		
Male	7	1 (14.5)
Female	12	0 (0)
subtotal	19	1 (5.3)
Total	149	4 (2.7)

RESULTS

Blood samples from 24 gray foxes were negative for circulating microfilariae. Nineteen of these foxes were examined at necropsy and also found negative for adult *D. immitis*.

Of 130 adult gray foxes examined, 3 (2%) contained adult *D. immitis* in the right ventricle and pulmonary arteries; an immature heartworm was found in the right ventricle in 1 of 19 (5%) juveniles.

Of the 149 gray foxes examined, 82 were male and 67 female. Three males (4%) and 1 female (1%) were infected with *D. immitis* (Table 1).

Two adult male gray foxes collected independently of this study contained six and eight *D. immitis*, respectively, in the right ventricle. One was dead on the road in Adams County, Mississippi (M.E. Ferriss, Pers. Comm.) and the other was trapped in Tallapoosa County, Alabama and held captive for 3 months (R.D. Powers, Pers. Comm.).

In this study the prevalence of infection was too low to statistically evaluate the significance of geographic distribution, sex or age.

Acknowledgements

We thank Dr. D. L. Diamond, Department of Pathology and Parasitology, and Dr. Paul R. Schnurrenberger, Department of Microbiology, both from Auburn University, School of Veterinary Medicine, and Dr. Annie K. Prestwood of the University of Georgia, College of Veterinary Medicine, Department of Parasitology, for their help and advice.

LITERATURE CITED

1. COFFIN, D.L. 1944. A case of *Dirofilaria immitis* infection in a captive bred timber wolf (*Canis occidentalis*, Richardson) N. Am. Vet. 25: 611-612.
2. CROWELL, W.A., T.R. KLEI and D.I. HALL. 1977. Occurrence of *Dirofilaria immitis* and associated pathology in coyotes and foxes from Louisiana. In: Proc. Heartworm Symposium, 1977. p. 10-13.
3. CRUM, J.M., V.F. NETTLES and W.R. DAVIDSON. 1978. Studies on endoparasites of the black bear (*Ursus americanus*) in the southeastern United States. J. Wildl. Dis. 14: 178-186.
4. ERICKSON, A.B. 1944. Helminths of Minnesota Canidae in relation to food habits, and a host list and key to the species reported from North America. Am. Midl. Nat. 32: 358-372.

DISCUSSION

All of the infections detected in this survey were single-sex infections. These data, and the rate of infection suggest that gray foxes presently are not reservoirs for *D. immitis* in Alabama and Georgia.

A high percentage of the dog population in this region is infected with *D. immitis*. Studies on dogs in animal shelters from Houston County, Alabama in 1973, indicated 73 of 98 (74%) infected; in 1978 14 of 33 (44%) from the same source were infected. (A.R. Dillon, Pers. Comm.).

Wild animals should not necessarily be considered as "dead end" hosts. *D. immitis* infection within wild mammal populations may have serious consequences that are as yet unknown. If the human population and their dogs continue their urban dispersion, the interaction of *D. immitis* in gray fox and dog populations possibly could become significant, perhaps to the detriment of one or both species (Stuht, unpubl.). If *D. immitis* is pathogenic to gray foxes, infection could be a detriment to survival of local populations of foxes.

5. FOIL, L. and T.C. ORIHHEL. 1975. *Dirofilaria immitis* (Leidy, 1856) in the beaver, *Castor canadensis*. J. Parasit. 61: 433.
6. FOX, M. 1944. Heartworms in a coon. Ft. Dodge Biochem. Rev. 12(2): 8.
7. FRANSEN, J.C., R.D. JORGENSEN and E.K. BOGGESS. 1976. Dirofilariasis in Iowa coyotes. J. Wildl. Dis. 12: 165-166.
8. GOBLE, F.C. 1942. Dog heartworms in the muskrat in New York. J. Mammal. 23: 346.
9. GRAHAM, J.M. 1975. Filariasis in coyotes from Kansas and Colorado. J. Parasit. 61: 513-516.
10. HERMAN, C.M. and D.L. PRICE. 1965. Epizootiologic studies of filarioids of the raccoon. J. Wildl. Manage. 29: 694-699.
11. KAZACOS, K.R. and E.O. EDBERG. 1979. *Dirofilaria immitis* infection in foxes and coyotes in Indiana. J. Am. vet. med. Ass. 175: 909-910.
12. KNOTT, J. 1939. A method for making microfilarial surveys on day blood. Trans. Roy. Soc. Trop. Med. and Hyg. 33: 191-196.
13. LORD, R.D., JR. 1961. A population study of the gray fox. Am. Midl. Nat. 66: 87-109.
14. MILLER, G.C. and R. HARKEMA. 1968. Helminths of some wild mammals in the southeastern United States. Proc. Helm. Soc. Wash. 35: 118-125.
15. MONSON, R.A., W.B. STONE and B.C. WEBER. 1973. Heartworms in foxes and wild canids in New York. N.Y. Fish and Game J. 20: 48-53.
16. OTTO, G.F. 1974. Occurrence of the heartworm in unusual locations and in unusual hosts. In: *Proc. of the Heartworm Symposium '74*. Bonner Springs, VM Publishing Inc., 1975. p. 613.
17. SCHLOTTHAUER, J.C. 1964. *Dirofilaria immitis* in the red fox (*Vulpes fulva*) in Minnesota. J. Parasit. 50(6): 801-802.
18. STONÉ, W.B. 1974. Heartworms and microfilaria in a gray fox. N.Y. Fish and Game J. 21: 87.
19. STUHT, J.N. and W.G. YOUATT. 1972. Heartworms and lung flukes from red foxes in Michigan. J. Wildl. Manage. 36: 166-170.
20. THORNTON, J.E., R.R. BELL and M.J. REARDON. 1974. Internal parasites of coyotes in southern Texas. J. Wildl. Dis. 10: 232-236.
21. WALTON, B.C., J.E. GLOVER and R.W. UPHAM. 1963. Lack of *Dirofilaria immitis* infection in wild foxes from an enzootic area. J. Parasit. 49: 526.
22. WILLIAMS, J.F. and A.W. DADE. 1976. *Dirofilaria immitis* infection in a wolverine. J. Parasit. 62: 174-175.
23. WOOD, J.E. 1958. Age structure and productivity of a gray fox population. J. Mammal. 39: 74-86.

Received for publication 21 May 1979