



PARASITES, DISEASES, AND HEALTH STATUS OF SYMPATRIC POPULATIONS OF SIKA DEER AND WHITE-TAILED DEER IN MARYLAND AND VIRGINIA 1

Authors: Davidson, William R., and Crow, Church B.

Source: Journal of Wildlife Diseases, 19(4) : 345-348

Published By: Wildlife Disease Association

URL: <https://doi.org/10.7589/0090-3558-19.4.345>

BioOne Complete (complete.BioOne.org) is a full-text database of 200 subscribed and open-access titles in the biological, ecological, and environmental sciences published by nonprofit societies, associations, museums, institutions, and presses.

Your use of this PDF, the BioOne Complete website, and all posted and associated content indicates your acceptance of BioOne's Terms of Use, available at www.bioone.org/terms-of-use.

Usage of BioOne Complete content is strictly limited to personal, educational, and non - commercial use. Commercial inquiries or rights and permissions requests should be directed to the individual publisher as copyright holder.

BioOne sees sustainable scholarly publishing as an inherently collaborative enterprise connecting authors, nonprofit publishers, academic institutions, research libraries, and research funders in the common goal of maximizing access to critical research.

PARASITES, DISEASES, AND HEALTH STATUS OF SYMPATRIC POPULATIONS OF SIKA DEER AND WHITE-TAILED DEER IN MARYLAND AND VIRGINIA¹

William R. Davidson and Church B. Crow

Southeastern Cooperative Wildlife Disease Study, Department of Parasitology, College of Veterinary Medicine, The University of Georgia, Athens, Georgia 30602, USA

ABSTRACT: In July 1981, investigations on parasites, diseases, and herd health status were conducted on sympatric populations of sika deer (*Cervus nippon*) and white-tailed deer (*Odocoileus virginianus*) from Blackwater National Wildlife Refuge (Maryland) and Chincoteague National Wildlife Refuge (Virginia) on the Delmarva Peninsula. Five adult deer of each species were collected from each location and subjected to thorough necropsy examinations and laboratory tests. White-tailed deer at both locations harbored protozoan, helminth, and arthropod parasites typically associated with this species throughout the southeastern United States. In contrast, sika deer at both locations harbored only light burdens of ticks, chiggers, and sarcocysts. Serologic tests for antibodies to seven infectious disease agents revealed evidence of exposure to bovine virus diarrhea (BVD) virus, infectious bovine rhinotracheitis virus, and parainfluenza₃ virus in white-tailed deer, but only BVD virus in sika deer. At both locations the general health status of sika deer was superior to that of white-tailed deer.

INTRODUCTION

White-tailed deer are the most widely distributed and abundant wild cervids in North America (Halls, 1978), and in many regions, they coexist with other native cervids, domestic livestock, and a variety of introduced exotic ungulates. As a result, numerous investigations have been made on the potential for cross-transmission of infectious agents between white-tailed deer and these hosts. In some instances, e.g., cerebrospinal parelaphostrongylosis, these disease potentials have far-reaching implications (Anderson and Prestwood, 1981).

Although many species have been investigated with regard to disease interactions with white-tailed deer, others including sika deer have not. The single report by Robinson et al. (1978) on elaeophorosis in sika deer in Texas provides the only suggestion of disease interaction between sika and white-tailed deer. This report presents an evaluation of the health status, diseases, and parasites of two sympatric populations of sika and white-tailed deer.

MATERIALS AND METHODS

This study was conducted at two locations: Blackwater National Wildlife Refuge (NWR), Dorchester County, Maryland, and Chincoteague National Wildlife Refuge, Accomack County, Virginia, both of which are on the Delmarva Peninsula. Both locations support sympatric wild populations of white-tailed and sika deer. The history of these sika deer populations has been documented elsewhere (Flyger and Warren, 1958; Flyger, 1960; Flyger and Davis, 1964; Feldhamer et al., 1978; Feldhamer, 1982). Five adult animals (≥ 1 yr of age) of each species were collected at each location by shooting in July 1981. Blood samples were obtained by cardiac puncture immediately after death, and the carcasses were refrigerated prior to necropsy within 12 hr. Detailed necropsy examinations, including parasite recovery and abomasal parasite counts (APC) (Eve and Kellogg, 1977; Nettles, 1981), were performed on each animal. Sarcocyst-infected muscle was fed to laboratory-reared dogs and cats to confirm the identity of the species present as described by Crum and Prestwood (1982). Physical condition was assessed by the method of Stockle et al. (1978). Serum samples were tested for antibodies to a number of infectious disease agents. Where appropriate, lesion specimens were obtained for pathogen isolation attempts. Sections of major organs along with gross lesions were obtained for histopathologic study.

RESULTS

The overall physical condition of sika deer at both locations was superior to that of white-tailed deer. All sika deer were rated in good physical condition, with some approaching the excellent category. White-tailed deer from each location had physical condition ratings of fair,

Received for publication 17 February 1983.

¹ This study was supported by an appropriation from the Congress of the United States. Funds were administered and research coordinated under the Federal Aid in Wildlife Restoration Act (50 Stat. 917) and through Contract Numbers 14-16-0009-81-014 and 14-16-0009-82-500, Fish and Wildlife Service, U.S. Department of the Interior.

TABLE 1. Number of reactors to serologic tests for selected infectious agents in sika and white-tailed deer from two locations on the Delmarva Peninsula.*

Disease agent	Blackwater NWR		Chincoteague NWR	
	Sika deer	White-tailed deer	Sika deer	White-tailed deer
Bovine virus diarrhea virus	2/5 ^b	0/5	0/5	2/5
Infectious bovine rhinotracheitis virus	0/5	0/5	0/5	1/5
Parainfluenza ₃ virus	0/5	0/5	0/5	1/5

* All tests for antibodies to the etiologic agents of leptospirosis, brucellosis, epizootic hemorrhagic disease and bluetongue were negative. Leptospirosis testing included the *pomona*, *grippotyphosa*, *hardjo*, *icterohemorrhagiae*, and *canicola* serotypes. Titers considered positive were: BVD and PI₃ ≥ 1:50 and IBR ≥ 1:2.

^b No. positive/no. tested.

except at Chincoteague NWR where one was rated poor and one was rated good.

Serologic tests for seven infectious agents revealed evidence of exposure to bovine virus diarrhea (BVD) virus, infectious bovine rhinotracheitis (IBR) virus, and parainfluenza₃ (PI₃) virus in white-tailed deer, but only BVD virus in sika deer (Table 1). Necropsy examinations

revealed gross lesions in a single sika and a single white-tailed deer. A 7-yr-old female sika deer from Blackwater NWR had enlarged iliac, inguinal, and cervical lymph nodes, enlarged and hemorrhagic tonsils, and a 1 cm diameter nodule extending from the surface into the parenchyma of the spleen. Histologically, all of these lesions were compatible with a necrotizing granulomatous lymphadenitis. Microbiologic cultures yielded *Staphylococcus aureus* and a beta-*Streptococcus* sp. from lymph nodes and a beta-*Streptococcus* sp., *Sarcina* sp. and *Serratia* sp. from the tonsil. Acid fast stains were negative. A 5-yr-old female white-tailed deer from Chincoteague NWR had massive disseminated abscesses involving the entire left psoas, the left longissimus dorsi and overlying skin, the pelvic cavity, the liver, the right dorsal lobe of the lung, and the right parotid lymph node. Microbiologic cultures of these abscesses revealed mixed infections of beta-*Streptococcus*, *Staphylococcus* sp., and *Aeromonas hydrophila*.

Parasitologic studies revealed 16 species of parasites, including three protozoans, two cestodes, seven nematodes, and four arthropods

TABLE 2. Parasites recovered from sika and white-tailed deer from two locations on the Delmarva Peninsula.*

Parasite	Blackwater NWR		Chincoteague NWR	
	Sika	White-tailed	Sika	White-tailed
Protozoans				
<i>Sarcocystis odocoileocanis</i>	20%, L, L ^b	60%, L, L-M	0	60%, M, L-M
<i>Theileria cervi</i> (77521) ^c	0	100%, L, L-M	0	80%, L, L-M
<i>Trypanosoma cervi</i> (77521)	0	60%, L, L-M	0	60%, L, L
Cestodes				
<i>Taenia hydatigena</i> (77522)	0	20%, 1, 1	0	0
<i>Moniezia benedeni</i> (77523)	0	0	0	20%, 3, 3
Nematodes				
<i>Apteragia odocoilei</i> (77524)	0	100%, 577, 363-834	0	100%, 768, 317-1,361
<i>Ostertagia dikmansii</i> (77525)	0	20%, 49, 49	0	80%, 49, 48-76
<i>Ostertagia mossi</i> (77526)	0	100%, 201, 92-486	0	80%, 112, 48-305
<i>Oesophagostomum venulosum</i> (77527)	0	20%, 1, 1	0	0
<i>Dictyocaulus viviparus</i> (77528)	0	80%, 3, 1-9	0	60%, 1, 1-2
<i>Parelaphostrongylus tenuis</i> (77529)	0	60%, 3, 1-11	0	20%, 1, 1
<i>Setaria yehi</i> (77530)	0	40%, 3, 4-9	0	20%, 1, 1
Protostrongylid larvae	0	80%, L, L-M	0	80%, L, L-M
Arthropods				
<i>Tricholipeurus parallelus</i>	0	20%, L, L	0	20%, L, L
<i>Amblyomma americanum</i>	60%, L, L	100%, L, L-M	100%, M, L-M	100%, M, L-H
<i>Derma-centor variabilis</i>	0	0	40%, L, L	0
<i>Eutrombicula splendens</i> (1983:325)	0	0	80%, L, L	60%, L, L

* Figures in columns are prevalence, mean number per infected animal, and maximum intensity of infection.

^b Actual counts not made; intensities rated as low (L), moderate (M), or high (H).

^c U.S. National Parasite Collection (Beltsville, Maryland) accession number (protozoans and helminths) and Bishop Museum (Honolulu, Hawaii) Collection number (chiggers).

(Table 2). White-tailed deer harbored 15 species, whereas sika deer harbored only four species. Abomasal parasite counts (APC) for white-tailed deer on Blackwater NWR and Chincoteague NWR were 788 and 928, respectively. Only mild lesions were associated with parasitism in white-tailed deer. One had a mild diffuse fibrinous peritonitis due to infection by *Setaria yehi*. Eight had scattered microgranulomas surrounding protostrongylid eggs and larvae in alveolar capillaries. One other had slightly cropped ears and subcutaneous inflammation associated with an infestation of *Amblyomma americanum*. Significant lesions attributable to parasitism were not found in sika deer.

DISCUSSION

Populations of sika deer at both locations could be considered "the picture of health" with substantial reserves of endogenous fat, limited evidence of infectious diseases, and a virtual absence of parasitism. This is in marked contrast to the only previous health investigation conducted on sika deer on the Delmarva Peninsula over two decades earlier. At that time, a large scale die-off of sika deer occurred on James Island due to overpopulation, malnutrition, and subsequent pine oil poisoning when the deer resorted to consuming the needles, bark, and roots of pines (Hayes and Shotts, 1959). Interestingly, examinations for endoparasites by Hayes and Shotts (1959) were also negative.

White-tailed deer at both locations were at a reasonable health status, although compared to sika deer they had limited endogenous fat reserves, evidence of exposure to more infectious diseases, and parasite faunas typical of white-tailed deer in the southeastern United States (Davidson et al., 1981). Abomasal parasite counts (Eve and Kellogg, 1977) and other herd health parameters (Eve, 1981) suggested that white-tailed deer herds at both locations were near carrying capacity and that natural mortality due to disease was of a covert nature (Eve, 1981).

Of particular interest was the occurrence of the meningeal worm (*Parelaphostrongylus tenuis*) in white-tailed deer at both locations. This helminth produces neurologic disease in numerous other wild ungulates (Anderson and Prestwood, 1981); however, its effects in sika deer have not been studied. Fish and Wildlife Service personnel reported that sika deer with signs of neurologic disease occasionally had been

encountered at Chincoteague NWR but no such observations were known for Blackwater NWR. Subsequent to this study, formalin fixed brains and spinal cords of two sika deer with neurologic disorders from Chincoteague NWR were examined. In neither case was evidence of cerebrospinal parelaphostrongylosis found; however, considering the range of cervid hosts in which *P. tenuis* causes clinical disease, the pathogenicity of *P. tenuis* in sika deer deserves further investigation.

General observations made during this study, the history of sika deer population growth on the Delmarva Peninsula (Flyger and Warren, 1958; Flyger, 1960; Flyger and Davis, 1964; Feldhamer et al., 1978; Feldhamer, 1982), and studies on interspecific competition between sika deer and white-tailed deer (Harmel, 1979; Armstrong and Harmel, 1981), all indicate that sika deer routinely outcompete white-tailed deer. Although previous studies (Harmel, 1979; Armstrong and Harmel, 1981) have focused on the importance of nutritional competition between these species, our findings suggest that differential susceptibilities to infectious diseases and parasitism also may be important factors in the interspecies equation.

ACKNOWLEDGMENTS

For assistance during field activities, the authors thank personnel with the U.S. Fish and Wildlife Service at Blackwater and Chincoteague National Wildlife Refuges, the Wildlife Administration of the Maryland Department of Natural Resources, and the Virginia Commission of Game and Inland Fisheries. Dr. J. L. Blue, Diagnostic Assistance Laboratory, College of Veterinary Medicine, The University of Georgia, kindly conducted serologic tests. Drs. A. K. Prestwood and J. M. Crum, Department of Parasitology, College of Veterinary Medicine, The University of Georgia, conducted *Sarcocystis* feeding trials. Mr. R. R. Gerrish, National Veterinary Services Laboratories, APHIS, USDA, and Dr. M. L. Goff, Bishop Museum, assisted with identification of arthropods. Dr. Sam Backos, Animal Health Laboratory, Maryland Department of Agriculture, provided assistance during examinations of the two clinically ill sika deer.

LITERATURE CITED

- ANDERSON, R. C., AND A. K. PRESTWOOD. 1981. Lungworms. *In* Diseases and Parasites of White-

- tailed Deer, W. R. Davidson, F. A. Hayes, V. F. Nettles, and F. E. Kellogg (eds.). Tall Timbers Research Station, Tallahassee, Florida, Misc. Publ. No. 7, pp. 266-317.
- ARMSTRONG, W. E., AND D. E. HARMEL. 1981. Exotic mammals competing with the natives. *Tex. Parks Wildl.* (February), pp. 6-7.
- CRUM, J. M., AND A. K. PRESTWOOD. 1982. Prevalence and distribution of *Sarcocystis* spp. among white-tailed deer of the southeastern United States. *J. Wildl. Dis.* 18: 195-203.
- DAVIDSON, W. R., F. A. HAYES, V. F. NETTLES, AND F. E. KELLOGG (eds.). 1981. Diseases and Parasites of White-tailed Deer. Tall Timbers Research Station, Tallahassee, Florida, Misc. Publ. No. 7, 458 pp.
- EVE, J. H. 1981. Management implication of diseases. *In Diseases and Parasites of White-tailed Deer*, W. R. Davidson, F. A. Hayes, V. F. Nettles, and F. E. Kellogg (eds.). Tall Timbers Research Station, Tallahassee, Florida, Misc. Publ. No. 7, pp. 413-423.
- , AND F. E. KELLOGG. 1977. Management implications of abomasal parasites in white-tailed deer. *J. Wildl. Manage.* 41: 169-177.
- FELDHAMER, G. A. 1982. Sika deer. *In Wild Mammals of North America*, J. A. Chapman and G. A. Feldhamer (eds.). The Johns Hopkins University Press, Baltimore, Maryland, pp. 1114-1123.
- , J. A. CHAPMAN, AND R. L. MILLER. 1978. Sika deer and white-tailed deer on Maryland's Eastern Shore. *Wildl. Soc. Bull.* 6: 155-157.
- FLYGER, V. F. 1960. Sika deer on islands in Maryland and Virginia. *J. Mammal.* 41: 140.
- , AND N. W. DAVIS. 1964. Distribution of sika deer (*Cervus nippon*) in Maryland and Virginia in 1962. *Chesapeake Sci.* 5: 212-213.
- , AND J. WARREN. 1958. Sika deer in Maryland—An additional big game animal or a possible pest. *Proc. Southeast. Assoc. Fish Wildl. Agencies* 12: 209-211.
- HALLS, L. D. 1978. White-tailed deer. *In Big Game of North America, Ecology and Management*, J. L. Schmidt and D. L. Gilbert (eds.). Stackpole Books, Harrisburg, Pennsylvania, pp. 43-65.
- HARMEL, D. 1979. Preliminary results of axis deer-white-tailed deer and sika deer-white-tailed deer survival study in the Edwards plateau of Texas. Second Annual Meeting Southeast Deer Study Group, Mississippi State, Mississippi, p. 14 (abstract).
- HAYES, F. A., AND E. B. SHOTTS. 1959. Pine oil poisoning in sika deer. *Southeast. Vet.* 10: 34-39.
- NETTLES, V. F. 1981. Necropsy procedures. *In Diseases and Parasites of White-tailed Deer*, W. R. Davidson, F. A. Hayes, V. F. Nettles, and F. E. Kellogg (eds.). Tall Timbers Research Station, Tallahassee, Florida, Misc. Publ. No. 7, pp. 6-16.
- ROBINSON, R. M., L. P. JONES, T. J. GALVIN, AND G. M. HARWELL. 1978. Elaeophorosis in sika deer in Texas. *J. Wildl. Dis.* 14: 136-141.
- STOCKLE, A. W., G. L. DOSTER, AND W. R. DAVIDSON. 1978. Endogenous fat as an indicator of physical condition of southeastern white-tailed deer. *Proc. Southeast. Assoc. Fish Wildl. Agencies* 32: 269-279.