

Dirofilaria immitis in a Raccoon (Procyon lotor)

Authors: Snyder, Daniel E., Hamir, Amir N., Hanlon, Cathleen A., and Rupprecht, Charles E.

Source: Journal of Wildlife Diseases, 25(1) : 130-131

Published By: Wildlife Disease Association

URL: <https://doi.org/10.7589/0090-3558-25.1.130>

BioOne Complete (complete.BioOne.org) is a full-text database of 200 subscribed and open-access titles in the biological, ecological, and environmental sciences published by nonprofit societies, associations, museums, institutions, and presses.

Your use of this PDF, the BioOne Complete website, and all posted and associated content indicates your acceptance of BioOne's Terms of Use, available at www.bioone.org/terms-of-use.

Usage of BioOne Complete content is strictly limited to personal, educational, and non - commercial use. Commercial inquiries or rights and permissions requests should be directed to the individual publisher as copyright holder.

BioOne sees sustainable scholarly publishing as an inherently collaborative enterprise connecting authors, nonprofit publishers, academic institutions, research libraries, and research funders in the common goal of maximizing access to critical research.

***Dirofilaria immitis* in a Raccoon (*Procyon lotor*)**

Daniel E. Snyder,¹ Amir N. Hamir,² Cathleen A. Hanlon,³ and Charles E. Rupprecht,³ ¹ U.S. Department of Agriculture, Agricultural Research Service, Animal Parasite Research Laboratory, P.O. Box 952, Auburn, Alabama 36831, USA; ² University of Pennsylvania, School of Veterinary Medicine, New Bolton Center, 382 West Street Road, Kennett Square, Pennsylvania 19348, USA; ³ The Wistar Institute of Anatomy and Biology, 3601 Spruce Street, Philadelphia, Pennsylvania 19104, USA

ABSTRACT: A single juvenile male raccoon (*Procyon lotor*) was found naturally infected with *Dirofilaria immitis*. Two immature female worms were found in the heart of this raccoon at necropsy. Lesions attributable to the presence of these parasites were not found. Histopathologic examination of various tissues did not reveal any microfilariae. The raccoon may serve as an aberrant definitive host for this parasite, however, patent infections have not been reported.

Key words: Raccoon, *Procyon lotor*, *Dirofilaria immitis*, heartworm, natural infection, case report.

Sixteen raccoons were live-trapped on Parramore Island, Virginia (37°32'N, 75°38'W) as part of baseline data collection prior to oral rabies vaccine field trials. The animals were killed and complete necropsies were performed. Formalin fixed tissues and organs were sent to the Laboratory of Large Animal Pathology, School of Veterinary Medicine, New Bolton Center (University of Pennsylvania, Kennett Square, Pennsylvania 19348, USA) for histopathological examination. Examination of the heart of one juvenile male raccoon yielded two immature female filarid-like nematodes in the right atrium. These specimens were identified as *Dirofilaria immitis* based on the description given by Orihel (1961) for immature females of this parasite. These specimens have been deposited in the U.S. National Parasite Collection (Beltsville, Maryland 20705, USA; Accession Number 80409). Lesions were not associated with the presence of these parasites. Microfilariae were not seen on histopathologic examination of various tissues. Nematodes were not found in the hearts of the other 15 raccoons that were examined.

The dog heartworm (*D. immitis*) has become widespread across the United

States. Wild canids (Hubert et al., 1980; Simmons et al., 1980; Agostine and Jones, 1982), the domestic cat (Donahoe, 1975) and a number of domestic (Klein and Stoddard, 1977) and wildlife (Goble, 1942; Foil and Orihel, 1975; Williams and Dade, 1976; Crum et al., 1978) hosts have been reported as experimental or natural infections. The raccoon has rarely been reported to be naturally infected (Fox, 1941; Herman and Price, 1965). One experimental report indicated that the raccoon could not support the development of *D. immitis* (Christensen and Shelton, 1978). The domestic dog and a few species of wild canids are the only definitive hosts that play any significant role in the perpetuation of *D. immitis*. Apparently under certain natural conditions, the raccoon may serve as an aberrant definitive host for this filarid. The role of the raccoon in the transmission and epidemiology of *D. immitis* is insignificant as evidenced by (1) the extreme rarity of natural infections and (2) the absence of sexually mature nematodes in the few naturally infected raccoons in these previously recorded infections. Natural infections in raccoons simply serve to indicate that the geographic area is an endemic locus of *D. immitis* and that appropriate mosquito intermediate hosts are prevalent.

LITERATURE CITED

- AGOSTINE, J. C., AND G. S. JONES. 1982. Heartworms (*Dirofilaria immitis*) in coyotes (*Canis latrans*) in New England. *Journal of Wildlife Diseases* 18: 343-345.
- CHRISTENSEN, B. M., AND M. E. SHELTON. 1978. Laboratory observations on the insusceptibility of raccoons to *Dirofilaria immitis*. *Journal of Wildlife Diseases* 14: 22-23.
- CRUM, J. M., V. F. NETTLES, AND W. R. DAVIDSON. 1978. Studies on endoparasites of the black bear

- (*Ursus americanus*) in the southeastern United States. *Journal of Wildlife Diseases* 14: 178-186.
- DONAHOE, J. M. R. 1975. Experimental infection of cats with *Dirofilaria immitis*. *The Journal of Parasitology* 61: 599-605.
- FOIL, L., AND T. C. ORIHIEL. 1975. *Dirofilaria immitis* (Leidy, 1856) in the beaver, *Castor canadensis*. *The Journal of Parasitology* 61: 433.
- FOX, M. 1941. Heartworms in a coon. *Fort Dodge Biochemical Review* 12: 8.
- GOBLE, F. C. 1942. Dog heartworms in the muskrat in New York. *Journal of Mammalogy* 23: 346.
- HERMAN, C. M., AND D. L. PRICE. 1965. Epizootiologic studies on filarioids of the raccoon. *The Journal of Wildlife Management* 29: 694-699.
- HUBERT, G. F., JR., T. J. KICK, AND R. D. ANDREWS. 1980. *Dirofilaria immitis* in red foxes in Illinois. *Journal of Wildlife Diseases* 16: 229-232.
- KLEIN, J. B., AND E. D. STODDARD. 1977. *Dirofilaria immitis* recovered from a horse. *Journal of the American Veterinary Medical Association* 171: 354-355.
- ORIHIEL, T. C. 1961. Morphology of the larval stages of *Dirofilaria immitis* in the dog. *The Journal of Parasitology* 47: 251-262.
- SIMMONS, J. M., W. S. NICHOLSON, E. P. HILL, AND D. B. BRIGGS. 1980. Occurrence of (*Dirofilaria immitis*) in gray fox (*Urocyon cinereoargenteus*) in Alabama and Georgia. *Journal of Wildlife Diseases* 16: 225-228.
- WILLIAMS, J. F., AND A. W. DADE. 1976. *Dirofilaria immitis* in a wolverine. *The Journal of Parasitology* 62: 174-175.

Received for publication 27 June 1988.