

## **EVALUATION OF A MULTIVALENT PASTEURELLA HAEMOLYTICA VACCINE IN BIGHORN SHEEP: PROTECTION FROM EXPERIMENTAL CHALLENGE**

Authors: Kraabel, Brita J., Miller, Michael W., Conlon, Jennifer A., and McNeil, Heather J.

Source: Journal of Wildlife Diseases, 34(2) : 325-333

Published By: Wildlife Disease Association

URL: <https://doi.org/10.7589/0090-3558-34.2.325>

---

BioOne Complete ([complete.BioOne.org](https://complete.BioOne.org)) is a full-text database of 200 subscribed and open-access titles in the biological, ecological, and environmental sciences published by nonprofit societies, associations, museums, institutions, and presses.

Your use of this PDF, the BioOne Complete website, and all posted and associated content indicates your acceptance of BioOne's Terms of Use, available at [www.bioone.org/terms-of-use](https://www.bioone.org/terms-of-use).

Usage of BioOne Complete content is strictly limited to personal, educational, and non - commercial use. Commercial inquiries or rights and permissions requests should be directed to the individual publisher as copyright holder.

---

BioOne sees sustainable scholarly publishing as an inherently collaborative enterprise connecting authors, nonprofit publishers, academic institutions, research libraries, and research funders in the common goal of maximizing access to critical research.

## EVALUATION OF A MULTIVALENT *PASTEURELLA HAEMOLYTICA* VACCINE IN BIGHORN SHEEP: PROTECTION FROM EXPERIMENTAL CHALLENGE

Brita J. Kraabel,<sup>1</sup> Michael W. Miller,<sup>1,6</sup> Jennifer A. Conlon,<sup>2,4</sup> and Heather J. McNeil<sup>3,5</sup>

<sup>1</sup> Colorado Division of Wildlife, Wildlife Research Center, 317 West Prospect Road, Fort Collins, Colorado 80526-2097, USA

<sup>2</sup> Fort Dodge Laboratories, 800 Fifth Street Northwest, Fort Dodge, Iowa 50501, USA

<sup>3</sup> Ayerst Veterinary Laboratories, 131 Malcolm Road, Guelph, Ontario N1K 1A8, Canada

<sup>4</sup> Present address: Rhone Merieux, Incorporated, 115 Transtech Drive, Athens, Georgia 30601, USA

<sup>5</sup> Present address: Department of Pathobiology, Ontario Veterinary College, University of Guelph, Guelph, Ontario N1G 2W1, Canada

<sup>6</sup> Corresponding author (e-mail: mike.miller@state.co.us)

**ABSTRACT:** The efficacy of a multivalent *Pasteurella haemolytica* vaccine (A1, A2, T10) in reducing morbidity and mortality associated with pneumonic pasteurellosis in bighorn sheep (*Ovis canadensis*) was examined. Fifteen captive bighorns were divided equally into three treatment groups based on vaccination status: control (no vaccination), one dose 10 days prior to challenge, or one or two doses 57 wk prior to challenge. At challenge, each bighorn received about  $6.2 \times 10^7$  colony forming units of *P. haemolytica* (biotype T, serotype 10, biogroup 4, CDCs, ribotype ECG; "Alamosa Canyon" strain) suspension sprayed into the proximal trachea. Vaccination reduced ( $P = 0.1$ ) mortality in bighorns vaccinated 10 days prior to challenge as compared to controls. Although mortality rates in bighorns vaccinated 57 wk prior to challenge did not differ from controls ( $P = 0.26$ ), a trend in reduced mortality was apparent. Ranked cumulative postmortem scores based on observed gross lesions and bacteriology did not differ ( $P \geq 0.14$ ) between vaccinated animals and control animals. Neutralizing antibody titers to *P. haemolytica* leukotoxin were elevated ( $P = 0.003$ ) at challenge in bighorns vaccinated 10 days before challenge, and neutralizing titers in bighorns from both vaccinated groups were elevated at death  $\leq 7$  days after challenge ( $P \leq 0.004$ ). In contrast, agglutinating antibody titers to *P. haemolytica* serotype A1 or T10 surface antigens did not differ between vaccinated and unvaccinated bighorns ( $P \geq 0.19$ ). Based on these data, we believe that this experimental *P. haemolytica* vaccine is safe and can stimulate protective immunity from pneumonic pasteurellosis in bighorn sheep. Further evaluation of this vaccine as a tool in preventing and managing pasteurellosis in wild bighorn sheep appears warranted.

**Key words:** Bighorn sheep, experimental study, *Ovis canadensis*, *Pasteurella haemolytica*, pasteurellosis, pneumonia, serology, vaccine.

### INTRODUCTION

Epidemics of respiratory disease have probably plagued bighorn sheep (*Ovis canadensis*) populations for at least a century (Buechner, 1960; Post, 1962). *Pasteurella* spp. are perhaps the most common pathogens isolated from bighorns during these epidemics (Potts, 1937; Marsh, 1938; Post, 1962; Onderka and Wishart, 1984; Festa-Bianchet, 1988; Miller et al., 1991). The inability to prevent or control mortality caused by pneumonic pasteurellosis represents a significant obstacle to successful long-term bighorn management (Onderka and Wishart, 1984; Festa-Bianchet, 1988; Hobbs and Miller, 1992).

Despite intensive study of the bighorn

pneumonia complex, few tools for its management have emerged. Although vaccination has been suggested as a potential management tool for nearly four decades, previous attempts to vaccinate bighorn sheep against pasteurellosis have yielded largely discouraging results. Early studies of autogenous bacterins showed some promise of protection (Rufi, 1961; Post, 1962; Nash, 1972), but these vaccines either failed in application (Howe, 1964) or were never fully evaluated or incorporated into bighorn management programs. Other attempts to prevent pneumonia in bighorns using an autogenous bacterin or pre-challenge inoculation with cytotoxic *P. haemolytica* also failed (Foreyt, 1992; Foreyt

and Silflow, 1996), and use of a modified-live *P. haemolytica* A1 vaccine apparently caused pasteurellosis in healthy bighorns (Onderka et al., 1988).

More recently, Miller et al. (1997) described a multivalent *P. haemolytica* vaccine (A1, A2, T10) that stimulated marked elevations in antibody titers to leukotoxin and surface antigens. The magnitude of antibody responses in vaccinated bighorns equaled or exceeded those associated with protection from experimental challenge in domestic sheep and cattle treated with similar vaccines (Miller et al., 1997). Here, we examined the efficacy of this multivalent vaccine in reducing morbidity and mortality in captive bighorn sheep challenged with a pathogenic strain of *P. haemolytica*.

#### MATERIALS AND METHODS

We used 17 captive Rocky Mountain bighorn sheep (*O. canadensis canadensis*) and four captive bighorn × mouflon f2 hybrid sheep (*O. canadensis canadensis* × *O. musimon*) in this experiment. All sheep were housed at the Colorado Division of Wildlife's (CDOW) Foothills Wildlife Research Facility (Fort Collins, Colorado, USA; 40°35'N, 105°10'W) throughout the study. Grass/alfalfa hay mix and a pelleted high-energy supplement were provided as prescribed under established feeding protocols for bighorn sheep in respective age/sex classes (Miller, 1990); fresh water and mineralized salt blocks were provided *ad libitum*. The resident bighorn herd was lungworm-free and had no previous evidence of infection with respiratory viruses (M. W. Miller, unpubl. data).

In evaluating vaccine efficacy, a repeatable representative model of natural pneumonic pasteurellosis in bighorn sheep was essential. Consequently, we conducted a pilot study to establish the effective minimum dose of *P. haemolytica* and optimal administration route needed to cause acute pneumonic pasteurellosis in bighorns. We used all four bighorn × mouflon hybrids and two bighorns in this pilot study; hybrid sheep were obtained from the Wyoming Game and Fish Department's Sybille Wildlife Research and Conservation Unit (SWRCU) (Wheatland, Wyoming, USA) in December 1995. Two or three individuals resided together in 100m<sup>2</sup> isolation pens throughout this study.

We used a field strain of *P. haemolytica* biotype T, serotype 10, biogroup 4.C:DS, ribotype

ECO ("Alamosa Canyon" strain) for challenge. This isolate was originally recovered from lung tissue collected in 1990 during a pasteurellosis epidemic in a free-ranging bighorn population in southcentral Colorado (M. W. Miller, unpubl. data) and had been passed <5 times on artificial media; previous *in vitro* evaluations of this isolate demonstrated marked leukotoxin production (Kraabel and Miller, 1997). Challenge doses were prepared by quick-thawing frozen *P. haemolytica* aliquots, placing them on 5% sheep blood agar plates (BBL Microbiology Systems, Becton Dickinson and Company, Cockeysville, Maryland, USA), and incubating them for 16 to 20 hr at 37 C. Isolated colonies were used to inoculate brain heart infusion broth (BHIB) (Difco Laboratories, Detroit, Michigan, USA); after 6 hr, this broth was diluted with fresh BHIB to an optical density at 525 nm (OD<sub>525</sub>) of 0.1 and incubated until the OD<sub>525</sub> reached 0.29 to 0.3. The culture was centrifuged at 4,000 × G for 15 min and resuspended in sterile phosphate buffered saline (PBS) solution at estimated concentrations of 10<sup>4</sup>, 10<sup>6</sup>, 10<sup>8</sup> or 10<sup>10</sup> colony forming units per ml (CFUs/ml). An aliquot of each resulting challenge dose level was plated, and doses were confirmed by counting colonies after 16 hr of incubation at 37 C.

We administered bacteria by sedating the sheep with 5 to 20 mg xylazine hydrochloride (HCL) (Wildlife Pharmaceuticals, Fort Collins, Colorado, USA) intravenously (IV) or intramuscularly (IM), opening their mouths, and visualizing the arytenoids with the aid of a laryngoscope. We initially administered the challenge dose by spraying culture solution onto the glottis (OP); subsequently, we delivered the bacteria into the proximal trachea (IT) (Table 1). Sedation was reversed with 30 to 50 mg yohimbine HCl (Wildlife Pharmaceuticals) IV immediately after challenge. One hybrid (MB4) was immobilized with a combination of 1.5 mg carfentanil HCl (Wildlife Pharmaceuticals), 90 mg ketamine HCl (Wildlife Pharmaceuticals), and 20 mg xylazine HCl (Wildlife Pharmaceuticals) IM for his second challenge; in this case, immobilization was reversed with 40 mg yohimbine HCl IV and 150 mg naltrexone HCl (Wildlife Pharmaceuticals), divided ¼ IV and ¾ subcutaneously.

Postchallenge, sheep were observed ≥3 times daily for signs of respiratory disease. In cases where the first selected dose/route combination failed to produce appreciable clinical signs or mortality in pilot animals ≤4 days postchallenge, those individuals were challenged again with a different dose/route combination (Table 1). Sheep that survived ≥4 days after a second challenge were sedated with 100

TABLE 1. Chronological listing of challenge doses in colony forming units (CFUs) per ml of *Pasteurella haemolytica* and routes used in a pilot study determining optimal techniques for vaccine challenge in captive bighorn sheep.

Dose (CFUs)/route <sup>a</sup>	Animal <sup>b</sup>	Outcome (duration) <sup>c</sup> /postmortem score <sup>d</sup>
10 <sup>4</sup> /OP	MB-2	alive/rc
10 <sup>6</sup> /OP	MB-3	alive/rc
10 <sup>8</sup> /OP	MB-1	alive/rc
10 <sup>10</sup> /OP	MB-2	died (<39 hr)/71
10 <sup>8</sup> /IT	MB-3	died (<72 hr)/65
10 <sup>10</sup> /IT	MB-1	died (<15 hr)/69
10 <sup>6</sup> /IT	MB-4	alive/rc
10 <sup>7</sup> /IT	MB-4	alive/19
10 <sup>7</sup> /IT	B-L395	died (<60 hr)/65
10 <sup>6</sup> /IT	B-Q95	alive/2

<sup>a</sup> OP = oropharyngeal; IT = intratracheal.

<sup>b</sup> MB = mouflon × bighorn hybrid sheep; B = bighorn sheep.

<sup>c</sup> Alive = survived >96 hr postchallenge; died = died (hr) postchallenge; rc = rechallenged.

<sup>d</sup> Maximum postmortem score = 72; see text for scoring methods.

to 300 mg xylazine HCl IV or IM and euthanized by IV injection of 100 mg/kg pentobarbital sodium solution (Beuthanasia®-D, Schering-Plough Animal Health Corporation, Kenilworth, New Jersey, USA).

All sheep were necropsied immediately after death. Gross lesions were photographed and scored using a system adapted from Jones et al. (1989). Each animal received a cumulative postmortem score (maximum = 72) based on severity of gross lung lesions, gross evidence of other organ involvement, and *P. haemolytica* isolation from select tissues. At necropsy, we diagramed areas of lung involvement and calculated a lung lesion score by estimating the proportion of each lung lobe affected from both dorsal and ventral aspects. Each lobe's contribution to the total score was extrapolated from estimated percentages of total lung volume occupied by each lobe in a normal domestic sheep's lung. The mean ventral and dorsal pneumonic scores were divided by 2 to give a final lung lesion score (maximum = 50). In addition, the amount of fibrin present on the visceral or parietal pleura was scored as 0 (none), 1 (small amount), 3 (moderate amount), or 6 (substantial amount) (maximum = 6). We scored other organ involvement by adding 1 point each for subcutaneous, epicardial, renal capsular, splenic, hepatic, or parietal pleural hemorrhages (maximum = 6). Representative samples of lung (at lesion margins, where available), liver, spleen, kidney, and right

carpus were refrigerated and shipped overnight on ice packs to the Caine Veterinary Teaching and Research Center (CVTRC) (Caldwell, Idaho, USA) for culture and biotyping using methods described by Miller et al. (1997); two points were added for each site yielding *P. haemolytica* (maximum = 10).

We used 15 captive Rocky Mountain bighorn sheep in the vaccine challenge trial. Three or four individuals resided in 100m<sup>2</sup> isolation pens throughout the study. Pen assignments were made to accommodate social differences and thereby minimize stress on cohoused individuals, but treatment assignments were randomly distributed across pens.

Bighorns were subdivided into three groups of five based on vaccination status: 0 (control, no vaccination), 1 (vaccinated 10 days prior to challenge), or 2 (vaccinated 57 wks prior to challenge). Bighorns in groups 0 and 1 had not been vaccinated previously, and were assigned randomly to respective groups; bighorns in group 2 had received experimental vaccine in February 1995 during a previous study (Miller et al., 1997). The experimental *P. haemolytica* vaccine (Langford Laboratories, Incorporated, Guelph, Ontario, Canada; lot 940902) evaluated here was the same vaccine previously described by Miller et al. (1997).

After an acclimation period of 12 days in isolation pens, we aseptically injected 2 ml of vaccine IM into bighorns in treatment group 1; bighorns in groups 0 and 2 received 2 ml 0.9% saline, aseptically injected IM. Ten days after vaccination or saline injection, each bighorn received about  $6.3 \times 10^7$  CFUs of *P. haemolytica* (biotype T, serotype 10, biogroup 4, CDS, ribotype E<sub>CO</sub>). The challenge dose was prepared as described previously to a final concentration  $1.25 \times 10^8$ /ml; sheep were inoculated with 500 µl of culture suspension diluted in 5 ml of PBS. To administer the challenge, we sedated each bighorn with 5 to 20 mg IV xylazine HCl, held open its mouth with a speculum, anesthetized the larynx with a 2% solution of lidocaine HCl (Sparhawk Laboratories, Kansas City, Kansas, USA), placed a spray nozzle into the glottis with aid of a laryngoscope, and sprayed diluted culture suspension in a fine mist into the proximal trachea. Sedation was reversed with 10 to 40 mg IV yohimbine HCl immediately after administration of the challenge dose.

This experiment was designed to evaluate the efficacy of vaccination in preventing or reducing pneumonic pasteurellosis. Although we anticipated some sheep would become clinically ill or die in response to challenge, our approach attempted to balance the need to evaluate clinical responses of challenged bighorns with the need to relieve pain and suffering in

affected animals. The health of bighorns was closely monitored by attending veterinarians  $\geq 3$  times daily (early morning, mid-day, and dusk) after challenge. Analgesic or anti-inflammatory drugs were not provided to affected animals because they might have interfered with vaccine-mediated responses to challenge. Moreover, because the goal of this research was to determine whether the vaccine protected bighorns in the face of infection with *P. haemolytica*, we did not euthanize animals demonstrating respiratory signs unless they became recumbant, extremely depressed, or moribund. All animals that died or were euthanized during the experiment were necropsied to determine cause of death; surviving bighorns were euthanized with 100 mg/kg pentobarbital sodium solution (Beuthanasia® -D) 7 days after challenge and necropsied. All necropsies and sampling were performed immediately after death and carcasses scored as described above. Representative tissue samples from all major organ systems also were collected and placed in 10% neutral buffered formalin and submitted for histological evaluation (Wyoming State Veterinary Laboratory, Laramie, Wyoming, USA).

In addition to postmortem samples, we collected 10 to 12 ml of blood via jugular venipuncture for antibody measurements from each bighorn immediately prior to vaccination, challenge, and death; we also collected oropharyngeal and nasal swabs from each bighorn in conjunction with sample collections for serology. All blood samples were held 1 to 4 hr at 22 C, centrifuged, and serum collected. Serum was stored at -20 C until analyzed. Swabs were placed in transport tubes containing modified Cary and Blair medium (Port-A-Cul®, Becton Dickinson and Company) and shipped overnight on ice packs to CVTRC for culture and analysis. Measurement of leukotoxin neutralizing antibodies and agglutinating antibodies to serotype-specific surface antigens, as well as culture and biogrouping of *P. haemolytica* from nasal and oropharyngeal swabs and tissue samples, followed methods described by Miller et al. (1997); titers were expressed as  $\log_2$ .

In evaluating vaccine efficacy, we compared (1) mortality rates, (2) postmortem scores, (3) serum neutralizing antibody titers to *P. haemolytica* leukotoxin, and (4) serum antibody titers to *P. haemolytica* surface antigen; in all cases, we compared each vaccine group to controls but did not compare between vaccine groups. Mortality rates were compared using Fisher's exact test (Mielke and Berry, 1992). Postmortem scores among groups were compared using Wilcoxon rank sum test with ties (SAS Institute, Inc., 1989). We analyzed serologic data using least squares analysis of vari-

ance (ANOVA) for general linear models (SAS Institute, Inc., 1989). Antibody responses were analyzed with ANOVA for a randomized complete block design with a repeated measures structure, with vaccine treatment group (0, 1, 2) as the sole main effect and time treated as a within subject effect (Morrison, 1976); we used least squares means for pairwise comparisons of antibody titers between each vaccine group and controls at vaccination, challenge, and postmortem. We used  $\alpha = 0.1$  for all statistical comparisons. Our challenge study was designed with sample sizes sufficient to offer power of  $\geq 0.63$  in detecting  $\geq 4$ -fold differences in mortality between treatment and control groups.

## RESULTS

Spraying about  $10^4$ ,  $10^6$ , or  $10^8$  CFUs of *P. haemolytica* into the oropharyngeal region of bighorn  $\times$  mouflon hybrid sheep failed to produce acute pneumonic pasteurellosis, and none of the sheep died (Table 1); however, about  $10^{10}$  CFUs sprayed OP caused death in one hybrid <39 hr later. Delivered IT, about  $10^6$  CFUs failed to cause clinical signs or significant lesions, but  $10^8$  or  $10^{10}$  CFUs IT caused fatal pneumonia and/or systemic disease within 72 hr (Table 1) in hybrids. Dosed at about  $10^7$  CFUs IT, the challenge strain caused severe pneumonia in a hybrid and fatal pneumonia and systemic disease in a bighorn (Table 1).

Vaccinating bighorns 10 days prior to challenge reduced ( $P = 0.1$ ) mortality (1/5) as compared to those for controls (4/5); although mortality (2/5) in bighorns vaccinated 57 wk prior to challenge did not differ from controls ( $P = 0.26$ ), a trend in reduced mortality was apparent (Fig. 1). Clinical signs were apparent in some animals <18 hr after challenge; eleven of the 15 bighorns, including all five controls, showed at least moderate signs of respiratory disease (depression, shallow or labored respiration, anorexia)  $\leq 72$  hr after challenge. Four of five controls died about 30 to 72 hr postchallenge; three vaccinated bighorns succumbed about 84 to 120 hr postchallenge. At necropsy, we observed evidence of acute pneumonia in all 15 big-




FIGURE 1. Vaccination reduced bighorns' susceptibility to fatal pasteurellosis resulting from experimental challenge. Mortality was lower ( $P = 0.1$ ) in bighorns vaccinated 10 days before challenge; mortality in bighorns vaccinated 57 wks prior to challenge did not differ from controls ( $P = 0.26$ ), but a reduced trend was apparent. Similarly, lesions tended to be less severe in vaccinated animals, although ranked postmortem scores did not differ between controls and 10 day vaccinates ( $P = 0.14$ ) or 57 wk vaccinates ( $P = 0.22$ ).

horns. Lung damage and other lesions tended to be less severe in vaccinated animals (Fig. 1), but ranked postmortem scores did not differ between controls and 10 day vaccinates ( $P = 0.14$ ) or 57 wk vaccinates ( $P = 0.22$ ).

Leukotoxin neutralizing antibody titers were elevated ( $P = 0.003$ ) at challenge in bighorns vaccinated 10 days previously as compared to titers in controls, and in both 10 day and 57 wk vaccinates at postmortem  $\leq 7$  days after challenge ( $P \leq 0.004$ ) (Fig. 2A). In contrast, titers of agglutinating antibody to *P. haemolytica* serotype A1 or T10 surface antigens did not differ between vaccinated and unvaccinated bighorns before, at, or after challenge ( $P \geq 0.19$ ) (Fig. 2B–C). All seven bighorns that succumbed had postchallenge leukotoxin neutralizing antibody titers  $\leq 4.0$ ; seven of eight challenge survivors had neutralizing titers  $\geq 3.5$  (Fig. 3).

We recovered *P. haemolytica* from all captive bighorns prior to challenge; in all, fourteen distinguishable biogroups were identified among 54 isolates. The challenge strain of *P. haemolytica* (biogroup 4-CDS) was not recovered from oropharyngeal or nasal swabs from any of the study bighorns at vaccination or at chal-




FIGURE 2. Vaccination stimulated both leukotoxin neutralizing and agglutinating antibody responses in bighorns. Leukotoxin neutralizing antibody titers (A) were elevated ( $P = 0.003$ ) at challenge in bighorns vaccinated 10 days previously, and in both 10 day and 57 wk vaccinates at postmortem  $\leq 7$  days after challenge ( $P \leq 0.004$ ). In contrast, titers of agglutinating antibody to *P. haemolytica* serotype A1 (B) or T10 (C) surface antigens did not differ between vaccinated and unvaccinated bighorns ( $P \geq 0.19$ ). Bars are mean observations; vertical lines are  $+1$  standard error of mean observations.



did not prevent bighorns from being colonized with the challenge strain.

Although the precise mechanisms for vaccine protection were not studied here, bighorn survival appeared strongly correlated to the magnitude of leukotoxin neutralizing antibody titers measured post-challenge (Fig. 3). Agglutinating antibody titers to *P. haemolytica* surface antigens alone were clearly insufficient to protect bighorns from experimental challenge. Titers to serotype T10 surface antigens were relatively high throughout the study across all treatment groups (Fig. 2C), probably reflecting previous oropharyngeal exposure to nonpathogenic *P. haemolytica* strains endemic in this captive population (Miller et al., 1997). Similarly, modest pre-challenge serotype A1 agglutinating titers rose in all three groups after challenge (Fig. 2B), but those elevated titers failed to protect controls from pasteurellosis. In contrast, vaccination stimulated marked elevation of leukotoxin neutralizing antibody titers in previously unvaccinated bighorns (Fig. 2A). Perhaps more importantly, challenge apparently stimulated anamnestic responses in previously vaccinated bighorns (Fig. 2A). Postchallenge leukotoxin neutralizing titers in seven of eight surviving bighorns approximated titers in free-ranging bighorns that survived naturally-occurring epidemics (M. W. Miller, unpubl. data), and also approximated titers in domestic sheep and cattle protected from experimental challenges (A. Alexander, pers. commun.; Conlon et al., 1995). Of the parameters we measured, leukotoxin neutralizing antibody titers seemed most useful in predicting the outcome of experimental challenge with pathogenic *P. haemolytica*. These observations are consistent with studies of immunity to pasteurellosis in domestic ruminants (Donachie, 1995; Shewen, 1995).

Responses to vaccination observed in this and a previous study (Miller et al., 1997) suggest several potential applications in managing pasteurellosis in wild bighorns. Because antibody titers in vac-

inated bighorns rise rapidly and remain elevated for 12 to 16 wk (Miller et al., 1997), there may be benefits to vaccinating wild sheep early in the course of a pneumonia epidemic. Perhaps of equal importance, the trend toward reduced mortality in bighorns vaccinated >1 yr prior to challenge demonstrates the potential utility of annual vaccination to prevent epidemics in some management situations. Stimulation of dramatic antibody responses by a single vaccination greatly enhances prospects for its use in wild bighorns.

Although bighorn pasteurellosis displays several epidemiological features that might favor vaccination as an effective control strategy (Barlow, 1996), the efficacy of vaccination in long-term management of bighorn populations remains untested. Because pasteurellosis epidemics in free-ranging populations are sometimes complicated by infections with respiratory viruses, chlamydia, or parasitic lungworms, vaccination could prove less effective under field conditions. Variable pathogenicity among *P. haemolytica* strains (Miller et al., 1991; Miller et al., 1995; Kraabel and Miller, 1997) could also affect vaccine efficacy. Moreover, because pasteurellosis among lambs born into herds with immune carriers can severely impair recruitment after an epidemic, we recognize that improving individual bighorn survival during an epidemic may not necessarily lead to improved population performance or survival thereafter. However, our data suggest that the multivalent *P. haemolytica* vaccine used here is safe and stimulates protective immunity in bighorn sheep. Consequently, we believe field evaluation of this vaccine through carefully designed management experiments in free-ranging bighorn sheep is warranted.

#### ACKNOWLEDGMENTS

Our study was supported by the CDOW, Federal Aid in Wildlife Restoration Project W-153-R-8, and Fort Dodge Laboratories. We thank the Wyoming Game and Fish Department for providing hybrids and some bighorns used in this study and CVTRC for providing



bacteriology services; we specifically recognize J. Bulgin, A. Case, A. Lotto, E. Zimmerman, G. Stout, A. Alexander, S. Chu, and R. Pacer for assistance in various facets of this study. Guidance on experimental design and data analyses was provided by D. Bowden and G. Gallo; the CDOW's Animal Care and Use Committee provided careful review of and useful suggestions on conduct of our study. D. L. Hunter and D. K. Onderka provided helpful reviews of an earlier version of this manuscript.

#### LITERATURE CITED

- ALEXANDER, A., M. ALLEY, AND J. CONLON. 1995. New vaccines for the control of ovine pneumonia. In Proceedings of 25th Sheep and Beef Cattle Seminar, G. Budge, (ed.). Publication Number 165, Veterinary Continuing Education, Massey University, Palmerston North, New Zealand, pp. 137-141.
- BARLOW, N. D. 1996. The ecology of wildlife disease control: Simple models revisited. *Journal of Applied Ecology* 33: 303-314.
- BUECHNER, H. K. 1960. The bighorn sheep in the United States, its past, present, and future. *Wildlife Monographs* 4: 174.
- CONLON, J. A. R., G. F. GALLO, P. E. SHEWEN, AND C. ADLAM. 1995. Comparison of protection of experimentally challenged cattle vaccinated once or twice with a *Pasteurella haemolytica* bacterial extract vaccine. *Canadian Journal of Veterinary Research* 59: 179-182.
- DONACHIE, W. 1995. Vaccine development against *Pasteurella haemolytica* infection in sheep. In *Haemophilus, Actinobacillus, and Pasteurella*, W. Donachie, F. A. Lainson, and J. C. Hodgson (eds.). Plenum Press, New York, New York, pp. 25-37.
- FESTA-BIANCHET, M. 1988. A pneumonia epizootic in bighorn sheep, with comments on preventive management. *Biennial Symposium of the Northern Wild Sheep and Goat Council* 6: 66-76.
- FOREYT, W. J. 1992. Failure of an experimental *Pasteurella haemolytica* vaccine to prevent respiratory disease and death in bighorn sheep after exposure to domestic sheep. *Biennial Symposium of the Northern Wild Sheep and Goat Council* 8: 155-163.
- , AND R. M. SILFLOW. 1996. Attempted protection of bighorn sheep (*Ovis canadensis*) from pneumonia using a nonlethal cytotoxic strain of *Pasteurella haemolytica*, biotype A, serotype 11. *Journal of Wildlife Diseases* 32: 315-321.
- , K. P. SNIPES, AND R. W. KASTEN. 1994. Fatal pneumonia following inoculation of healthy bighorn sheep with *Pasteurella haemolytica* from healthy domestic sheep. *Journal of Wildlife Diseases* 30: 137-145.
- HOBBS, N. T., AND M. W. MILLER. 1992. Interactions between pathogens and hosts: Simulation of pasteurellosis epizootics in bighorn sheep populations. In *Wildlife 2001: Populations*, D. R. McCullough and R. H. Barrett, (eds.). Elsevier Science Publishers, Ltd., London, England, pp. 997-1007.
- HOWE, D. L. 1964. Etiology of pneumonia in bighorn sheep. In *Federal Aid in Wildlife Restoration, Game and Fish Laboratory Research, Research Project Segment, Job Completion Report, Project Number FW-3-R-11, Work Plan 1, Job 2W*. Wyoming Game and Fish Department, Cheyenne, Wyoming, pp. 16-18.
- JONES, G. E., W. DONACHIE, A. D. SUTHERLAND, D. P. KNOX, AND J. S. GILMOUR. 1989. Protection of lambs against experimental pneumonic pasteurellosis by transfer of immune serum. *Veterinary Microbiology* 20: 59-71.
- KRAABEL, B. J., AND M. W. MILLER. 1997. Effect of simulated stress on susceptibility of bighorn sheep neutrophils to *Pasteurella haemolytica* leukotoxin. *Journal of Wildlife Diseases* 33: 558-566.
- MARSH, H. 1938. Pneumonia in Rocky Mountain bighorn sheep. *Journal of Mammalogy* 19: 214-219.
- MIELKE, P. W., AND K. J. BERRY. 1992. Fisher's exact probability test for cross-classification tables. *Educational and Psychological Measurement* 52: 97-101.
- MILLER, M. W. 1990. Animal and pen support facilities for mammals research. In *Wildlife Research Report, Mammals Research, Federal Aid Projects, Job Progress Report, Project W-153-R-3, Work Plan 1a, Job 1*. Colorado Division of Wildlife, Fort Collins, Colorado, pp. 45-63.
- , N. T. HOBBS, AND E. S. WILLIAMS. 1991. Spontaneous pasteurellosis in captive Rocky Mountain bighorn sheep (*Ovis canadensis canadensis*): Clinical, laboratory, and epizootiological observations. *Journal of Wildlife Diseases* 27: 534-542.
- , J. A. CONLON, H. J. MCNEIL, J. M. BULGIN, AND A. C. S. WARD. 1997. Evaluation of a multivalent *Pasteurella haemolytica* vaccine in bighorn sheep: Safety and serologic responses. *Journal of Wildlife Diseases* 33: 738-748.
- , B. J. KRAABEL, J. A. CONLON, H. J. MCNEIL, AND J. M. BULGIN. 1995. Strategies for managing infectious diseases in mountain sheep populations. *Wildlife Research Report, Mammals Research, Federal Aid Projects, Job Progress Report, Project W-153-R-8, WP2a, J4*. Colorado Division of Wildlife, Fort Collins, Colorado, pp. 151-161.
- MORRISON, D. F. 1976. *Multivariate statistical methods*, 2nd Edition. McGraw-Hill Book Co., New York, New York, 415 pp.
- NASH, P. 1972. Development of a *Pasteurella* bacterin for bighorn sheep (*Ovis canadensis cana-*

- densis*). Ph.D. Thesis, Colorado State University, Fort Collins, Colorado, 102 pp.
- ONDERKA, D. K., AND W. D. WISHART. 1984. A major bighorn sheep dieoff from pneumonia in southern Alberta. *Biennial Symposium of the Northern Wild Sheep and Goat Council* 4: 356–363.
- , S. A. RAWLUK, AND W. D. WISHART. 1988. Susceptibility of Rocky Mountain bighorn sheep and domestic sheep to pneumonia induced by bighorn and domestic livestock strains of *Pasteurella haemolytica*. *Canadian Journal of Veterinary Research* 52: 439–444.
- POST, G. 1962. Pasteurellosis of Rocky Mountain bighorn sheep (*Ovis canadensis*). *Wildlife Disease* 23: 1–13.
- POTTS, M. K. 1937. Hemorrhagic septicemia in the bighorn of Rocky Mountain National Park. *Journal of Mammalogy* 18: 105–106.
- RUFI, V. G. 1961. Study of pasteurellosis in bighorn sheep. In *Federal Aid in Wildlife Restoration, Game & Fish Laboratory Research, Investigations Projects, Job Completion Report, Project Number FW-3-R-8, Work Plan 1, Job 2W*. Wyoming Game and Fish Department, Cheyenne, Wyoming, pp. 21–27.
- SAS INSTITUTE, INC. 1989. SAS/STAT® User's Guide, Version 6, 4th Edition, Volume 2. SAS Institute, Cary, North Carolina, 846 pp.
- SHEWEN, P. E. 1995. Host response to infection with HAP: Implications for vaccine development. In *Haemophilus, Actinobacillus, and Pasteurella*, W. Donachie, F. A. Lainson, and J. C. Hodgson (eds.). Plenum Press, New York, New York, pp. 165–171.
- SILFLOW, R. M., W. J. FOREYT, S. M. TAYLOR, W. W. LAEGRIED, H. D. LIGGITT, AND R. W. LEID. 1989. Comparison of pulmonary defense mechanisms in Rocky Mountain bighorn sheep (*Ovis canadensis canadensis*) and domestic sheep. *Journal of Wildlife Diseases* 25: 514–520.
- SPRAKER, T. R., C. P. HIBLER, G. G. SCHOONVELD, AND W. S. ADNEY. 1984. Pathologic changes and microorganisms found in bighorn sheep during a stress-related die-off. *Journal of Wildlife Diseases* 20: 319–327.

*Received for publication 24 April 1997.*