



Appendix 6

Source: A Rapid Biodiversity Assessment of the Nakauvadra Range, Ra Province, Fiji: 125

Published By: Conservation International

URL: <https://doi.org/10.1896/054.057.0121>

BioOne Complete (complete.BioOne.org) is a full-text database of 200 subscribed and open-access titles in the biological, ecological, and environmental sciences published by nonprofit societies, associations, museums, institutions, and presses.

Your use of this PDF, the BioOne Complete website, and all posted and associated content indicates your acceptance of BioOne's Terms of Use, available at www.bioone.org/terms-of-use.

Usage of BioOne Complete content is strictly limited to personal, educational, and non - commercial use. Commercial inquiries or rights and permissions requests should be directed to the individual publisher as copyright holder.

BioOne sees sustainable scholarly publishing as an inherently collaborative enterprise connecting authors, nonprofit publishers, academic institutions, research libraries, and research funders in the common goal of maximizing access to critical research.

Appendix 6
Sites of interest for terrestrial insects in the Nakauvadra Range, Ra Province, Fiji
Hilda Waqa-Sakiti

| No. | Name | Latitude (S) | Longitude (E) | Elevation (m) | Comments |
|---------|--|--------------|---------------|---------------|---|
| ento 1 | <i>Nesobasis angolicolis</i> <i>Melanesobasis</i> sp. | 17°26'15"S | 178°05'36"E | 406 | dry creek, 2° forest edge |
| ento 2 | <i>Papilio schemltzi</i> | 17°26'19"S | 178°05'42"E | 476 | 1st sighting of <i>P. schemltzi</i> , host plant <i>Micromellum minutum</i> found within study area |
| ento 3 | Leaf litter 1 | 17°26'27"S | 178°05'45"E | 541 | mid-altitude, on gentle slope |
| ento 4 | Base camp 1 <i>Phasmotaenia inermis</i> (syn. <i>Hermarchus</i>) <i>Liocheles australasiae</i> (scorpion) | 17°27'22"S | 178°05'54"E | 329 | near confluence of Volivoli and Nabiya creek |
| ento 5 | Light trap 1 | 17°27'20"S | 178°05'52"E | 327 | adjacent to Nabiya Creek |
| ento 6 | Leaf litter 2 | 17°27'29"S | 178°05'35"E | 503 | mid-altitude, on very steep slope |
| ento 7 | Leaf litter 3 | 17°27'25"S | 178°05'55"E | 328 | low altitude leaf litter sampling |
| ento 8 | Light trap 2 | 17°27'21"S | 178°05'56"E | 308 | |
| ento 9 | Tree beating 1 | 17°27'20"S | 178°05'56"E | 327 | |
| ento 10 | Tree beating 2 | 17°27'21"S | 178°05'56"E | 309 | |
| ento 11 | <i>Nisyrius spinulosus</i> (synonym to <i>Cotylosoma</i>) | 17°27'29"S | 178°05'11"E | 424 | found within dakua (<i>Agathis macrophylla</i>) plot camouflaged by lichens on the tree bark |
| ento 12 | <i>P. schmeltzi</i> | 17°27'29"S | 178°05'57"E | 330 | numerous sightings of <i>P. scheltzi</i> and <i>Danaus plexippus</i> |
| ento 13 | Base camp 2 Buprestidae (jewel beetle) | 17°27'46"S | 178°07'46"E | 275 | A rarely encountered beetle family |
| ento 14 | Light trap 3 | 17°27'20"S | 178°05'52"E | 327 | |
| ento 15 | Light trap 4 | 17°27'47"S | 178°07'43"E | 249 | |
| ento 16 | Leaf litter 4 | 17°27'47"S | 178°07'45"E | 253 | |