



Map

Source: A Rapid Biological Assessment of the Konashen Community Owned Conservation Area, Southern Guyana: 21

Published By: Conservation International

URL: <https://doi.org/10.1896/054.051.0105>

BioOne Complete (complete.BioOne.org) is a full-text database of 200 subscribed and open-access titles in the biological, ecological, and environmental sciences published by nonprofit societies, associations, museums, institutions, and presses.

Your use of this PDF, the BioOne Complete website, and all posted and associated content indicates your acceptance of BioOne's Terms of Use, available at www.bioone.org/terms-of-use.

Usage of BioOne Complete content is strictly limited to personal, educational, and non - commercial use. Commercial inquiries or rights and permissions requests should be directed to the individual publisher as copyright holder.

BioOne sees sustainable scholarly publishing as an inherently collaborative enterprise connecting authors, nonprofit publishers, academic institutions, research libraries, and research funders in the common goal of maximizing access to critical research.



Large-headed Ant (*Daceton armigerum*) showing large mandibles



Freshwater Crab (*Fredius* sp.) can sometimes be found outside of water, foraging on the forest floor



Blue Poison Dart Frog (*Dendrobates tinctorum*)



Mouse possum (*Marmosa* sp.)



Forest floor of a typical, annually inundated forest along the Kamao River



Emerald Tree Boa (*Corallus caninus*) coiled in a tree



Ichthyologist Carlos Lasso with a freshly caught catfish



Surinam Toad (*Pipa pipa*) mimicking leaf litter at the bottom of a forest pond



Tukeit Hill Frog (*Allophryne ruthveni*)



Bess Beetle (*Passalus* sp.)



Peacock Katydid (*Pterochroza ocellata*) amid leaf litter



Lobster Katydid (*Panoploscelis specularis*), one of the largest katydids of the Neotropics



A pristine stream in the Acarai Mountains, the first site of the RAP survey




Worm Lizard (*Amphisbaena vanzolini*) is legless and without functional eyes



Smooth-fronted Caiman (*Paleosuchus trigonatus*) young, this species prefers more turbid waters than other species of the genus



Giant Bird-eating Spider (*Theraphosa blondi*), the largest spider in the world



Entomologist Christopher Marshall and assistants collecting symbiotic moths from the fur of Pale-throated Three-toed Sloth (*Bradypus tridactylus*)