



Appendix 6

Source: A Rapid Biodiversity Assessment of the Nakauvadra Range, Ra Province, Fiji: 125

Published By: Conservation International

URL: <https://doi.org/10.1896/054.057.0121>

BioOne Complete (complete.BioOne.org) is a full-text database of 200 subscribed and open-access titles in the biological, ecological, and environmental sciences published by nonprofit societies, associations, museums, institutions, and presses.

Your use of this PDF, the BioOne Complete website, and all posted and associated content indicates your acceptance of BioOne's Terms of Use, available at www.bioone.org/terms-of-use.

Usage of BioOne Complete content is strictly limited to personal, educational, and non - commercial use. Commercial inquiries or rights and permissions requests should be directed to the individual publisher as copyright holder.

BioOne sees sustainable scholarly publishing as an inherently collaborative enterprise connecting authors, nonprofit publishers, academic institutions, research libraries, and research funders in the common goal of maximizing access to critical research.

Appendix 6
Sites of interest for terrestrial insects in the Nakauvadra Range, Ra Province, Fiji
Hilda Waqa-Sakiti

No.	Name	Latitude (S)	Longitude (E)	Elevation (m)	Comments
ento 1	<i>Nesobasis angolicolis</i> <i>Melanesobasis</i> sp.	17°26'15"S	178°05'36"E	406	dry creek, 2° forest edge
ento 2	<i>Papilio schemltzi</i>	17°26'19"S	178°05'42"E	476	1st sighting of <i>P. schemltzi</i> , host plant <i>Micromellum minutum</i> found within study area
ento 3	Leaf litter 1	17°26'27"S	178°05'45"E	541	mid-altitude, on gentle slope
ento 4	Base camp 1 <i>Phasmotaenia inermis</i> (syn. <i>Hermarchus</i>) <i>Liocheles australasiae</i> (scorpion)	17°27'22"S	178°05'54"E	329	near confluence of Volivoli and Nabiya creek
ento 5	Light trap 1	17°27'20"S	178°05'52"E	327	adjacent to Nabiya Creek
ento 6	Leaf litter 2	17°27'29"S	178°05'35"E	503	mid-altitude, on very steep slope
ento 7	Leaf litter 3	17°27'25"S	178°05'55"E	328	low altitude leaf litter sampling
ento 8	Light trap 2	17°27'21"S	178°05'56"E	308	
ento 9	Tree beating 1	17°27'20"S	178°05'56"E	327	
ento 10	Tree beating 2	17°27'21"S	178°05'56"E	309	
ento 11	<i>Nisyrius spinulosus</i> (synonym to <i>Cotylosoma</i>)	17°27'29"S	178°05'11"E	424	found within dakua (<i>Agathis macrophylla</i>) plot camouflaged by lichens on the tree bark
ento 12	<i>P. schmeltzi</i>	17°27'29"S	178°05'57"E	330	numerous sightings of <i>P. scheltzi</i> and <i>Danaus plexipus</i>
ento 13	Base camp 2 Buprestidae (jewel beetle)	17°27'46"S	178°07'46"E	275	A rarely encountered beetle family
ento 14	Light trap 3	17°27'20"S	178°05'52"E	327	
ento 15	Light trap 4	17°27'47"S	178°07'43"E	249	
ento 16	Leaf litter 4	17°27'47"S	178°07'45"E	253	