

## Map

Source: A Rapid Biological Assessment of the Atewa Range Forest

Reserve, Eastern Ghana: 31

Published By: Conservation International

URL: https://doi.org/10.1896/054.047.0105


BioOne Complete (complete.BioOne.org) is a full-text database of 200 subscribed and open-access titles in the biological, ecological, and environmental sciences published by nonprofit societies, associations, museums, institutions, and presses.

Your use of this PDF, the BioOne Complete website, and all posted and associated content indicates your acceptance of BioOne's Terms of Use, available at <a href="https://www.bioone.org/terms-of-use">www.bioone.org/terms-of-use</a>.

Usage of BioOne Complete content is strictly limited to personal, educational, and non - commercial use. Commercial inquiries or rights and permissions requests should be directed to the individual publisher as copyright holder.

BioOne sees sustainable scholarly publishing as an inherently collaborative enterprise connecting authors, nonprofit publishers, academic institutions, research libraries, and research funders in the common goal of maximizing access to critical research.

## Map



## **Photos from the Atewa RAP Survey**

Al photos aken by ot as eck except here noted)



Butterfly (Kallimoides rumia)



A new species of tick spider (Ricinoides sp. n.)



Tadpoles of the clawed frog (Siluarana tropicalis)



A female of *Orthetrum julia* (the Julia Skimmer), one of Africa's most common dragonfly species



Green tree viper (Atheris chlorechis)



Atewa tree fern (Cyathea manniana)



Rhinoceros viper (Bitis nasicornis)



A new species of katydid (Tetraconcha sp. n.)



Mudpuddling male of the Atewa Dotted Border, *Mylothris atewa*, a narrowly endemic species that occurs only in the upland rainforest of the Atewa Range



A driver ant (Dorylus sp.) attacking a termite



A stream of driver ants (Dorylus sp.)



*Hyperolius bobirensis* is known only from two other Ghanaian sites. Characteristics are its large size and the granular back skin.



RAP scientist Natalie Weber with a bat (Hipposidesos gigas)



Tree frogs in amplexus (Chiromantis rufescens)



Decayed leaf katydid (Weissenbornia praestantissima)



Leaf katydid (Poreuomena lamottei)



The Atewa RAP team



The aquatic *Conraua derooi* is Critically Endangered and may have its largest populations in the Atewa Range Forest Reserve