
Bibliography and additional reading

- Agar, I. T., Hess-Pierce, B., Sourour, M. M., and Kader, A. A. (1998). Quality of fruit and oil of black-ripe olives is influenced by cultivar and storage period. *Journal of Agricultural and Food Chemistry* 46(9), 3415–3421.
- Agar, I. T., Hess-Pierce, B., Sourour, M. M., and Kader, A. A. (1999). Identification of optimum preprocessing storage conditions to maintain quality of black-ripe ‘Manzanillo’ olives. *Postharvest Biology and Technology* 15(1), 53–64.
- Allaby, M. (1998). ‘Oxford Dictionary of Plant Sciences.’ (Oxford University Press, Oxford.)
- Aloni, N. (2000). Irrigation and sustainable nutrition – modern olive groves need supplementary nutrients. *The Olive Press, Spring Edition*, 15–16.
- Amelio, M., and De Mauro, E. (2000). Naturally fermented black olives of Taggiasca variety (*Olea europaea* L.). *Grasas Y Aceites* 51(6), 429–438.
- Angerosa, F., Lanza, B., d’Alessandro, N., Marsilio, V., and Cumitini, S. (1999). Olive oil off-odour compounds produced by *Aspergillus* and *Penicillium*. In ‘Proceedings of the 3rd International Symposium on Olive Growing’. (Eds I. T. Metzidakis and D. G. Voyiatzis.) *Acta Horticulturae*, 474(2), 695–699.
- Anon. (2003). Table olives and the world market. *Olivae* 99, 45–47.
- Anon. (2004). Report: Olives in a nutshell. *Choice, July edition*, 24–27.
- Anon. (2005). Reviving olives in Robinvale. *Australian and New Zealand Olivegrower and Processor* 42 (Mar.–Apr.), 31.
- Anon. (2006). Food microbiology and risk assessment conferences. *Food Australia* 58(1–2), 32–40.
- Antognozzi, E., and Proietti, P. (1995). Effects of CPPU (cytokinin) on table olive trees (*Ascolana Tenera*) under nonirrigated and irrigated conditions. In ‘Quality of Fruit and Vegetables – Pre and post-harvest factors and technology’. (Eds D. Gerasopoulos, C. H. Olympos and H. Passam.) *Acta Horticulturae* 379, 159–166.