

## Acknowledgment

We thank J. H. Frank (Department of Entomology, University of Florida, Gainesville), R. C. Froeschner (Department of Entomology, Smithsonian Institution, Washington, D.C. [SI]), R. W. Hodges (Systematic Entomology Laboratory, USDA-ARS, Washington, D.C. [SEL]), I. M. Kerzhner (Zoological Institute, Leningrad), J. A. Slater (Department of Ecology & Evolutionary Biology, University of Connecticut, Storrs), and the late D. R. Whitehead (SEL) for their useful comments on this manuscript. We also are grateful to I. M. Kerzhner for pointing out several nomenclatural changes involving some of the Holarctic Miridae, sharing with us his opinion that *Pithanus hrabei* might occur in North America, and for making available new records from the USSR; to M. D. Schwartz (Canadian National Collection, Agriculture Canada, Ottawa, Ontario) and J. Soroka (Research Station, Agriculture Canada, Saskatoon, Saskatchewan) for searching their collections for specimens of *P. hrabei* and records of other species treated in this paper; to P. Štys (University of Karlovy, Prague) for the record of *C. virgula* males; to P. Lindskog (Naturhistoriska Riksmuseet, Stockholm) and R. T. Schuh (American Museum of Natural History, New York) for comments on the identity of *Psallus* sp. currently identified as *P. albipennis*; to J. H. Frank for allowing us to use the manuscript on biogeographic terminology he and E. D. McCoy had in press during preparation of this manuscript; to J. D. Lattin (Department of Entomology, Oregon State University, Corvallis) for the warm