

5 Subfamily Mirinae

Tribe Mirini

Adelphocoris lineolatus (Goeze)

(Fig. 3)

Distribution (Map 5). Knight (1922) gave Cape Breton Island, Nova Scotia, as the first North American record and (1930a) listed Ames, Ia., as the first U.S. record. Knight (1941) recorded its presence in Minnesota in 1934; Illinois, Missouri, and South Dakota in 1935; Nebraska and Wisconsin in 1936; and Kansas and Manitoba in 1939. Additional records are New Brunswick (Hughes 1943); Quebec (Moore 1950); Ontario (Phillips 1951); Saskatchewan (Arnason et al. 1951); North Dakota and Indiana (USDA 1952a, b); Delaware (USDA 1953a); Maryland (USDA 1953b); New Jersey, Pennsylvania, Virginia, and West Virginia (USDA 1953c); Maine (USDA 1954a); Ohio (USDA 1954c); New York (Neunzig & Gyrisco 1955); Michigan (USDA 1956b); Massachusetts (USDA 1958b); Kentucky (USDA 1960a); Vermont (MacCollom 1958); Wyoming (USDA 1961); Connecticut and Rhode Island (USDA 1969a); Alberta and Montana (Craig 1971); and North Carolina (McPherson et al. 1983).

New record. **United States:** COLORADO: Mesa Co., COLM Fruita Entrance, 29 July 1962, McCoy & Miller (UCM).