

Labor and Inflammation Increase the Expression of Oxytocin Receptor in Human Amnion 1

Authors: Terzidou, Vasso, Blanks, Andrew M., Kim, Sung Hye, Thornton, Steven, and Bennett, Phillip R.

Source: *Biology of Reproduction*, 84(3) : 546-552

Published By: Society for the Study of Reproduction

URL: <https://doi.org/10.1095/biolreprod.110.086785>

BioOne Complete (complete.BioOne.org) is a full-text database of 200 subscribed and open-access titles in the biological, ecological, and environmental sciences published by nonprofit societies, associations, museums, institutions, and presses.

Your use of this PDF, the BioOne Complete website, and all posted and associated content indicates your acceptance of BioOne's Terms of Use, available at www.bioone.org/terms-of-use.

Usage of BioOne Complete content is strictly limited to personal, educational, and non - commercial use. Commercial inquiries or rights and permissions requests should be directed to the individual publisher as copyright holder.

BioOne sees sustainable scholarly publishing as an inherently collaborative enterprise connecting authors, nonprofit publishers, academic institutions, research libraries, and research funders in the common goal of maximizing access to critical research.

Labor and Inflammation Increase the Expression of Oxytocin Receptor in Human Amnion¹

Vasso Terzidou,^{2,3} Andrew M. Blanks,⁴ Sung Hye Kim,³ Steven Thornton,⁴ and Phillip R. Bennett³

Imperial College London,³ Parturition Research Group, Institute of Reproductive and Developmental Biology, Hammersmith Hospital Campus, London, United Kingdom
University of Warwick,⁴ Clinical Sciences Research Institute, Warwick Medical School, UHCW, Coventry, United Kingdom

ABSTRACT

The oxytocin/oxytocin receptor (OXT/OXTR) system plays an important role in the regulation of parturition. The amnion is a major source of prostaglandins and inflammatory cytokine synthesis, which increase both before and during labor. Amnion is a noncontractile tissue; therefore, the role played by OXT/OXTR in this tissue will be fundamentally different from the role played in myometrial contractions. In the present study, we demonstrate increased *OXTR* mRNA and protein concentrations in human amnion epithelial cells associated with the onset of labor. We show that incubation of primary human amnion epithelial cells with IL1B results in a rapid, transient up-regulation of *OXTR* mRNA expression, which peaks in prelabor samples after 6 h. Incubation of prelabor amnion epithelial cells with OXT results in a marked increase of prostaglandin E₂ synthesis, and we demonstrate that OXT activates the extracellular signal-regulated protein kinase signal transduction pathway to stimulate up-regulation of cyclo-oxygenase 2 in human amnion epithelial cells. The increased ability of human amnion to produce prostaglandins in response to OXT treatment suggests a complementary role for the OXT/OXTR system in the activation of human amnion and the onset of labor.

amnion, oxytocin, oxytocin receptor, parturition, prostaglandins

INTRODUCTION

The oxytocin receptor (OXTR) is a member of the G protein-coupled superfamily, and uterine sensitivity to oxytocin (OXT) markedly increases around the onset of labor. This is associated with both an up-regulation of *OXTR* mRNA levels and a strong increase in myometrial OXTR density, reaching a peak during early labor [1]. The physiological role of the OXT/OXTR system in myometrium is well understood, because OXT represents one of the best-known contraction mediators. The increase in OXTR before labor is not confined to the myometrium. In rabbit amnion, a 200-fold up-regulation of

OXTR occurs at the end of pregnancy [2], and OXTR binding to OXT significantly increases in human fetal membranes with the onset of labor [3]. Studies in several species have shown OXT to stimulate uterine prostaglandin (PG) synthesis when administered near estrus and to stimulate PGE₂ production. This suggests a role for OXT/OXTR in the activation of the amnion that occurs at the time of labor.

The amnion plays an important role in the onset of human labor. It is a major source of PG and inflammatory cytokine synthesis, which increase both before and during labor. Increased PG synthesis in the amnion appears to occur first in the region overlying the cervix, where it is thought to mediate cervical ripening [4]. PGs, especially PGE₂ synthesized in the amnion, are thought to cross to the decidua and myometrium to mediate cervical ripening and lower segment remodeling and initiate contractions. Roseblade et al. [5] have suggested that when using physiological concentrations of PGE₂ in an in vitro model, only small amounts cross the membranes without being metabolized. However, Johnston et al. [6] have demonstrated marked transfer of PGE₂ across ovine fetal membranes in vivo after intra-amniotic PGE₂ injection. Amnion PG synthesis occurs principally via cyclo-oxygenase 2 (PTGS2), which itself is regulated by nuclear factor kappa B (NFkB) [7–10]. Labor, both term and preterm, is associated with increased expression and activity of PTGS2 in the amnion [11], and basal NFkB DNA binding and transcriptional activity increase in human amnion with labor onset. The NFkB changes persist in culture and represent a form of amnion differentiation, which we believe commits to labor and delivery [7]. Both basal and IL1B-induced NFkB DNA binding involves the p50-p65 heterodimer as well as the p65 and p50 homodimers [12]. Using whole-genome cDNA arrays, we recently studied the range of genes for which expression is increased in amnion with high levels of NFkB activity compared to low levels. OXTR expression showed the second-greatest difference, with a 24-fold increase in expression (after PTGS2, which was increased 40-fold), therefore strongly suggesting a role for NFkB in the regulation of OXTR in amnion [13].

Mitogen-activated protein kinases (MAPKs) are important signal pathway components and function as integrators of mitogenic and other signals originating from G protein-coupled receptors. Members from the three subgroups of the MAPK family—extracellular signal-regulated protein kinase (ERK), stress-activated protein kinase c-Jun NH₂-terminal kinase (MAPK8), and p38 MAPK (MAPK14)—mediate responses to proinflammatory stimuli and affect gene expression at the transcriptional and posttranscriptional levels. Each subtype of the MAPK family is activated by its own kinase system. The activation of MAPK3/1 (ERK1/2) involves activation of RAF followed by MAPK kinase (MAP2K) 1/2. When MAPK3/1

¹Supported by an Action Medical Research Project Grant (SP4454) and a National Institute for Health, Biomedical Research Centre Grant (DSRR_NG0122/2009).

²Correspondence: V. Terzidou, Parturition Research Group, Institute of Reproductive and Developmental Biology, Hammersmith Hospital Campus, Du Cane Road, East Acton, London W12 0NN, U.K. FAX: 44 207 5942189; e-mail: v.terzidou@imperial.ac.uk

Received: 21 June 2010.

First decision: 23 July 2010.

Accepted: 4 October 2010.

© 2011 by the Society for the Study of Reproduction, Inc.

This is an Open Access article, freely available through *Biology of Reproduction's* Authors' Choice option.

eISSN: 1529-7268 <http://www.biolreprod.org>

ISSN: 0006-3363

gets activated, it exerts its effects via phosphorylation of proteins and/or transcription factors that can alter gene expression [14].

In rat myometrium during pregnancy, regulation of ERK activity is controversial. It has been reported to increase with advancing gestation, although some studies have documented a decline immediately before the onset of parturition [15], in contrast to others that have documented a rise with the onset of labor [16]. In nonpregnant women, expression of MAPK14 and MAPK3/1 was uniform throughout the uterus, whereas in pregnant women before and after labor onset, expression of MAPK14 and MAPK3 was significantly elevated in the upper compared to the lower uterine segment [17]. The role of OXT in MAPK activation is not well defined. Cytosolic phospholipase A₂ (cPLA₂) is an important enzyme providing substrate for cyclo-oxygenases and is activated by various agents that stimulate its phosphorylation on Ser505 by MAPKs [18, 19].

In the present study, we have investigated the labor-associated changes in OXTR expression and the effect of IL1B in human amnion epithelial cells. We provide evidence suggesting for a role of the OXT/OXTR system in PG production in human amnion, and we propose that an ERK-mediated PTGS2 up-regulation is involved in this process.

MATERIALS AND METHODS

Cell Preparation and Culture

Fetal membranes were obtained by elective cesarean section before labor at term (prelabor) or after spontaneous vaginal delivery at term (postlabor), where term was defined as 37–42 completed weeks of pregnancy. Institutional ethics committee approval was granted for the present study, and patients gave informed consent. Amnion epithelial cells were prepared from tissue as previously described [20]. In brief, amnion was separated from chorion and washed in PBS. Next, the membrane was cut into strips and incubated in 0.5 mmol/L of ethylenediaminetetra-acetic acid (EDTA; BDH Laboratory Supplies Ltd.) for 15 min at room temperature. The strips were then rinsed twice in PBS, followed by incubation in 2.5 g/L of dispase (Life Technologies) for 40 min at 37°C. Amnion epithelial cells were isolated by vigorous shaking for 3 min, after which strips were removed. Individual cells were collected by centrifugation at 400 × g for 10 min and grown in Dulbecco modified Eagle medium supplemented with 10% fetal calf serum (Sigma), 2 mmol/L of L-glutamine, 100 U/ml of penicillin, and 100 µg/ml of streptomycin (Life Technologies) at 5% CO₂. All amnion epithelial cells used were primary cultures (no passages) and were usually cultured for 3–4 days before treatment.

Real-Time RT-PCR

Total RNA was extracted using RNA STAT-60 reagent (AMS Biotechnology) according to the manufacturer's specifications. After quantification, 1 µg of total RNA was digested with 0.5 µl of DNase I in 1× DNase I reaction buffer (Invitrogen) in a total volume of 5 µl at room temperature for 15 min. The reaction was terminated by the addition of 0.5 µl of 25 mM EDTA and incubation at 65°C for 15 min. The whole of this reaction was subsequently used for first-strand cDNA synthesis with SuperScript First-Strand Synthesis System for RT-PCR (Invitrogen) according to the manufacturer's instructions. Gene expression was verified by real-time RT-PCR using the ABI PRISM 7700 Sequence Detection System (Applied Biosystems). TaqMan primers and probes to analyze the different transcripts were designed using the Primer Express software (Applied Biosystems). The data were analyzed using Sequence Detector version 1.7 software (Applied Biosystems), and the values obtained were normalized according to transcript levels of glyceraldehyde phosphate dehydrogenase (GAPDH) or ribosomal protein L19. RT-PCR expression data were assessed using the delta Ct method. The sequences used are shown in Table 1. Statistical significance was determined by ANOVA.

Protein Extraction, SDS-PAGE, Western Blot Analysis, and Immunodetection

Confluent amnion epithelial cells were rinsed in PBS and lysed in a buffer containing 10 mM Hepes, 10 mM KCl, 0.1 mM EDTA, 0.1 mM ethylene glycol tetraacetic acid, 2 mM dithiothreitol (DTT), 1% (v/v) Nonidet P-40 (NP-

40), and Complete Protease Inhibitor Tablets (Roche Diagnostics). Cell lysates were incubated on ice for 10 min, and NP-40 was added to a final concentration of 1% (v/v). Lysates were vortexed and centrifuged for 30 sec at 4°C and 12000 × g. The supernatants were retained as the cytosolic protein fraction. The pellets were resuspended in a buffer containing 10 mM Hepes, 10 mM KCl, 0.1 mM EDTA, 0.1 mM ethylene glycol tetraacetic acid, 2 mM DTT, 400 mM NaCl, 1% (v/v) NP-40, and Complete Protease Inhibitor Tablets. Samples were then shaken vigorously for 15 min on ice. Nuclear protein extracts were obtained in the supernatant after centrifugation for 5 min at 12000 × g and 4°C. For whole-cell protein, cells were lysed on ice for 20 min in radioimmuno-precipitation assay buffer (1% NP-40, 1% sodium deoxycholate, 1% Triton X-100, 0.1% SDS, 150 mM NaCl, 10 mM Tris [pH 8.0], and 2 mM NaF). Protein concentrations were determined using BioRad assay (Bio-Rad Laboratories). Protein samples (40 µg) were denatured by boiling for 5 min and then separated by electrophoresis on an 8% SDS-PAGE gel for 80 min at 150 V. Transfer from gel to polyvinylidene fluoride membrane (Anachem) took place in a semidry chamber with three buffer systems as described by Pohnke et al. [21]. The membrane was activated in methanol, blocked in 5% milk protein solution (Marvel Lincs) for 1 h at room temperature, and then washed and hybridized with the primary antibodies (1:1000 dilution) overnight at 5°C in a fresh blocking buffer (1× PBS, 1% milk protein, and 0.1% Tween-20), followed by incubation with horseradish peroxidase (HRP)-conjugated secondary antibodies (1:2000 dilution). Signal detection was carried out using ECL Plus (Amersham). To confirm equal loading of each well, the blot membrane was treated with a stripping buffer (2% SDS, 62.5 mM Tris-HCl [pH 6.7], and 100 mM 2-mercaptoethanol) for 30 min at 50°C, washed in PBS-Tween-20, and then preblocked and reprobed with an antibody to human β-actin.

Antibodies and Materials

The following antibodies were obtained from Santa Cruz Biotechnologies: goat anti-PTGS2 (COX-2) (C20), goat anti-OXTR (C-terminus), and HRP-conjugated secondary antibodies raised against goat, rabbit, and mouse immunoglobulin Gs. Rabbit monoclonal antibodies to phospho-cPLA₂, phospho-MAPK14 (p38 MAPK), phospho-MAPK3/1 (ERK1/2 p44/42 MAPK), and mouse monoclonal MAPK8 (SAPK/JNK) were from Cell Signaling Technology. The monoclonal anti-smooth muscle-actin antibody was from Sigma. PGE₂ enzyme immunoassays were from Amersham (GE Healthcare UK Ltd.). Cells before treatment with OXT (Alliance Pharmaceuticals) or IL1B (R&D Systems) were serum-deprived for 16 h and stimulated for the required times.

Statistical Analysis

Multiple comparisons were analyzed using one-way ANOVA. The effects of OXT and IL1B treatments with nonstimulated effects were compared using *t*-test. Differences of *P* < 0.05 were considered to be statistically significant.

RESULTS

OXTR Is Expressed in Humans, and Both mRNA and Protein Expression Significantly Increase in Association with Labor

Primary human amnion epithelial cell cultures were established from patients before and after the onset of labor. Using real-time RT-PCR, we found that the expression of *OXTR* mRNA in postlabor primary amnion epithelial cells was 16-fold higher than in prelabor cells (*P* < 0.001; *n* = 4 sets of amnion epithelial cells isolated from 4 patients each group, duplicate samples) (Fig. 1A). Whole-cell protein was extracted, and Western blot analysis demonstrated that *OXTR* protein expression also significantly increased in association with labor (Fig. 1, B and C).

IL1B Increases OXTR Expression in Prelabor Amnion to Postlabor Levels But Does Not Further Increase Expression in Postlabor Cells


Time-course experiments were performed to assess *OXTR* mRNA expression after treatment with IL1B (1 ng/ml) for 30 min as well as 1, 2, 4, 6, and 24 h before and after the onset of labor. Treatment with IL1B resulted in a significant up-

TABLE 1. The primer pairs, probes, and GenBank accession numbers for the primer pairs used.

mRNA	Sequences*	GenBank/EMBL Accession no.
<i>OXTR</i>	Primers	
	F : GATGGGAAGGGTGGT R : CAAGGACCCAGCATTTGTT	NM_000916
<i>PTGS2</i>	F : GCTCAAACATGATGTTTGCATT R : GCTGGCCCTCGCTTATGA	AY151286
	F : TCTTAGCCCTTGGATTCCT R : ATTCTTAGCCCGGGATTCAG	BC008280
<i>GAPDH</i>	F : GAAGGTGAAGTCCGGAGT R : GAAGATGGTGATGGGATTTTC	XM_068376
	F : GCGGAAGGTACAGCCAAT R : GCAGCCGGCGCAA	NM_000981
<i>OXTR</i>	Probes	
	CCTCCTGACCTCAAAGTGTATTTGCCTTTAAGC ATTTGGTCGTATTGGGCGCCTGGTCACC	NM_000916 XM_068376

* F, Forward; R, Reverse

FIG. 1. Expression of *OXTR* in human amnion epithelial cells. **A)** Expression of *OXTR* mRNA measured by real-time RT-PCR in prelabor (L-) and postlabor (L+) amnion epithelial cells. Both L- and L+ cells were treated with IL1B (1 ng/ml) for 6 h. Results are corrected for *GAPDH* expression and given as the fold-increase in *OXTR/GAPDH* mRNA expression (n = 4 sets of amnion epithelial cells isolated from 4 patients each group, duplicate samples; ANOVA: ^a*P* < 0.05, L- a IL1B-stimulated vs. nonstimulated cells; ^b*P* < 0.05, L- vs. L+ cells. **B)** Western blot analysis for *OXTR* in amnion epithelial cells taken before (-) and after (+) the onset of labor. Controls with β -actin confirmed equal protein loading. **C)** Densitometric quantification of *OXTR* protein expression normalized against actin protein levels (n = 3 sets of amnion epithelial cells isolated from 3 patients; ANOVA: **P* < 0.05). **D)** Time course of *OXTR* mRNA measured by real-time RT-PCR in prelabor (-) amnion epithelial cells after treatment with IL1B (1 ng/ml). Results are corrected for *GAPDH* expression and given as the percentage of untreated cells for each time point (n = 4 sets of amnion epithelial cells isolated from 4 patients in each group, duplicate samples; ANOVA: **P* < 0.05 vs. nonstimulated cells). Treatment with IL1B resulted in a significant up-regulation of *OXTR* expression that peaked 6 h after treatment.






FIG. 2. OXT-stimulated PGE₂ production in human amnion epithelial cells. PGE₂ production by prelabor cells after treatment with OXT (10⁻⁷ M) for 15 min, 30 min, 2 h, and 6 h is shown. Results are expressed as the percentage of control (nonstimulated cells) production for each time point (n = 3 sets of amnion epithelial cells isolated from 3 patients, triplicate samples; ANOVA: *P < 0.05).

regulation of *OXTR* mRNA expression. This peaked in the prelabor cells at 6 h after treatment and resulted in a 17-fold induction of *OXTR* expression (P < 0.001; n = 4 sets of amnion epithelial cells isolated from 4 patients, duplicate samples) (Fig. 1D). The level of expression that was observed in the prelabor cells was significantly less than that in the postlabor cells. Treatment with IL1B in prelabor cells brought the level of *OXTR* expression to that observed in postlabor cells. In postlabor cells, IL1B caused a further, but nonsignificant, increase in *OXTR* expression (Fig. 1A).

PGE₂ Release by Primary Human Amnion Significantly Increases after OXT Stimulation

Time-course experiments were performed with prelabor cells to assess the effect of OXT treatment in PGE₂ production. OXT (10⁻⁷ M) stimulation for 6 h resulted in a 40-fold increase of PGE₂ production (P < 0.05; n = 3 sets of amnion epithelial cells isolated from 3 patients, triplicate samples) (Fig. 2).

Changes in *PTGS2* and Phospho-cPLA₂ in Amnion after OXT Stimulation

Amnion epithelial cells from patients before the onset of labor were treated for 6 h with either OXT (10⁻⁷ M) or IL1B (1 ng/ml). Treatment with IL1B resulted in a 3-fold increase in *PTGS2* mRNA expression (P < 0.001; n = 7 sets of amnion epithelial cells isolated from 7 patients, duplicate samples) and OXT induced a 2-fold increase in *PTGS2* mRNA expression (P < 0.005; n = 7, as described above) (Fig. 3A), whereas the levels of PGE synthase (*PTGES*) mRNA were not affected (Fig. 3B). Western blot analysis confirmed this increase in protein level (Fig. 3, C and F). Phospho-cPLA₂ also increased after OXT treatment (Fig. 3, D and G). Western blot analysis for β-actin confirmed equal loading (Fig. 3E).

OXT Activates the MAPK Pathway in Amnion Epithelial Cells, Specifically via ERK

To determine the identity of the MAPK potentially involved in OXT-stimulated *PTGS2* up-regulation, immunoblotting of amnion cell lysates was performed using antibodies to dually




FIG. 3. OXT stimulation effects in the PG biosynthetic pathway in human amnion epithelial cells. Prelabor cells were treated with OXT (10⁻⁷ M) or IL1B (1 ng/ml) for 6 h, after which mRNA was extracted and real-time RT-PCR was performed. **A**) Real-time RT-PCR for *PTGS2*. Results are corrected for ribosomal protein L19; treatment with IL1B resulted in a 3-fold increase in *PTGS2* mRNA expression (n = 7 sets of amnion epithelial cells isolated from 7 patients/duplicate samples; ANOVA: *P < 0.001 vs. stimulated cells), and OXT induced a 2-fold increase in *PTGS2* mRNA expression (n = 7, as above; ANOVA: *P < 0.005 vs. stimulated cells). **B**) Real-time RT-PCR for *PTGES*. Results are corrected for L19. **C–E**) Prelabor amnion epithelial cells were treated with OXT (10⁻⁷ M) or IL1B (1 ng/ml) for 6 h, after which cytoplasmic protein was extracted. Western blot analysis for *PTGS2* (C), phospho-cPLA₂ (D), and β-actin (E; control) is shown. OXT stimulation up-regulates both enzymes, as does treatment with IL1B. *PTGS2* is degraded when it catalyzes PG synthesis, which explains the down-regulation observed after 24 h of treatment. Controls with β-actin confirmed protein equal loading. **F** and **G**) Densitometric quantification of *PTGS2* and cPLA₂ protein expression normalized against actin protein levels (n = 3 sets of amnion epithelial cells isolated from 3 patients; ANOVA: *P < 0.02 vs. nonstimulated cells).

phosphorylated MAPK3/1, phospho-MAPK14, and phospho-MAPK8. We found that OXT stimulation increases phospho-MAPK3/1, whereas phospho-MAPK14 and phospho-MAPK8 were unaffected (Fig. 4, A–C). Pretreatment of amnion




FIG. 4. OXT stimulation effects in the MAPK pathway in human amnion epithelial cells. Prelabor cells were treated with OXT (10^{-7} M) or IL1B (1 ng/ml), after which cytoplasmic protein was extracted. **A–D**) Western blot analysis for phospho-MAPK8 (**A**), MAPK14 (**B**), phospho-MAPK3/1 (**C**), and β -actin (**D**; control). Controls with β -actin confirmed equal protein loading. **E**) Densitometric quantification of phospho-MAPK3 and phospho-MAPK1 expression normalized against actin protein levels ($n = 3$ sets of amnion epithelial cells isolated from 3 patients; ANOVA: $*P < 0.05$ vs. nonstimulated cells).

epithelial cells with the MAP2K1/2 inhibitor UO126 (10 mM; Sigma) for 2 h before OXT stimulation resulted in a significant decrease in the OXT-stimulated increase of PTGS2 (Fig. 5, A and B).

DISCUSSION

We have demonstrated an increase in *OXTR* mRNA and protein concentrations in human amnion epithelial cells associated with the onset of labor, as has been previously shown in human choriodecidua and fundal myometrium [1, 22, 23].

In the present study, incubation of primary human amnion epithelial cells with IL1B results in a rapid, transient up-regulation of *OXTR* mRNA expression. This peaked in prelabor amnion epithelial cells at 6 h after treatment and resulted in a 17-fold induction of *OXTR* mRNA expression. This response of *OXTR* expression to IL1B is similar to that which we have previously reported in myometrium [15], and further supports the concept that *OXTR* is regulated in a paracrine as well as endocrine fashion. Rauk et al. [24] as well as Mitchell and Schmid [25] have reported that IL1B down-regulates *OXTR* mRNA in myometrial cells. Schmid et al. [26]




FIG. 5. MAP2K1/2 inhibition of the OXT-induced PTGS2 up-regulation in human amnion epithelial cells. Prelabor cells were pretreated with the MAP2K1/2 inhibitor UO126 (Sigma) for 2 h before OXT stimulation (10^{-7} M), after which cytoplasmic protein was extracted. **A** and **B**) Western blot analysis for PTGS2 (**A**) and β -actin (**B**; control). Controls with β -actin confirmed protein equal loading. **C**) Densitometric quantification of PTGS2 expression normalized against actin protein levels ($n = 3$ sets of amnion epithelial cells isolated from 3 patients; ANOVA: $*P < 0.04$, IL1B-treated vs. nonstimulated cells; $^{\circ}P < 0.05$, OXT-treated vs. nonstimulated cells; $^{\S}P < 0.05$, UO126+OXT-treated vs. OXT-treated cells).

used an immortalized cell line derived from nonpregnant human myometrium, whereas Rauk et al. [24], using pregnant primary myocytes and measuring *OXTR* mRNA by Southern blot analysis, showed an early, transient increase in *OXTR* expression, followed by a significant decrease after 25 h. To us, this seems to be counterintuitive. It is well established that IL1B concentrations within the uterus increase at the time of both term and preterm labor [27], as does the expression of several “labor-associated proteins,” such as PTGS2 and IL8, each of which is up-regulated by IL1B [8, 10, 28–30]. *OXTR* expression is also up-regulated at term, at a time when IL1B concentrations are high and PTGS2 and IL8 expression increases. Our data suggest that IL1B increases *OXTR* expression in primary human amnion epithelial cells. The time course of the effect is similar to that seen with stimulation of both PTGS2 and IL8 by IL1B [28, 31]. The level of expression that was observed in the prelabor amnion epithelial cells was significantly less than that in the postlabor cells. Incubation with IL1B in prelabor cells brought the level of *OXTR* expression to that observed in postlabor cells. In postlabor cells, IL1B caused only a small, nonsignificant further increase in *OXTR* expression. Loudon et al. [28] have previously shown that PGE₂ production is lower in prelabor than in postlabor

cells and that PGE₂ production in prelabor cells can be stimulated by IL1B to levels similar to that found in postlabor cells. This shows that the pathways for PG synthesis are activated in postlabor cells, and our present data suggest that up-regulation of *OXTR* is a feature of amnion activation and is sensitive to inflammatory cytokines.

Amnion is not a contractile tissue; therefore, the physiological role of the OXT/OXTR system in amnion must be in some respect different from its role in myometrium. In rabbit amnion [2], up-regulation of *OXTR* occurs at the end of pregnancy, and OXT has been shown to stimulate PGE₂ production. Similarly, we have found that release of PGE₂ by human amnion epithelial cells is significantly increased after OXT stimulation. This supports a role for OXT/OXTR in the activation of the amnion that occurs at the time of labor. The present study shows that OXT causes increased PG synthesis through up-regulation of PTGS2. It is established that the PTGS2 enzyme, which mediates the committing and rate-limited step of PG biosynthesis, generating a PGH₂ intermediate that is converted to the terminal PGs, is central to increased PGs synthesis in the human amnion at the time of labor [10, 32, 33].

In cultured bovine endometrial cells, OXT has been shown to stimulate PGF_{2α} via activation of the phospholipase C second-messenger cascade, activation of phospholipase A₂ [34], and induction of PTGS2 [35]. In ovariectomized ewes, injection of OXT induced an increase in serum PGFM (stable metabolite of PGF_{2α}), which was associated with an increase in endometrial concentrations of PTGS2 [36]. In rabbit [37], the OXT-stimulated increase in PGE₂ is virtually immediate, with concentrations in the medium reaching near-maximal values by 15 min, which is too rapid to involve new synthesis of PTGS2. This suggests that in rabbit, OXT leads to increased availability of PG precursors, which are synthesized into PGs via pre-existing PTGS2. Our studies in human amnion show that PG synthesis occurred between 2 and 6 h following OXT treatment and was associated with increased PTGS2 expression but no changes in PTGES expression. This shows that in human amnion, unlike rabbit amnion but similar to bovine endometrium, transcriptional regulation of PTGS2 is required for the OXT-stimulated induction of PG synthesis.

We previously demonstrated that OXT can be extracted and measured from the amnion in samples taken both before and after the onset of spontaneous labor [23]. We demonstrated that labor was associated with a significant increase in OXT peptide in amnion, chorion, and decidua. The source of the measured OXT, which acts as ligand to the OXTR in amnion, is unclear, but three possibilities exist. First, neurohypophysial OXT may diffuse from the maternal circulation across the amnion to the epithelium. Although the amnion is avascular, we have found large amounts of OXT to be present in decidua, chorion, and amnion where the placenta was delivered shortly after administration of a bolus of exogenous Syntocinon (Novartis Pharmaceuticals UK Ltd) to the mother, suggesting that OXT (and OXT analogues) can diffuse from the maternal circulation to the amnion (unpublished data). Second, it has been proposed that the fetus may secrete OXT into the amniotic fluid [38]. Third, we have demonstrated that choriodecidua is capable of synthesizing full neurophysin-OXT peptide directly [23]. Studies in rat uterine epithelial cells have demonstrated a clear polarization in OXT and OXTR expression, whereby OXT is secreted from the apical membrane to act on apical OXTR via an autocrine loop [39]. This is particularly pertinent, because in rodents, the uterine epithelium is the primary source of PGs and produces the PGF_{2α} surge required to initiate luteolysis, progesterone withdrawal, and the onset of labor.

In CHO cells transfected with the rat OXTR, stimulation with OXT resulted in the specific phosphorylation and activation of MAPK1 [40]. We have found that OXT activated the ERK signal transduction pathway to stimulate up-regulation of PTGS2 in human amnion epithelial cells. We also have found both MAPK3 and MAPK1 to be activated in human amnion after 2 h of OXT treatment, which does not apply to the other principal MAPKs, MAPK14 and MAPK8.

In summary, we have found that human *OXTR* expression increases in postlabor amnion epithelial cells and that treatment with IL1B stimulates *OXTR* expression before the onset of labor. The increased ability of human amnion to produce PGE₂ in response to OXT treatment suggests a complementary role of the OXT/OXTR system in the activation of human amnion and in the onset of labor. In addition to its role in mediating contractions, the role of OXT in biochemical processes that lead to the onset of labor suggests that the potential clinical use of OXT antagonists requires re-evaluation.

REFERENCES

- Kimura T, Tanizawa O, Mori K, Brownstein MJ, Okayama H. Structure and expression of a human oxytocin receptor. *Nature* 1992; 356:526–529 [published erratum appears in *Nature* 1992; 357:176].
- Hinko A, Soloff MS. Characterization of oxytocin receptors in rabbit amnion involved in the production of prostaglandin E₂. *Endocrinology* 1992; 130:3547–3553.
- Benedetto MT, De Cicco F, Rossiello F, Nicosia AL, Lupi G, Dell'Acqua S. Oxytocin receptor in human fetal membranes at term and during labor. *J Steroid Biochem* 1990; 35:205–208.
- Van Meir CA, Ramirez MM, Matthews SG, Calder AA, Keirse MJ, Challis JR. Chorionic prostaglandin catabolism is decreased in the lower uterine segment with term labor. *Placenta* 1997; 18:109–114.
- Roseblade CK, Sullivan MH, Khan H, Lumb MR, Elder MG. Limited transfer of prostaglandin E₂ across the fetal membrane before and after labor. *Acta Obstet Gynecol Scand* 1990; 69:399–403.
- Johnston PC, Greer IA, Brooks AN. The transfer of prostaglandin E₂ across ovine fetal membranes in vivo. *J Soc Gynecol Invest* 1996; 3: 303–308.
- Allport VC, Pieber D, Slater DM, Newton R, White JO, Bennett PR. Human labor is associated with nuclear factor-kappaB activity which mediates cyclo-oxygenase-2 expression and is involved with the “functional progesterone withdrawal.” *Mol Hum Reprod* 2001; 7:581–586.
- Slater DM, Dennes WJ, Campa JS, Poston L, Bennett PR. Expression of cyclo-oxygenase types-1 and -2 in human myometrium throughout pregnancy. *Mol Hum Reprod* 1999; 5:880–884.
- Sawdy RJ, Slater DM, Dennes WJ, Sullivan MH, Bennett PR. The roles of the cyclo-oxygenases types one and two in prostaglandin synthesis in human fetal membranes at term. *Placenta* 2000; 21:54–57.
- Slater D, Dennes W, Sawdy R, Allport V, Bennett P. Expression of cyclo-oxygenase types-1 and -2 in human fetal membranes throughout pregnancy. *J Mol Endocrinol* 1999; 22:125–130.
- Zakar T, Olson DM, Teixeira FJ, Hirst JJ. Regulation of prostaglandin endoperoxide H₂ synthase in term human gestational tissues. *Acta Physiol Hung* 1996; 84:109–118.
- Lindstrom TM, Bennett PR. The role of nuclear factor kappa B in human labor. *Reproduction* 2005; 130:569–581.
- Lim S, Khanjani S, Terzidou V, Lee YS, Teoh T, Bennett PR. Whole-genome array and si-RNA investigation of the function of NF-kB in human amnion epithelial cells. *Reprod Sci* 2008; 15:260.
- Anderson NG, Maller JL, Tonks NK, Sturgill TW. Requirement for integration of signals from two distinct phosphorylation pathways for activation of MAP kinase. *Nature* 1990; 343:651–653.
- Ruzycky AL. Down-regulation of the mitogen-activated protein kinase cascade immediately before parturition in the rat myometrium. *J Soc Gynecol Invest* 1998; 5:304–310.
- Li Y, Je HD, Malek S, Morgan KG. ERK1/2-mediated phosphorylation of myometrial caldesmon during pregnancy and labor. *Am J Physiol Regul Integr Comp Physiol* 2003; 284:R192–R199.
- Otun HA, MacDougall MW, Bailey J, Europe-Finner GN, Robson SC. Spatial and temporal expression of the myometrial mitogen-activated protein kinases p38 and ERK1/2 in the human uterus during pregnancy and labor. *J Soc Gynecol Invest* 2005; 12:185–190.

18. Lin LL, Wartmann M, Lin AY, Knopf JL, Seth A, Davis RJ. cPLA2 is phosphorylated and activated by MAP kinase. *Cell* 1993; 72:269–278.
19. Lin LL, Lin AY, Knopf JL. Cytosolic phospholipase A₂ is coupled to hormonally regulated release of arachidonic acid. *Proc Natl Acad Sci U S A* 1992; 89:6147–6151.
20. Bennett PR, Rose MP, Myatt L, Elder MG. Preterm labor: stimulation of arachidonic acid metabolism in human amnion cells by bacterial products. *Am J Obstet Gynecol* 1987; 156:649–655.
21. Pohnke Y, Kempf R, Gellersen B. CCAAT/enhancer-binding proteins are mediators in the protein kinase A-dependent activation of the decidual prolactin promoter. *J Biol Chem* 1999; 274:24808–24818.
22. Chibbar R, Miller FD, Mitchell BF. Synthesis of oxytocin in amnion, chorion, and decidua may influence the timing of human parturition. *J Clin Invest* 1993; 91:185–192.
23. Blanks AM, Vatish M, Allen MJ, Ladds G, de Wit NC, Slater DM, Thornton S. Paracrine oxytocin and estradiol demonstrate a spatial increase in human intrauterine tissues with labor. *J Clin Endocrinol Metab* 2003; 88:3392–3400.
24. Rauk PN, Friebe-Hoffmann U. Interleukin-1beta down-regulates the oxytocin receptor in cultured uterine smooth muscle cells. *Am J Reprod Immunol* 2000; 43:85–91.
25. Mitchell BF, Schmid B. Oxytocin and its receptor in the process of parturition. *J Soc Gynecol Investig* 2001; 8:122–133.
26. Schmid B, Wong S, Mitchell BF. Transcriptional regulation of oxytocin receptor by interleukin-1beta and interleukin-6. *Endocrinology* 2001; 142:1380–1385.
27. Romero R, Mazor M, Brandt F, Sepulveda W, Avila C, Cotton DB, Dinarello CA. Interleukin-1alpha and interleukin-1beta in preterm and term human parturition. *Am J Reprod Immunol* 1992; 27:117–123.
28. Loudon JA, Elliott CL, Hills F, Bennett PR. Progesterone represses interleukin-8 and cyclo-oxygenase-2 in human lower segment fibroblast cells and amnion epithelial cells. *Biol Reprod* 2003; 69:331–337.
29. Gilroy DW, Saunders MA, Wu KK. COX-2 expression and cell cycle progression in human fibroblasts. *Am J Physiol Cell Physiol* 2001; 281:C188–C194.
30. Slater D, Allport V, Bennett P. Changes in the expression of the type-2 but not the type-1 cyclo-oxygenase enzyme in chorion-decidua with the onset of labor. *Br J Obstet Gynaecol* 1998; 105:745–748.
31. Bartlett SR, Sawdy R, Mann GE. Induction of cyclo-oxygenase-2 expression in human myometrial smooth muscle cells by interleukin-1beta: involvement of p38 mitogen-activated protein kinase. *J Physiol* 1999; 520(pt 2):399–406.
32. Slater DM, Berger LC, Newton R, Moore GE, Bennett PR. Expression of cyclo-oxygenase types 1 and 2 in human fetal membranes at term. *Am J Obstet Gynecol* 1995; 172:77–82.
33. Slater D, Berger L, Newton R, Moore G, Bennett P. The relative abundance of type 1 to type 2 cyclo-oxygenase mRNA in human amnion at term. *Biochem Biophys Res Commun* 1994; 198:304–308.
34. Lee JS, Silvia WJ. Cellular mechanisms mediating the stimulation of ovine endometrial secretion of prostaglandin F_{2alpha} in response to oxytocin: role of phospholipase A₂. *J Endocrinol* 1994; 141:491–496.
35. Asselin E, Drolet P, Fortier MA. Cellular mechanisms involved during oxytocin-induced prostaglandin F_{2alpha} production in endometrial epithelial cells in vitro: role of cyclo-oxygenase-2. *Endocrinology* 1997; 138:4798–4805.
36. Burns PD, Tsai SJ, Wiltbank MC, Hayes SH, Graf GA, Silvia WJ. Effect of oxytocin on concentrations of prostaglandin H synthase-2 mRNA in ovine endometrial tissue in vivo. *Endocrinology* 1997; 138:5637–5640.
37. Jeng YJ, Liebenthal D, Strakova Z, Ives KL, Hellmich MR, Soloff MS. Complementary mechanisms of enhanced oxytocin-stimulated prostaglandin E₂ synthesis in rabbit amnion at the end of gestation. *Endocrinology* 2000; 141:4136–4145.
38. Dawood MY, Wang CF, Gupta R, Fuchs F. Fetal contribution to oxytocin in human labor. *Obstet Gynecol* 1978; 52:205–209.
39. Morel G, Pechoux C, Raccurt M, Zingg HH. Intrauterine oxytocin system. Compartmental distribution of oxytocin and oxytocin receptors in rat endometrial epithelium. *Cell Tissue Res* 2001; 304:377–382.
40. Strakova Z, Copland JA, Lolait SJ, Soloff MS. ERK2 mediates oxytocin-stimulated PGE₂ synthesis. *Am J Physiol* 1998; 274:E634–E641.