

## **Forty-Ninth Supplement to the American Ornithologists' Union Check-List of North American Birds**

Authors: Banks, Richard C., Chesser, R. Terry, Cicero, Carla, Dunn, Jon L., Kratter, Andrew W., et al.

Source: *The Auk*, 125(3) : 758-768

Published By: American Ornithological Society

URL: <https://doi.org/10.1525/auk.2008.9708>

---

BioOne Complete ([complete.BioOne.org](https://complete.BioOne.org)) is a full-text database of 200 subscribed and open-access titles in the biological, ecological, and environmental sciences published by nonprofit societies, associations, museums, institutions, and presses.

Your use of this PDF, the BioOne Complete website, and all posted and associated content indicates your acceptance of BioOne's Terms of Use, available at [www.bioone.org/terms-of-use](https://www.bioone.org/terms-of-use).

Usage of BioOne Complete content is strictly limited to personal, educational, and non - commercial use. Commercial inquiries or rights and permissions requests should be directed to the individual publisher as copyright holder.

---

BioOne sees sustainable scholarly publishing as an inherently collaborative enterprise connecting authors, nonprofit publishers, academic institutions, research libraries, and research funders in the common goal of maximizing access to critical research.



## FORTY-NINTH SUPPLEMENT TO THE AMERICAN ORNITHOLOGISTS' UNION CHECK-LIST OF NORTH AMERICAN BIRDS

RICHARD C. BANKS,<sup>1,11,12</sup> R. TERRY CHESSER,<sup>1</sup> CARLA CICERO,<sup>2</sup> JON L. DUNN,<sup>3</sup>  
ANDREW W. KRATTER,<sup>4</sup> IRBY J. LOVETTE,<sup>5</sup> PAMELA C. RASMUSSEN,<sup>6</sup> J. V. REMSEN, JR.,<sup>7</sup>  
JAMES D. RISING,<sup>8</sup> DOUGLAS F. STOTZ,<sup>9</sup> AND KEVIN WINKER<sup>10</sup>

<sup>1</sup>U.S. Geological Survey, Patuxent Wildlife Research Center, National Museum of Natural History, MRC-111,  
P.O. Box 37012, Washington, D.C. 20013, USA;

<sup>2</sup>Museum of Vertebrate Zoology, University of California, 3101 Valley Life Sciences Building, Berkeley,  
California 94720, USA;

<sup>3</sup>Rural Route 2, Box 52R, Bishop, California 93514, USA;

<sup>4</sup>Florida Museum of Natural History, P.O. Box 117800, University of Florida, Gainesville, Florida 32611, USA;

<sup>5</sup>Cornell Laboratory of Ornithology, 159 Sapsucker Woods Road, Ithaca, New York 14850, USA;

<sup>6</sup>Museum and Department of Zoology, Michigan State University, West Circle Drive, East Lansing,  
Michigan 48824, USA;

<sup>7</sup>Museum of Natural Science, Louisiana State University, Foster Hall 119, Baton Rouge, Louisiana 70803, USA;

<sup>8</sup>Department of Zoology and Evolutionary Biology, University of Toronto, 25 Harbord Street,  
Toronto, Ontario M5S 3G5, Canada;

<sup>9</sup>Environmental and Conservation Programs, Field Museum of Natural History, 1400 S. Lake Shore Drive,  
Chicago, Illinois 60605, USA; and

<sup>10</sup>University of Alaska Museum, 907 Yukon Drive, Fairbanks, Alaska 99775, USA

This is the eighth Supplement since publication of the 7th edition of the *Check-list of North American Birds* (American Ornithologists' Union [AOU] 1998). It summarizes decisions made by the AOU's Committee on Classification and Nomenclature—North and Middle America between 1 January and 31 December 2007. The Committee has continued to operate in the manner outlined in the 42nd Supplement (AOU 2000). Kevin Winker became a member of the Committee in 2007. Changes in this Supplement fall into the following categories: (1) one genus (*Creagrus*) and three species (*Creagrus furcatus*, *Phylloscopus proregulus*, and *Turdus philomelos*) are added to the main list (including one transferred from the Appendix) because of new distributional information; (2) one species is removed from the list (*Buteogallus subtilis*) by being merged with another on the list; (3) two species are changed by being split from extralimital species (*Anas zonorhyncha* and *Nonnula frontalis*); (4) six genera are added (*Helicolestes*, *Chroicocephalus*, *Hydrocoloeus*, *Leucophaeus*, *Epinecrophylla*, and *Magumma*) and one is replaced (*Pyrilia*) because of generic splits; (5) two genera (*Lysurus* and *Buarremon*) are lost by merger (with *Arremon*); (6) 18 scientific names are changed by transfer from one genus to another (*Helicolestes hamatus*, *Chroicocephalus*

*philadelphia*, *C. cirrocephalus*, *C. ridibundus*, *Hydrocoloeus minutus*, *Leucophaeus modestus*, *L. atricilla*, *L. pipixcan*, *Pyrilia pyrilia*, *P. haematotis*, *Colaptes rubiginosus*, *C. auricularis*, *Epinecrophylla fulviventris*, *Arremon crassirostris*, *A. brunneinucha*, *A. virenticeps*, *A. torquatus*, and *Magumma parva*); (7) 13 English names are changed (*Phoenicopterus ruber* becomes American Flamingo, two species of *Colibri* become Violetear rather than Violetear, *Goethalsia bella* becomes Pirre [rather than Rufous-cheeked] Hummingbird, *Cnipodectes subbrunneus* becomes Brownish Twistwing rather than Brownish Flycatcher, six species of *Turdus* become Thrush rather than Robin, *Chlorothraupis carmioli* becomes Carmiol's [rather than Olive] Tanager, and Troupial becomes Venezuelan Troupial); (8) distribution statements of four species are changed by splits of extralimital taxa (*Pelecanus occidentalis*, *Phoenicopterus ruber*, *Conopias albivittatus*, and *Icterus icterus*); and (9) one generic name in the Appendix is changed (*Columba goodsoni* becomes *Patagioenas goodsoni*).

Additionally, a new classification and sequence of genera and species is adopted for gulls of the subfamily Larinae. A new sequence is adopted for the species of tinamou (Tinamidae) and for

<sup>11</sup> Authors are members of the Committee on Classification and Nomenclature—North and Middle America, of the American Ornithologists' Union, listed alphabetically after the Chairman.

<sup>12</sup> E-mail: [banksr@si.edu](mailto:banksr@si.edu)

species in the genus *Tangara*. The flamingos (Phoenicopteridae) are moved to a position immediately following the grebes (Podicipedidae). In higher-level classification, a suborder (Eurylaimi) and family (Eurylaimidae) are added to our list for the species *Sapayoa aenigma*, previously *incertae sedis*. The family Furnariidae is divided into three subfamilies. The family Formicariidae is limited to anthrushes of the genera *Formicarius* and (extralimital) *Chamaeza*; the genus *Pittasoma* is transferred to the family Conopophagidae (adding a family to the *Check-list* area), and the genera *Grallaria*, *Hyllopezus*, and *Grallaricula* are placed in a newly recognized family, Grallariidae.

Several of the changes in this Supplement were made so that the North and Middle American and South American (Rensen et al. 2008) lists will conform. The species changed belong to groups that are primarily South American.

Literature that provides the basis for the Committee's decisions is cited at the end of the Supplement, and citations not already in the Literature Cited of the 7th edition (with Supplements) become additions to it. An updated list of the bird species known from the AOU *Check-list* area can be found at [www.AOU.org/checklist/index.php3](http://www.AOU.org/checklist/index.php3).

The following changes to the 7th edition (page numbers refer thereto) and its Supplements result from the Committee's actions:

pp. xvii–liv. Change the number in the title of the list of species to 2,048. Insert the following names in the proper position as indicated by the text of this Supplement:

*Anas zonorhyncha* Eastern Spot-billed Duck. (A)  
*Helicolestes hamatus* Slender-billed Kite.  
*Creagrus furcatus* Swallow-tailed Gull. (A)  
*Pyrilia haematotis* Brown-hooded Parrot.  
*Pyrilia pyrilia* Saffron-headed Parrot.  
*Nonnula frontalis* Gray-cheeked Nunlet.  
*Colaptes rubiginosus* Golden-olive Woodpecker.  
*Colaptes auricularis* Gray-crowned Woodpecker.  
 EURYLAIMIDAE  
 Sclerurinae  
 Furnariinae  
 Dendrocolaptinae  
 CONOPOPHAGIDAE  
 GRALLARIIDAE  
*Epinecrophylla fulviventris* Checker-throated Antwren.  
*Phylloscopus proregulus* Pallas's Leaf-Warbler. (A)  
*Turdus philomelos* Song Thrush. (A)  
*Magumma parva* Anianiau.

Delete the following names:

*Anas poecilorhyncha* Spot-billed Duck. (A)  
*Rostrhamus hamatus* Slender-billed Kite.  
*Buteogallus subtilis* Mangrove Black-Hawk.  
*Pionopsitta pyrilia* Saffron-headed Parrot.  
*Pionopsitta haematotis* Brown-hooded Parrot.  
*Nonnula ruficapilla* Gray-cheeked Nunlet.  
*Piculus rubiginosus* Golden-olive Woodpecker.

*Piculus auricularis* Gray-crowned Woodpecker.  
*Myrmotherula fulviventris* Checker-throated Antwren.  
*Hemignathus parvus* Anianiau.

Change the generic names and position:

*Lysurus crassirostris* to *Arremon crassirostris*  
*Buarremon brunneinucha* to *Arremon brunneinucha*  
*Buarremon virenticeps* to *Arremon virenticeps*  
*Buarremon torquatus* to *Arremon torquatus*

Change the English names of the following species:

*Phoenicopus ruber* to American Flamingo.  
*Colibri delphinae* to Brown Violetear.  
*Colibri thalassinus* to Green Violetear.  
*Goethalsia bella* to Pirre Hummingbird.  
*Cnipodectes subbrunneus* to Brownish Twistwing.  
*Turdus nigrescens* to Sooty Thrush.  
*Turdus infuscatus* to Black Thrush.  
*Turdus plebejus* to Mountain Thrush.  
*Turdus grayi* to Clay-colored Thrush.  
*Turdus nudigenis* to Bare-eyed Thrush.  
*Turdus assimilis* to White-throated Thrush.  
*Chlorothraupis carmioli* to Carmiol's Tanager.  
*Icterus icterus* to Venezuelan Troupial.

Reverse the sequence of:

*Nothocercus bonapartei* Highland Tinamou.  
*Tinamus major* Great Tinamou.

Move PHOENICOPTERIFORMES and *Phoenicopus ruber* to follow PODICIPEDIFORMES.

Replace the listing of the Larinae with the following:

*Creagrus furcatus* Swallow-tailed Gull. (A)  
*Rissa tridactyla* Black-legged Kittiwake.  
*Rissa brevirostris* Red-legged Kittiwake.  
*Pagophila eburnea* Ivory Gull.  
*Xema sabini* Sabine's Gull.  
*Chroicocephalus philadelphia* Bonaparte's Gull.  
*Chroicocephalus cirrocephalus* Gray-hooded Gull. (A)  
*Chroicocephalus ridibundus* Black-headed Gull.  
*Hydrocoloeus minutus* Little Gull.  
*Rhodostethia rosea* Ross's Gull.  
*Leucophaeus modestus* Gray Gull. (A)  
*Leucophaeus atricilla* Laughing Gull.  
*Leucophaeus pipixcan* Franklin's Gull.  
*Larus belcheri* Belcher's Gull. (A)  
*Larus crassirostris* Black-tailed Gull. (A)  
*Larus heermanni* Heermann's Gull.  
*Larus canus* Mew Gull.  
*Larus delawarensis* Ring-billed Gull.  
*Larus occidentalis* Western Gull.  
*Larus livens* Yellow-footed Gull.  
*Larus californicus* California Gull.  
*Larus argentatus* Herring Gull.  
*Larus michahellis* Yellow-legged Gull. (A)  
*Larus thayeri* Thayer's Gull.  
*Larus glaucooides* Iceland Gull.

*Larus fuscus* Lesser Black-backed Gull. (N)  
*Larus schistisagus* Slaty-backed Gull.  
*Larus glaucescens* Glaucous-winged Gull.  
*Larus hyperboreus* Glaucous Gull.  
*Larus marinus* Great Black-backed Gull.  
*Larus dominicanus* Kelp Gull.

Reverse the sequence of:

*Pyrilia pyrilia* Saffron-headed Parrot.  
*Pyrilia haematotis* Brown-hooded Parrot.

Move *Sapayoa aenigma* to follow EURYLAIMIDAE.

Move the three species of *Sclerurus* to follow the newly inserted *Sclerurinae*.

Rearrange the species in *Tangara* to the following sequence:

*Tangara palmeri*  
*Tangara cabanisi*  
*Tangara cucullata*  
*Tangara larvata*  
*Tangara guttata*  
*Tangara fucosa*  
*Tangara dowii*  
*Tangara inornata*  
*Tangara lavinia*  
*Tangara gyrola*  
*Tangara florida*  
*Tangara icterocephala*

p. 1. After the family Tinamidae insert the following: **Notes.**—The sequence of genera and species in this family is based on the phylogeny indicated by the data of Bertelli et al. (2002) and Bertelli and Porzecanski (2004).

Reverse the sequence of the tinamou genera (and included species) *Tinamus* and *Nothocercus*.

p. 9. Insert the Order Phoenicopteriformes and its included account, as modified below, after the account for *Aechmophorus clarkii*, transferring it from p. 54. Under the heading for the order, insert: **Notes.**—To recognize the close relationship to the order Podicipediformes shown by several genetic studies (Van Tuinen et al. 2001, Chubb 2004, Cracraft et al. 2004, Ericson et al. 2006), the Phoenicopteriformes are placed next to the Podicipediformes in the linear sequence of orders. They were formerly considered more closely related to the Ciconiiformes.

p. 17. *Procellaria parkinsoni* is added to the list of species known from the United States because of an accepted record off California. Add the following paragraph to the section on Distribution:

Accidental off central California (one photographed about 18 miles [29 km] off Pt. Reyes, Marin County, 1 October 2005; Stallcup and Preston 2006).

p. 31. *Pelecanus thagus* is recognized as distinct from *P. occidentalis*. No reasons were given for the merger by Peters (1931) or subsequent authors who continued to treat them as conspecific.

In the account for *P. occidentalis*, delete information on the *thagus* group and the words “[*occidentalis* group].” Change Notes to read: Formerly included *P. thagus* Molina, 1782 [Peruvian Pelican], now considered distinct (e.g., Sibley and Monroe 1990, Ridgely and Greenfield 2001) on the basis of much larger size, differences in color of plumage and soft parts (Wetmore 1945), and absence of interbreeding.

p. 40. *Ardea cinerea* is added to lists of bird species known to occur in Canada and in the United States because of records in Newfoundland and the Pribilof Islands. Add the following paragraph to the section on Distribution:

Accidental in Canada (Avalon Peninsula, Newfoundland, 11 October 1996; Renner and Linegar 2007) and Alaska (St. Paul Island, Pribilofs, 1 August 1999 and 1–2 October 2007; Gibson et al., in press); a sight record for Bermuda (7 October 2005; Dobson 2005).

p. 54. Greater Flamingo of the Old World, *Phoenicopterus roseus*, is recognized as a species distinct from American birds, *P. ruber*. No convincing evidence was cited for their merger (AOU 1983).

Change the name of *Phoenicopterus ruber* to American Flamingo.

In the Distribution section of the account for *P. ruber*, delete the phrase “[*ruber* group]” and all information for the *roseus* group. Change the Notes to read: Formerly included *P. roseus* Pallas, 1811 [Greater Flamingo], separated on the basis of differences in color of plumage and bill, and in displays and vocalizations (Sangster 1997).

p. 54. To recognize the apparent close relationship to the order Podicipediformes shown by several genetic studies (Van Tuinen et al. 2001, Chubb 2004, Cracraft et al. 2004, Ericson et al. 2006), the order Phoenicopteriformes is moved ahead in the sequence to follow the grebes and should be moved from p. 54 to p. 9, following the account of *Aechmophorus clarkii*.

p. 69. *Anas zonorhyncha* is treated as a separate species from *Anas poecilorhyncha*. Replace the account of the latter with the following:

*Anas zonorhyncha* Swinhoe. Eastern Spot-billed Duck.

*Anas zonorhyncha* Swinhoe, 1866, Ibis 2 (sec. ser.):394. (Ningpo, China.)

**Habitat.**—Small streams and ponds.

**Distribution.**—Breeds from Transbaikalia and the Amur River south through the eastern half of China and winters in southeastern China.

Casual in Alaska in the Aleutians (Attu, Adak; photographs; Gibson and Byrd 2007) and on Kodiak Island (specimen; Trapp and MacIntosh 1978).

**Notes.**—Formerly considered conspecific with *A. poecilorhyncha* J. R. Forster, 1781 [Indian Spot-billed Duck] but separated by Leader (2006) on the basis of sympatric breeding at Hong Kong in southern China.

pp. 89, 90. The genus *Helicolestes*, now in the synonymy of *Rostrhamus*, is restored for the species *hamatus*, because most of the evidence for merging the genera (Amadon 1964) is now suspected to be the result of convergence. Remove the citation for *Helicolestes* from *Rostrhamus* and insert it at the top of p. 90 under the heading:

Genus **HELICOLESTES** Bangs and Penard

**Notes.**—Formerly merged with the genus *Rostrhamus*, following Amadon (1964), but now treated as a separate genus because of lack of evidence of relationship.

Change *Rostrhamus hamatus* (Temminck) to *Helicolestes hamatus* (Temminck). At the end of that species account, add:

**Notes.**—Formerly treated in the genus *Rostrhamus*.

pp. 97–98. *Buteogallus subtilis* is treated as a subspecies of *B. anthracinus*. Add Mangrove Forest to the Habitat section of the *B. anthracinus* account.

Modify the Distribution section of *B. anthracinus* as modified in the previous Supplement (Banks et al. 2007) by inserting “[*anthracinus* group]” after the words “Resident” and “accidental,” and by adding “[*subtilis* group]” and the distribution statement now under *B. subtilis* to the end of the first paragraph. Change the Notes to read: **Notes.**—Formerly known as Black Hawk. Groups: *B. anthracinus* [Common Black-Hawk] and *B. subtilis* (Thayer and Bangs, 1905) [Mangrove Black-Hawk]. An analysis of morphological characters (Clark 2007) has shown that *B. subtilis* is better treated as a subspecies of *B. anthracinus* than as a separate species. See notes after *B. gundlachi*.

Delete the account for *Buteogallus subtilis*.

p. 184 et seq. Pons et al. (2005) proposed a genus-level reclassification of the subfamily Larinae on the basis of studies of mitochondrial DNA; they found that the existing broadly defined genus *Larus* was paraphyletic. Their classification included the splitting of the present genus *Larus* into four genera and the merging of *Rhodostethia* into *Hydrocoloeus*. We accept that classification in principle but disagree with the merging of *Rhodostethia* into *Hydrocoloeus*. The phylogeny of Pons et al. (2005, fig. 1) also suggests that many relationships within the subfamily are best represented by a new linear arrangement of genera and species. We have accepted this new linear sequence except in the case of the white-headed gull group in *Larus*. Support for the perceived relationships of most species in this group is poor, and we prefer to retain the sequence of species currently in use until their relationships are better resolved.

Under the heading Subfamily LARINAE: Gulls, insert **Notes.**—The recognition and sequence of genera largely follows that of Pons et al. (2005).

Remove the citations for *Hydrocoloeus* and *Chroicocephalus* from the synonymy of *Larus*; these will be used later for newly recognized genera. Rearrange the genera and the accounts for the

species in Larinae (from the 7th edition and Supplements) as follows, with new generic names, headings, and notes inserted as indicated. The first species was previously in the Appendix and is newly placed on the main list.

Genus **CREAGRUS** Bonaparte

*Creagrus* Bonaparte, 1854, Naumannia 1854, p. 213. Type, by original designation, *Larus furcatus* Néboux.

*Creagrus furcatus* (Néboux). Swallow-tailed Gull.

*Larus furcatus* Néboux, 1846, Voy. Vénus, Atlas, Zool., Ois., pl. 10. (rade de Monterey, Haute-Californie [error = Galapagos Islands].)

**Habitat.**—Breeds on rocky islands, nesting on cliffs, crevices and caves; otherwise pelagic (feeds at night).

**Distribution.**—Breeds in the Galapagos Islands and Malpelo Island, Colombia.

**Ranges** along the Pacific coast of South America south to central Peru and, uncommonly, to central Chile. Casual well off the coast of Panama and Costa Rica (recent records documented with photos; North American Birds 60:452–454, 2006). Also reported at Osa Peninsula, Costa Rica, 14 March 2003 (North American Birds 57:415, 2003). Accidental in California (Monterey Bay, 6–8 June 1985; photo; Heindel and Garrett 1995, and west of the Farallon Islands, 3 March 1996, McCaskie and San Miguel 1999).

Genus **RISSA**

*Rissa tridactyla* Black-legged Kittiwake

*Rissa brevirostris* Red-legged Kittiwake

Genus **PAGOPHILA**

*Pagophila eburnea* Ivory Gull

Genus **XEMA**

*Xema sabini* Sabine's Gull

Genus **CHROICOCEPHALUS** Eyton, 1836

Insert the citation for this name now in the synonymy of *Larus*.

Insert: **Notes.**—Formerly included in *Larus* but separated on the basis of genetic data (Pons et al. 2005) that indicate that that genus would be paraphyletic if the following species were included.

*Chroicocephalus philadelphia* Bonaparte's Gull

*Chroicocephalus cirrocephalus* (Vieillot) Gray-hooded Gull

*Chroicocephalus ridibundus* (Linnaeus) Black-headed Gull

Genus **HYDROCOLOEUS** Kaup, 1829

Insert the citation for this name now in the synonymy of *Larus*.

Insert: **Notes.**—Formerly included in *Larus* but separated on the basis of genetic data (Pons et al. 2005) that indicate that that genus would be paraphyletic if the following species were included.

*Hydrocoloeus minutus* (Pallas) Little Gull

Genus **RHODOSTETHIA**

**Notes.**—Merged with *Hydrocoloeus* by Pons et al. (2005).

*Rhodostethia rosea* Ross's Gull

Genus **LEUCOPHAEUS** Bruch, 1853

*Leucophaeus* Bruch, 1853, Journ. für Ornithol. 1853, p. 108. Type by original designation, *Larus scoresbii* Traill.

**Notes.**—Formerly included in *Larus* but separated on the basis of genetic data (Pons et al. 2005) that indicate that that genus would be paraphyletic if the following species were included.

*Leucophaeus modestus* (Tschudi) Gray Gull  
*Leucophaeus atricilla* (Linnaeus) Laughing Gull  
*Leucophaeus pipixcan* (Wagler) Franklin's Gull

Genus **LARUS**

*Larus belcheri* Belcher's Gull  
*Larus crassirostris* Black-tailed Gull  
*Larus heermanni* Heermann's Gull  
*Larus canus* Mew Gull  
*Larus delawarensis* Ring-billed Gull  
*Larus occidentalis* Western Gull  
*Larus livens* Yellow-footed Gull  
*Larus californicus* California Gull  
*Larus argentatus* Herring Gull

Insert at beginning of notes: Includes the North American *L. smithsonianus* Coues, 1862, separated as a species by Pons et al. (2005) and the Siberian *L. vegae* Palmen, 1887, separated as a distinct species by Crochet et al. (2002).

*Larus michahellis* Yellow-legged Gull  
*Larus thayeri* Thayer's Gull  
*Larus glaucooides* Iceland Gull  
*Larus fuscus* Lesser Black-backed Gull  
*Larus schistisagus* Slaty-backed Gull  
*Larus glaucescens* Glaucous-winged Gull  
*Larus hyperboreus* Glaucous Gull  
*Larus marinus* Great Black-backed Gull  
*Larus dominicanus* Kelp Gull

p. 194. *Creagrus furcatus* is moved from the Appendix to the main list. A report of a bird in Monterey Bay in 1985 was rejected by the California Bird Records Committee (CBRC) on the basis of uncertain origin of the bird (DeBenedictis 1996). More recent reports resulted in reevaluation of that report, and it is now accepted by the CBRC and the American Birding Association Checklist Committee (Rottenborn and Morlan 2000, Pranty et al. 2007). It would have followed *Xema sabini* in the 7th edition; its proper position is indicated in the listing above.

pp. 240–241. On the basis of mtDNA sequence data, Ribas et al. (2005) removed several species of South and Middle American parrots from the genus *Pionopsitta* Bonaparte, 1854 and placed them in the genus *Gypopsitta* Bonaparte, 1856 (type species *Psittacus vulturinus* Kuhl, 1820). Among these species was *P. pyrilia* (Bonaparte, 1853), which is the type species of the genus *Pyrilia* Bonaparte, 1856. Most citations for these generic names (e.g., Cory 1918, Peters 1937) indicate that they originate from the same paper, but *Pyrilia* was named earlier in 1856 in a different paper and must be used when the two type species are placed in the same genus.

Replace the heading for the genus *Pionopsitta* with:

Genus **PYRILIA** Bonaparte

*Pyrilia* Bonaparte, 1856 (2 June), Compt. Rend. Acad. Sci. Paris 42:956. Type, by original designation and tautonomy, *Psittacula pyrilia* Bonaparte.

*Gypopsitta* Bonaparte, 1856 (August), Naumannia, Beilage no. 1, Consp. Psitt., genus 25. Type, by monotypy, *Psittacus vulturinus* Wagler = *Psittacus vulturinus* Kuhl.

**Notes.**—Formerly merged with the South American genus *Pionopsitta* Bonaparte, 1854, but separated on the basis of mitochondrial DNA sequence data by Ribas et al. (2005), who removed several species of South and Middle American parrots from *Pionopsitta* and placed them in the genus *Gypopsitta* Bonaparte, 1856, but *Pyrilia* Bonaparte, 1856 has priority.

Replace the name *Pionopsitta pyrilia* (Bonaparte) with *Pyrilia pyrilia* (Bonaparte).

Replace the name *Pionopsitta haematotis* (Sclater and Salvin) with *Pyrilia haematotis* (Sclater and Salvin). Reverse the sequence of those species so that *pyrilia* follows *haematotis*.

p. 286. Remove the hyphen in the English name of *Colibri delphinae*, changing it to Brown Violetear (as in Remsen et al. 2008).

p. 287. Remove the hyphen in the English name of *Colibri thalassinus*, changing it to Green Violetear (as in Remsen et al. 2008). In the Notes, change Mountain Violet-ear to Mountain Violetear.

p. 296. Change the English name of *Goethalsia bella* to Pirre Hummingbird (as in Wetmore 1968, Ridgely 1976, Dickinson 2003, and Remsen et al. 2008). Change notes accordingly.

p. 326. *Nonnula frontalis* is separated from the allopatric *N. ruficapilla*. No evidence was presented for their merger and most classifications (e.g., Sibley and Monroe 1990, Rasmussen and Collar 2002, Dickinson 2003, Remsen et al. 2008) treat them as distinct. The English name is retained for the form in our area.

Change the species name and citation of *N. ruficapilla* to:

***Nonnula frontalis*** (Sclater). Gray-cheeked Nunlet.  
*Malacoptila frontalis* Sclater, 1854, Ann. Mag. Nat. Hist. (2) 13:479. (Nova Grenada = interior of Colombia.)

Change the Distribution by removing the term “[*frontalis* group]” and all mention of the *ruficapilla* group. Change the Notes to: Formerly treated as conspecific with *N. ruficapilla* (Tschudi, 1844) [Rufous-capped Nunlet] of South America. Because most sources treat the two as distinct species (Hilty and Brown 1986, Rasmussen and Collar 2002, Dickinson 2003, Remsen et al. 2008), and no evidence supporting a close relationship has ever been presented, we consider that treating *ruficapilla* and *frontalis* as distinct species is the best course.

p. 343. The species now listed as *Piculus rubiginosus* and *P. auricularis* are transferred to the genus *Colaptes*. Move the accounts of these two species to precede that of *Colaptes punctigula* under the heading of the genus *Colaptes*, under the names: ***Colaptes rubiginosus*** (Swainson). Golden-olive Woodpecker. ***Colaptes auricularis*** (Salvin and Godman). Gray-crowned Woodpecker.

In the Notes of each species, change the generic name or abbreviation to *Colaptes* or *C.*, and add the statement: Formerly placed in the genus *Piculus*, but studies of morphological and genetic characters (Benz et al. 2006, Moore et al. 2006) indicate that they are members of *Colaptes*.

To the synonymy of the genus *Colaptes*, after the citation for *Chrysoptilus*, insert:

*Chloronerpes* Swainson, 1837, *Classif. Birds*, 2, p. 307. Type by subsequent designation (G. R. Gray, 1840), *C. rubiginosus* Swainson = *Picus rubiginosus* Swainson.

p. 347. The species *Sapayoa aenigma* has been shown by DNA sequence data to be more closely related to Old World suboscines than to any New World group (Fjeldså et al. 2003, Chesser 2004), and to be embedded in the Old World broadbill family Eurylaimidae (Irestedt et al. 2006, Moyle et al. 2006). After the heading for the Order Passeriformes, insert headings:

Suborder EURYLAIMI: Broadbills, Asities, and Pittas

Family EURYLAIMIDAE: Broadbills

Move the heading for the genus *Sapayoa* and the account for *Sapayoa aenigma* from p. 416 to follow this newly added family.

p. 347 et seq. Independent genetic data sets (Irestedt et al. 2002, 2006; Fjeldså et al. 2003; Chesser 2004) indicate that the family Furnariidae (including the former Dendrocolaptidae, merged into Furnariidae in an earlier Supplement [Banks et al. 2006]) should be divided into three subfamilies: (1) Sclerurinae, containing the genus *Sclerurus* and the South American genus *Geositta*; (2) Furnariinae, containing the remaining genera in the Furnariidae; and (3) Dendrocolaptinae, including the genera in the former family Dendrocolaptidae. The sequence of the genera in the two latter subfamilies does not change.

Change the heading Family FURNARIIDAE: Ovenbirds to Family FURNARIIDAE: Ovenbirds, Woodcreepers, and Leaf-tossers.

Delete the Notes and insert a center heading:

Subfamily SCLERURINAE: Leaf-tossers and Miners

Move the accounts of the genus *Sclerurus* and the included species from pages 353–354 to follow this new subfamily heading.

After the account of *Sclerurus guatemalensis*, insert a center heading:

Subfamily FURNARIINAE: Ovenbirds

Following the account for *Lochmias nematura* on p. 354, insert a center heading:

Subfamily DENDROCOLAPTINAE: Woodcreepers

pp. 364–365. Isler et al. (2006) found that the genus *Myrmotherula* is not monophyletic and named a new genus, *Epinecrophylla*, for the stipple-throated species, to include the species *fulviventris* of our area and seven species endemic to South America.

Delete the Notes under the genus *Myrmotherula*. Before the heading of the genus *Herpsilochmus* on p. 365, insert the heading:

Genus **EPINECROPHYLLA** Isler and Brumfield

*Epinecrophylla* Isler and Brumfield, 2006, in Isler et al., *Proc. Biol. Soc. Wash.* 116: 523. Type species *Formicivora haematonota* Sclater, 1857.

Insert the account for *Myrmotherula fulviventris*, moved from p. 364, with the heading:

***Epinecrophylla fulviventris*** (Lawrence). Checker-throated Antwren.

Add the following: **Notes**.—Formerly placed in the genus *Myrmotherula*, but separated on the basis of genetic data and morphological, vocal, and ecological characters (Isler et al. 2006).

pp. 370–371. Strong genetic evidence (Irestedt et al. 2002; Chesser 2004; Rice 2005a, b) indicates that the family Formicariidae should be limited to anthrushes of the genera *Formicarius* and (extralimital) *Chamaeza*. The genus *Pittasoma* is closely related to the South American *Conopophaga* and belongs with it in the family Conopophagidae. The genera *Grallaria*, *Hylopezus*, and *Grallaricula* are now placed in a newly recognized family Grallariidae. Both of the latter two families, new to our list, were established by Sclater and Salvin (1873).

Remove the words “and Antpittas” from the heading of the family Formicariidae.

After the account of *Formicarius rufipectus* and before the genus *Pittasoma*, insert:

Family CONOPOPHAGIDAE: Gnateaters.

After the citation of the genus *Pittasoma*, add: **Notes**.—Formerly treated as part of the family Formicariidae, but now moved to the Conopophagidae to reflect relationships with the genus *Conopophaga* (Krabbe and Schulenberg 2003, Rice 2005a).

After the account of *Pittasoma michleri* and before the genus *Grallaria*, insert:

Family GRALLARIIDAE: Antpittas.

Insert: **Notes**.—Members of this family were previously included in the family Formicariidae but are placed in their own family because genetic data (Irestedt et al. 2002) indicate that their inclusion in the Formicariidae would make it non-monophyletic.

p. 383. Change the English name of *Cnipodectes subbrunneus* to Brownish Twistwing, as in most recent South American works, e.g., Remsen et al. 2008. Change the Notes to read: Formerly known as Brownish Flycatcher.

p. 409. *Conopias parvus* is considered a species distinct from *C. albobittatus*. Remove the term “[albobittatus group]” and all mention of the *parvus* group from the section on Distribution. Change the Notes to read: **Notes**.—Formerly considered conspecific with *C. parvus* (Pelzeln, 1868) [Yellow-throated Flycatcher] of South America, but separated because of vocal differences (Ridgely and Greenfield 2001, Fitzpatrick 2004). Sometimes placed in the genus *Coryphotriccus*.

p. 414. A new record of *Tyrannus caudifasciatus* in Florida places the species back on the list of birds known to occur in the

United States, from which it was removed by Banks et al. (2002). Replace the second paragraph of the Distribution section with:

Accidental in Florida (Key West, Monroe County, 8–27 March 2007; *North American Birds* 61:432, 2007). Analysis of photos (J. S. Greenlaw *in litt.*) indicates that the subspecies was likely *caudifasciatus* or perhaps *caymanensis*. All prior reports from south Florida are considered questionable (Smith et al. 2000). A sight report for the central Bahamas (Long Island).

Delete last two sentences in notes.

p. 416. Delete the word “seven” from the first sentence and the entire second sentence in the Notes under the heading *Genera INCERTAE SEDIS*.

p. 463. The page number in the citation of the genus *Poecile* should be 114, not 92.

p. 490. *Phylloscopus proregulus* is added to the main list on the basis of a new distributional record. Following the account of *P. fuscatus*, before that of *P. inornatus* (added to the list by Banks et al. 2002), insert the following:

*Phylloscopus proregulus* (Pallas). Pallas’s Leaf-Warbler.

*Motacilla Proregulus* Pallas, 1811, Zoographia Rosso-Asiat. 1:490. (Ingoda River, southern Transbaikalia.)

**Habitat.**—Breeds in mature coniferous and mixed forest, usually with dense scrub undergrowth.

**Distribution.**—Breeds from southwestern Siberia east to Transbaikalia, northern Mongolia, northern Manchuria, Amurland, Ussuriland, and Sakhalin.

**Winters** in southeastern China and northern Indochina. Frequent in fall migration to Scandinavia and northwest Europe, particularly the United Kingdom, and casually to the Mediterranean region and Iceland.

Accidental in Alaska (Gambell, St. Lawrence Island, 25 September 2006; photos; Lehman and Rosenberg 2007).

**Notes.**—We follow Alström and Olsson (1990) in treating this species as monotypic.

p. 508. Song Thrush (*Turdus philomelos*) is added to the main list because of new distributional records. Following the account for *Turdus iliacus*, insert the following:

*Turdus philomelos* Brehm. Song Thrush.

*Turdus philomelos* Brehm, 1831, Handb. Naturgesch. Vög. Deutschl., p. 382. (Mitteldeutschland.)

**Habitat.**—Breeds in a variety of woodland types.

**Distribution.**—Breeds from British Isles and Europe east across Siberia to Lake Baikal and south to northern Iran.

**Winters** in the Mediterranean Basin and southern Asia with small numbers to North Africa and the Arabian Peninsula.

Wanders to Iceland in late fall.

Accidental in northeastern Greenland (specimen) at Clavering Ø in 1982; (Boertmann 1994) and Canada (11–17 November 2006, Saint-Fulgence, Quebec; Auchu et al. 2007; photos).

pp. 508–512. The English group names of several American species in the genus *Turdus* are changed from Robin to Thrush, to agree with the treatments by Ridgely and Tudor (1989), Sibley and Monroe (1990), Gill and Wright (2006), and Remsen et al. (2008). These changes are as follows:

*Turdus nigrescens* Sooty Robin becomes Sooty Thrush

*Turdus infuscatus* Black Robin becomes Black Thrush

*Turdus plebejus* Mountain Robin becomes Mountain Thrush

*Turdus grayi* Clay-colored Robin becomes Clay-colored Thrush

*Turdus nudigenis* Bare-eyed Robin becomes Bare-eyed Thrush

(We are awaiting proposals for a potential name change for this species because African *T. tephronotus* typically bears the same English name)

*Turdus assimilis* White-throated Robin becomes White-throated Thrush

p. 573. Change the English name of *Chlorothraupis carmioli* to Carmiol’s Tanager (as in Meyer de Schauensee 1970, Dickinson 2003, Remsen et al. 2008), from Olive Tanager to avoid confusion with *C. olivacea*.

pp. 586–589. A gene-based phylogeny (Burns and Naoki 2004) has indicated that relationships in the genus *Tangara* are best expressed by a new linear arrangement of the species. Rearrange the species in our list to the following sequence:

*Tangara palmeri*

*Tangara cabanisi*

*Tangara cucullata*

*Tangara larvata*

*Tangara guttata*

*Tangara fucosa*

*Tangara dowii*

*Tangara inornata*

*Tangara lavinia*

*Tangara gyrola*

*Tangara florida*

*Tangara icterocephala*

Add the following after the citation for the genus *Tangara*:  
**Notes.**—The sequence of species in this genus is based on the phylogeny indicated by the genetic data of Burns and Naoki (2004).

pp. 600–602. The genera *Lysurus* and *Buarremon* are merged into the genus *Arremon* to reflect relationships found by study of mitochondrial and nuclear DNA (Cadena et al. 2007).

Delete the headings for the genera *Lysurus* and *Buarremon*; move their citations to follow the citation for the genus *Arremon*. Move the account for *Lysurus crassirostris* to follow that of *Arremon aurantirostris*, and the accounts of the three species of *Buarremon* to follow that of *crassirostris*, with the new headings:

*Arremon crassirostris* (Cassin)

*Arremon brunneinucha* (Lafresnaye)

*Arremon virenticeps* (Bonaparte)

*Arremon torquatus* (Lafresnaye and d’Orbigny)

In the Notes under *A. crassirostris*, change the generic name *Lysurus* to *Arremon* and the initial *L.* to *A.*; add: Formerly treated in the genus *Lysurus* but merged into *Arremon* to reflect relationships found by Cadena et al. (2007).



In the Notes under those species formerly in *Buarremon* (creating a new Note for *A. virenticeps*) insert: Formerly treated in the genus *Buarremon*. Cadena et al. (2007) found that genetic data indicate that *Buarremon* as traditionally defined is paraphyletic with respect to *Arremon* and also probably *Lysurus*. Change the generic abbreviations from *B.* to *A.*

p. 607. Remove the parentheses around the names of the authors of *Aimophila humeralis* and *A. sumichrasti*. The Code, Article 51.3.1 (International Commission on Zoological Nomenclature 1999) states that “Parentheses are not used when the species-group name was originally combined with an incorrect spelling or an emendation of the generic name. . . .”

p. 623. The page numbers in the citation for *Zonotrichia leucophrys* should be 403, 426, not 340.

p. 652. The species *Icterus icterus* is divided into three species; the two populations separated, currently called groups, are extralimital to our area.

Change the English name of *Icterus icterus* (Linnaeus) to Venezuelan Troupial. From the Distribution, remove the phrases “[*icterus* group]” and the sections on the *croconotus* and *jamacaii* groups. Change Notes to: Formerly included two South American populations now separated as the species *Icterus croconotus* (Wagler, 1829) [Orange-backed Troupial] and *I. jamacaii* (Gmelin, 1788) [Campo Troupial] on the basis of limited sympatry without signs of interbreeding between *jamacaii* and *croconotus* (Pacheco and Olmos 2006) and pronounced vocal differences among the three (Jaramillo and Burke 1999, Ridgely and Greenfield 2001, Hilty 2002). Retain the last sentence.

pp. 674–675. The species listed as *Hemignathus parvus* is transferred to the monotypic genus *Magumma* on the basis of studies of mtDNA (Tarr and Fleischer 1993, Fleischer et al. 2001) and morphology (Conant et al. 1998, Pratt 2005). After the account for *Hemignathus munroi* and before the genus *Oreomystis*, insert:

Genus **MAGUMMA** Mathews

*Rothschildia* Perkins in Wilson and Evans, 1899, Aves Hawaiiensis, p. xxi. Type, by monotypy, *Himatione parva* Stejneger.

*Magumma* Mathews, 1925, Bull. Brit. Ornithol. Club 45: 93. New name for *Rothschildia* Perkins, preoccupied.

**Notes.**—Formerly included in *Hemignathus* Lichtenstein, but separated on the basis of genetic and morphological differences (Tarr and Fleischer 1993, Fleischer et al. 2001, Pratt 2005).

Insert the account of *Hemignathus parvus* (p. 674) with the heading:

*Magumma parva* (Stejneger). Anianiau.

The account remains the same, but change Notes to: Formerly included in the genus *Hemignathus*, but see above.

p. 692. Delete the account for *Creagrus furcatus*, now moved to the main list.

p. 693. Change *Columba goodsoni* Hartert to *Patagioenas goodsoni* (Hartert).

p. 705 et seq. In the list of French names for North American Birds, make the following changes:

Insert the following names in the proper position as indicated by the text of this Supplement:

<i>Anas zonorhyncha</i>	Canard de Chine
<i>Helicolestes hamatus</i>	
<i>Creagrus furcatus</i>	
<i>Pyrilia haematotis</i>	
<i>Pyrilia pyrilia</i>	
<i>Nonnula frontalis</i>	Barbacou à joues grises
<i>Colaptes rubiginosus</i>	
<i>Colaptes auricularis</i>	
EURYLAIMIDAE	
CONOPOPHAGIDAE	
GRALLARIIDAE	
<i>Epinecrophylla fulviventris</i>	
<i>Phylloscopus proregulus</i>	Pouillot de Pallas
<i>Turdus philomelos</i>	Grive musicienne
<i>Magumma parva</i>	Anianiau de Kauai

Delete the following names:

*Anas poecilorhyncha*  
*Rostrhamus hamatus*  
*Buteogallus subtilis*  
*Pionopsitta pyrilia*  
*Pionopsitta haematotis*  
*Nonnula ruficapilla*  
*Piculus rubiginosus*  
*Piculus auricularis*  
*Myrmotherula fulviventris*  
*Hemignathus parvus*

Change the French names of the following species:

<i>Phoenicopterus ruber</i>	Flamant des Caraïbes
<i>Geotrygon veraguensis</i>	Colombe de Veraguas
<i>Aratinga nana</i>	Conure aztèque
<i>Coccyzus pluvialis</i>	Tacco de pluie
<i>Coccyzus ruficularis</i>	Tacco cabrite
<i>Threnetes ruckeri</i>	Ermite de Rücker
<i>Anthracothorax veraguensis</i>	Mango de Veraguas
<i>Melanerpes hoffmannii</i>	Pic de Hoffmann
<i>Pittasoma michleri</i>	Pittasome à tête noire
<i>Baoelophus ridgwayi</i>	Mésange des genévriers
<i>Saltator grossus</i>	Saltator ardoisé
<i>Pterodroma defilippiana</i>	Pétrel de De Filippi

Reverse the sequence of *Tinamus major* and *Nothocercus bonapartei*.

Change the generic names and position:

*Lysurus crassirostris* to *Arremon crassirostris*  
*Buarremon brunneinucha* to *Arremon brunneinucha*  
*Buarremon virenticeps* to *Arremon virenticeps*  
*Buarremon torquatus* to *Arremon torquatus*

Move PHOENICOPTERIDAE and *Phoenicopterus ruber* to follow PODICIPEDIDAE.

Replace the listing of the Laridae from *Larus atricilla* to *Pagophila eburnea* with the following, with no change in French names:

*Creagrus furcatus*  
*Rissa tridactyla*  
*Rissa brevirostris*  
*Pagophila eburnea*  
*Xema sabini*  
*Chroicocephalus philadelphia*  
*Chroicocephalus cirrocephalus*  
*Chroicocephalus ridibundus*  
*Hydrocoloeus minutus*  
*Rhodostethia rosea*  
*Leucophaeus modestus*  
*Leucophaeus atricilla*  
*Leucophaeus pipixcan*  
*Larus belcheri*  
*Larus crassirostris*  
*Larus heermanni*  
*Larus canus*  
*Larus delawarensis*  
*Larus occidentalis*  
*Larus livens*  
*Larus californicus*  
*Larus argentatus*  
*Larus michahellis*  
*Larus thayeri*  
*Larus glaucoides*  
*Larus fuscus*  
*Larus schistisagus*  
*Larus glaucescens*  
*Larus hyperboreus*  
*Larus marinus*  
*Larus dominicanus*

Move *Sapayoa aenigma* to follow the newly inserted EURYLAIMIDAE.

Move the three species of *Sclerurus* to the beginning of the FURNARIIDAE.

Rearrange the species in *Tangara* to the following sequence:

*Tangara palmeri*  
*Tangara cabanisi*  
*Tangara cucullata*  
*Tangara larvata*  
*Tangara guttata*  
*Tangara fucosa*  
*Tangara dowii*  
*Tangara inornata*  
*Tangara lavinia*  
*Tangara gyrola*  
*Tangara florida*  
*Tangara icterocephala*

Delete *Creagrus furcatus* from its listing in the appendix.

Change *Columba goodsoni* to *Patagioenas goodsoni*.

Taxonomic proposals considered but not accepted by the Committee include: recognition of *Chondrohierax uncinatus wilsonii* of Cuba as a species (Johnson et al. 2007); resurrection of the genus *Rupornis* for the Roadside Hawk (Riesing et al. 2003); the merger of *Rhodostethia* into *Hydrocoloeus* (Pons et al. 2005); the separation of *Larus smithsonianus* and *L. vegae* from *L. argentatus* (Crochet et al. 2002, Pons et al. 2005, Olson and Banks 2007); the separation of *Pionopsitta* (now *Pyrilia*) *h. coccinillaris* from *P. haematotis* (Ribas et al. 2005); the transfer of *Veniliornis fumigatus* to *Picooides* (Moore et al. 2006), held in anticipation of further changes in *Picooides*; the division of *Icterus spurius* into two species (Kiere et al. 2007); and the transfer of the New World species of *Carpodacus* to the genus *Burrica* (Arnaiz-Villena et al. 2007). Proposals to change the English names of *Gallinula chloropus*, *Brotogeris versicolurus*, and *Microbates cinereiventris* were rejected. Finally, a broad proposal to alter the hyphenation of English names was not accepted for reasons outlined elsewhere (Auk 124:1472, 2007).

#### ACKNOWLEDGMENTS

Normand David serves as the Committee's advisor for classical languages in relation to scientific names, and Michel Gosselin is the authority for French names. We also thank K. J. Burns, C. D. Cadena, T. Donegan, R. Fraga, D. D. Gibson, M. J. Iliff, A. Jaramillo, M. Mendenhall, A. P. Peterson, H. D. Pratt, F. G. Stiles, J. Tobias, and K. J. Zimmer for assistance, suggestions, and comments.

#### LITERATURE CITED

- ALSTRÖM, P., AND U. OLSSON. 1990. Taxonomy of the *Phylloscopus proregulus* complex. *Bulletin of the British Ornithologists' Club* 110:38–43.
- AMADON, D. 1964. Taxonomic notes on birds of prey. *American Museum Novitates* 2166:1–24.
- AMERICAN ORNITHOLOGISTS' UNION. 1983. Check-list of North American Birds, 6th ed. American Ornithologists' Union, Washington, D.C.
- AMERICAN ORNITHOLOGISTS' UNION. 1998. Check-list of North American Birds, 7th ed. American Ornithologists' Union, Washington, D.C.
- AMERICAN ORNITHOLOGISTS' UNION. 2000. Forty-second supplement to the American Ornithologists' Union *Check-list of North American Birds*. *Auk* 117:847–858.
- ARNAIZ-VILLENA, A., J. MOSCOSO, V. RUIZ-DEL-VALLE, J. GONZALEZ, R. REGUERA, M. WINK, AND J. I. SERRANO-VELA. 2007. Bayesian phylogeny of Fringillinae birds: Status of the singular African Oriole Finch *Linurgus olivaceus* and evolution and heterogeneity of the genus *Carpodacus*. *Acta Zoologica Sinica* 53:826–834.
- AUCHU, C., C. GIRARD, AND G. SAVARD. 2007. First record of Song Thrush (*Turdus philomelos*) in North America. *North American Birds* 61:166–168.

- BANKS, R. C., R. T. CHESSER, C. CICERO, J. L. DUNN, A. W. KRATTER, I. J. LOVETTE, P. C. RASMUSSEN, J. V. REMSEN, JR., J. D. RISING, AND D. F. STOTZ. 2007. Forty-eighth supplement to the American Ornithologists' Union *Check-list of North American Birds*. *Auk* 124:1109–1115.
- BANKS, R. C., C. CICERO, J. L. DUNN, A. W. KRATTER, P. C. RASMUSSEN, J. V. REMSEN, JR., J. D. RISING, AND D. F. STOTZ. 2002. Forty-third supplement to the American Ornithologists' Union *Check-list of North American Birds*. *Auk* 119:897–906.
- BANKS, R. C., C. CICERO, J. L. DUNN, A. W. KRATTER, P. C. RASMUSSEN, J. V. REMSEN, JR., J. D. RISING, AND D. F. STOTZ. 2006. Forty-seventh supplement to the American Ornithologists' Union *Check-list of North American Birds*. *Auk* 123:926–936.
- BENZ, B. W., M. B. ROBBINS, AND A. T. PETERSON. 2006. Evolutionary history of woodpeckers and allies (Aves: Picidae): Placing key taxa on the phylogenetic tree. *Molecular Phylogenetics and Evolution* 40:389–399.
- BERTELLI, S., N. P. GIANNINI, AND P. A. GOLOBOFF. 2002. A phylogeny of the tinamous (Aves: Palaeognathiformes) based on integumentary characters. *Systematic Biology* 51:959–979.
- BERTELLI, S., AND A. L. PORZECANSKI. 2004. Tinamou (Tinamidae) systematics: A preliminary combined analysis of morphology and molecules. *Ornithologia Neotropical* 15 (Supplement):293–299.
- BOERTMANN, D. 1994. An annotated checklist to the birds of Greenland. *Meddelelser om Grønland, Bioscience*, no. 38.
- BURNS, K. J., AND K. NAOKI. 2004. Molecular phylogenetics and biogeography of Neotropical tanagers in the genus *Tangara*. *Molecular Phylogenetics and Evolution* 32:838–854.
- CADENA, C. D., J. KLICKA, AND R. E. RICKLEFS. 2007. Evolutionary differentiation in the Neotropical montane region: Molecular phylogenetics and phylogeography of *Buarremon* brush-finches (Aves, Emberizidae). *Molecular Phylogenetics and Evolution* 44:993–1016.
- CHESSER, R. T. 2004. Molecular systematics of New World suboscine birds. *Molecular Phylogenetics and Evolution* 32:11–24.
- CHUBB, A. L. 2004. New nuclear evidence for the oldest divergence among neognath birds: The phylogenetic utility of ZENK (i). *Molecular Phylogenetics and Evolution* 30:140–151.
- CLARK, W. S. 2007. Taxonomic status and distribution of Mangrove Black Hawk *Buteogallus (anthracinus) subtilis*. *Bulletin of the British Ornithologists' Club* 127:110–117.
- CONANT, S., H. D. PRATT, AND R. J. SHALLENBERGER. 1998. Reflections on a 1975 expedition to the lost world of the Alaka'i and other notes on the natural history, systematics, and conservation of Kaua'i birds. *Wilson Bulletin* 110:1–22.
- CORY, C. B. 1918. Catalogue of birds of the Americas. Field Museum of Natural History Publications, Zoological Series, vol. 13, pt. 2, no. 1.
- CRACRAFT, J., F. K. BARKER, M. BRAUN, J. HARSHMAN, G. J. DYKE, J. FEINSTEIN, S. STANLEY, A. CIBOIS, P. SCHIKLER, P. BERESFORD, AND OTHERS. 2004. Phylogenetic relationships among modern birds (Neornithes): Toward an avian tree of life. Pages 468–489 in *Assembling the Tree of Life* (J. Cracraft and M. J. Donoghue, Eds.). Oxford University Press, Oxford, United Kingdom.
- CROCHET, P.-A., J.-D. LEBRETON, AND F. BONHOMME. 2002. Systematics of large white-headed gulls: Patterns of mitochondrial DNA variation in western European taxa. *Auk* 119:603–620.
- DEBENEDICTIS, P. A. 1996. 1995 ABA Checklist report. *Birding* 28:399–405.
- DICKINSON, E. C., ED. 2003. *The Howard and Moore Complete Checklist of the Birds of the World*, 3rd ed. Princeton University Press, Princeton, New Jersey.
- DOBSON, A. 2005. Grey Heron—New to Bermuda. *Bermuda Audubon Society Newsletter* 16, no. 3.
- ERICSON, P. G. P., C. L. ANDERSON, T. BRITTON, A. ELZANOWSKI, U. S. JOHANSSON, M. KÄLLERSJÖ, J. I. OHLSON, T. J. PARSONS, D. ZUCCON, AND G. MAYR. 2006. Diversification of Neoaves: Integration of molecular sequence data and fossils. *Biology Letters* 2:543–547.
- FITZPATRICK, J. W. 2004. Family Tyrannidae (tyrant-flycatchers). Pages 170–462 in *Handbook of the Birds of the World*, vol. 9: Cotingas to Pipits and Wagtails (J. del Hoyo, A. Elliott, and D. Christie, Eds.). Lynx Edicions, Barcelona, Spain.
- FJELDSÅ, J., M. IRESTEDT, AND P. G. P. ERICSON. 2005. Molecular data reveal some major adaptational shifts in the early evolution of the most diverse avian family, the Furnariidae. *Journal of Ornithology* 146:1–13.
- FJELDSÅ, J., D. ZUCCON, M. IRESTEDT, U. S. JOHANSSON, AND P. G. P. ERICSON. 2003. *Sapayoa aenigma*: A New World representative of 'Old World suboscines'. *Proceedings of the Royal Society of London, Series B (Supplement)* 270:S238–S241.
- FLEISCHER, R. C., C. L. TARR, H. F. JAMES, B. SLIKAS, AND C. E. MCINTOSH. 2001. Phylogenetic placement of the Po'ouli, *Melamprosops phaeosoma*, based on mitochondrial DNA sequence and osteological characters. Pages 98–103 in *Evolution, Ecology, Conservation, and Management of Hawaiian Birds: A Vanishing Avifauna* (J. M. Scott, S. Conant, and C. Van Riper III, Eds.). *Studies in Avian Biology*, no. 22.
- GIBSON, D. D., AND G. V. BYRD. 2007. *Birds of the Aleutian Islands, Alaska*. Series in Ornithology, no. 1.
- GIBSON, D. D., S. C. HEINL, AND T. G. TOBISH, JR. in press. Report of the Alaska Checklist Committee, 2003–2007. *Western Birds*.
- GILL, F., AND M. WRIGHT. 2006. *Birds of the World: Recommended English Names*. Princeton University Press, Princeton, New Jersey.
- HEINDEL, M. T., AND K. L. GARRETT. 1995. *Sixteenth Annual Report of the California Bird Records Committee*. *Western Birds* 26:1–33.
- HILTY, S. L. 2002. *Birds of Venezuela*, 2nd ed. Princeton University Press, Princeton, New Jersey.
- HILTY, S. L., AND W. L. BROWN. 1986. *A Guide to the Birds of Colombia*. Princeton University Press, Princeton, New Jersey.
- INTERNATIONAL COMMISSION ON ZOOLOGICAL NOMENCLATURE. 1999. *International Code of Zoological Nomenclature*, 4th ed. International Commission on Zoological Nomenclature, London.
- IRESTEDT, M., J. FJELDSÅ, U. S. JOHANSSON, AND P. G. P. ERICSON. 2002. Systematic relationships and biogeography of the tracheophone suboscines (Aves: Passeriformes). *Molecular Phylogenetics and Evolution* 23:499–512.
- IRESTEDT, M., J. I. OHLSON, D. ZUCCON, M. KÄLLERSJÖ, AND P. G. P. ERICSON. 2006. Nuclear DNA from old collections of avian study skins reveals the evolutionary history of the Old World suboscines (Aves, Passeriformes). *Zoologica Scripta* 35:567–580.

- ISLER, M. I., D. R. LACERDA, P. R. ISLER, S. J. HACKETT, K. V. ROSENBERG, AND R. T. BRUMFIELD. 2006. *Epinecrophylla*, a new genus of antwrens (Aves: Passeriformes: Thamnophilidae). *Proceedings of the Biological Society of Washington* 119:522–527.
- JARAMILLO, A., AND P. BURKE. 1999. *New World Blackbirds: The Icterids*. A. & C. Black, London.
- JOHNSON, J. A., R. THORSTROM, AND D. P. MINDELL. 2007. Systematics and conservation of the Hook-billed Kite including the island taxa from Cuba and Grenada. *Animal Conservation* 10:349–359.
- KIERE, L. M., C. M. HOFMANN, I. E. TRACY, T. W. CRONIN, J. LEIPS, AND K. E. OMLAND. 2007. Using color to define species boundaries: Quantitative analysis in the Orchard Oriole complex supports the recognition of two species. *Condor* 109:692–697.
- KRABBE, N., AND T. S. SCHULENBERG. 2003. Family Formicariidae (ground-antbirds). Pages 682–731 in *Handbook of the Birds of the World*, vol. 8: Broadbills to Tapaculos (J. del Hoyo, A. Elliott, and D. Christie, Eds.). Lynx Edicions, Barcelona, Spain.
- LEADER, P. J. 2006. Sympatric breeding of two Spot-billed Duck *Anas poecilorhyncha* taxa in southern China. *Bulletin of the British Ornithologists' Club* 126:248–252.
- LEHMAN, P. E., AND G. H. ROSENBERG. 2007. First North American record of Pallas's Warbler (*Phylloscopus proregulus*) at Gambell, Alaska. *North American Birds* 61:4–8.
- MCCASKIE, G., AND M. SAN MIGUEL. 1999. Report of the California Bird Records Committee: 1996 records. *Western Birds* 30:57–85.
- MEYER DE SCHAUENSEE, R. 1970. *A Guide to the Birds of South America*. Livingston, Wynnewood, Pennsylvania.
- MOORE, W. S., A. C. WEIBEL, AND A. AGIUS. 2006. Mitochondrial DNA phylogeny of the woodpecker genus *Veniliornis* (Picidae, Picinae) and related genera implies convergent evolution of plumage patterns. *Biological Journal of the Linnean Society* 87: 611–624.
- MOYLE, R. G., R. T. CHESSER, R. O. PRUM, P. SCHIKLER, AND J. CRACRAFT. 2006. Phylogeny and evolutionary history of Old World subsocial birds (Aves: Eurylaimides). *American Museum Novitates* 3544.
- OLSON, S. L., AND R. C. BANKS. 2007. Leptotypification of *Larus smithsonianus* Coues, 1862 (Aves: Laridae). *Proceedings of the Biological Society of Washington* 120:382–386.
- PETERS, J. L. 1931. *Check-list of Birds of the World*, vol. 1. Harvard University Press, Cambridge, Massachusetts.
- PETERS, J. L. 1937. *Check-list of Birds of the World*, vol. 3. Harvard University Press, Cambridge, Massachusetts.
- PONS, J.-M., A. HASSANIN, AND P.-A. CROCHET. 2005. Phylogenetic relationships within the *Laridae* (Charadriiformes: Aves) inferred from mitochondrial markers. *Molecular Phylogenetics and Evolution* 37:686–699.
- PRANTY, B., J. L. DUNN, S. C. HEINL, A. W. KRATTER, P. E. LEHMAN, M. W. LOCKWOOD, B. MACTAVISH, AND K. J. ZIMMER. 2007. Annual Report of the ABA Checklist Committee: 2007. *Birding* 39(6):24–31.
- PRATT, H. D. 2005. *The Hawaiian Honeycreepers: Drepanidinae*. Oxford University Press, Oxford, United Kingdom.
- RASMUSSEN, P. C., AND N. J. COLLAR. 2002. Family Bucconidae (puffbirds). Pages 102–139 in *Handbook of the Birds of the World*, vol. 7: Jacamars to Woodpeckers (J. del Hoyo, A. Elliott, and J. Saragta, Eds.). Lynx Edicions, Barcelona, Spain.
- REMSEN, J. V., JR., C. D. CADENA, A. JARAMILLO, M. NORES, J. F. PACHECO, M. B. ROBBINS, T. S. SCHULENBERG, F. G. STILES, D. F. STOTZ, AND K. J. ZIMMER. 2008. A classification of the bird species of South America. American Ornithologists' Union, Washington, D.C. [ONLINE.] Available at [www.museum.lsu.edu/~Remsen/SACCBaseline.html](http://www.museum.lsu.edu/~Remsen/SACCBaseline.html).
- RENNER, M., AND P. D. LINEGAR. 2007. The first specimen record of Gray Heron (*Ardea cinerea*) for North America. *Wilson Journal of Ornithology* 119:134–136.
- RIBAS, C. C., R. GABAN-LIMA, C. Y. MIYAKI, AND J. CRACRAFT. 2005. Historical biogeography and diversification within the Neotropical parrot genus *Pionopsitta* (Aves: Psittacidae). *Journal of Biogeography* 32:1409–1427.
- RICE, N. H. 2005a. Further evidence for paraphyly of the Formicariidae (Passeriformes). *Condor* 107:910–915.
- RICE, N. H. 2005b. Phylogenetic relationships of antpitta genera (Passeriformes: Formicariidae). *Auk* 122:673–683.
- RIDGELY, R. S. 1976. *A Guide to the Birds of Panama*. Princeton University Press, Princeton, New Jersey.
- RIDGELY, R. S., AND P. J. GREENFIELD. 2001. *The Birds of Ecuador*, vol. 1: Status, Distribution, and Taxonomy. Cornell University Press, Ithaca, New York.
- RIDGELY, R. S., AND G. TUDOR. 1989. *The Birds of South America*, vol. 1: The Oscine Passerines. University of Texas Press, Austin.
- RIESING, M. J., L. KRUCKENHAUSER, A. GAMAU, AND E. HARING. 2003. Molecular phylogeny of the genus *Buteo* (Aves: Accipitridae) based on mitochondrial marker sequences. *Molecular Phylogenetics and Evolution* 27:328–342.
- ROTTENBORN, S., AND J. MORLAN. 2000. Report of the California Bird Records Committee: 1997 Records. *Western Birds* 31:1–37.
- SANGSTER, G. 1997. Species limits in flamingos, with comments on lack of consensus in taxonomy. *Dutch Birding* 19:193–198.
- SCLATER, P. L., AND O. SALVIN. 1873. *Nomenclator Avium Neotropicalium*. J. W. Elliot, London.
- SIBLEY, C. G., AND B. L. MONROE, JR. 1990. *Distribution and Taxonomy of Birds of the World*. Yale University Press, New Haven, Connecticut.
- SMITH, P. W., G. E. WOOLFENDEN, AND A. SPRUNT IV. 2000. The Loggerhead Kingbird in Florida: The evidence revisited. *North American Birds* 54:235–240.
- STALLCUP, R., AND E. W. PRESTON. 2006. First record of Parkinson's Petrel (*Procellaria parkinsoni*) for the continental United States. *North American Birds* 60:166–169.
- TARR, C. L., AND R. C. FLEISCHER. 1993. Mitochondrial-DNA variation and evolutionary relationships in the Amakihi complex. *Auk* 110:825–831.
- TRAPP, J. L., AND R. A. MACINTOSH. 1978. First North American specimen of the Spotbill Duck. *Western Birds* 9:127–128.
- VAN TUINEN, M., D. B. BUTVILL, J. A. W. KIRSCH, AND S. B. HEDGES. 2001. Convergence and divergence in the evolution of aquatic birds. *Proceedings of the Royal Society London, Series B* 268:1345–1350.
- WETMORE, A. 1945. A review of the forms of the Brown Pelican. *Auk* 62:577–586.
- WETMORE, A. 1968. *The Birds of the Republic of Panamá*, part 2. *Smithsonian Miscellaneous Collections*, vol. 150.