

CHOLULA MINUTA, A NEW SPECIES OF MYODOCHINI (LYGAEOIDEA: RHYPAROCHROMIDAE) FROM JAMAICA

Author: Cervantes Peredo, Luis

Source: Florida Entomologist, 87(2) : 173-175

Published By: Florida Entomological Society

URL: [https://doi.org/10.1653/0015-4040\(2004\)087\[0173:CMANSO\]2.0.CO;2](https://doi.org/10.1653/0015-4040(2004)087[0173:CMANSO]2.0.CO;2)

BioOne Complete (complete.BioOne.org) is a full-text database of 200 subscribed and open-access titles in the biological, ecological, and environmental sciences published by nonprofit societies, associations, museums, institutions, and presses.

Your use of this PDF, the BioOne Complete website, and all posted and associated content indicates your acceptance of BioOne's Terms of Use, available at www.bioone.org/terms-of-use.

Usage of BioOne Complete content is strictly limited to personal, educational, and non - commercial use. Commercial inquiries or rights and permissions requests should be directed to the individual publisher as copyright holder.

BioOne sees sustainable scholarly publishing as an inherently collaborative enterprise connecting authors, nonprofit publishers, academic institutions, research libraries, and research funders in the common goal of maximizing access to critical research.

**CHOLULA MINUTA, A NEW SPECIES OF MYODOCHINI
(LYGAEOIDEA: RHYPAROCHROMIDAE) FROM JAMAICA**

LUIS CERVANTES PEREDO

Instituto de Ecología, A.C., Apartado Postal 63 CP 91000, Xalapa, Veracruz, Mexico

E-mail: cervantl@ecologia.edu.mx

ABSTRACT

A new species of *Cholula* (Myodochini) from Jamaica is described. This represents the first record of this genus for the Caribbean. *Cholula minuta* can be differentiated from other species of the genus mainly by its size. It is one of the smallest species described to date, being similar in size only to *C. parvus*, but *C. minuta* is unicolorous, while *C. parvus* has a mixture of black, brown and white coloration.

Key Words: *Cholula*, Lygaeidae, Rhyparochromidae, Jamaica

RESUMEN

Se describe una nueva especie de *Cholula* (Myodochini) de Jamaica. Ésta representa el primer registro de este género para el Caribe. *Cholula minuta* puede diferenciarse de otras especies del género principalmente debido a su tamaño. Es una de las especies más pequeñas descritas hasta ahora, es similar en tamaño a *C. parvus*, pero *C. minuta* es de un solo color, mientras que *C. parvus* es de una coloración mezclada de negro, pardo y blanco.

Translation provided by author.

This paper describes a new species of *Cholula* in order to make the name available for a review of West Indian lygaeids that is in preparation by J. A. Slater and R. Baranowski. The genus *Cholula* includes 12 species of Neotropical distribution. None of the species has been recorded previously from the Caribbean; six species are reported from Mexico (*C. bracteicola* Cervantes & Pacheco, *C. irrorandus* (Distant), *C. lactifera* Brailovsky, *C. lymphata* Brailovsky, *C. maculatus* (Distant), and *C. scapha* Brailovsky), five from Guatemala (*C. bicolor* Distant, *C. irrorandus*, *C. parvus* (Distant), *C. variegata* Distant, and *C. vigena* (Distant)), three from Panama (*C. discoloria* Distant, *C. firmus* (Distant), and *C. vigena*), and one from Honduras (*C. parvus*) (Brailovsky 1981; Cervantes & Pacheco 2003; Distant 1882-1893).

Cholula minuta Cervantes new species
(Fig. 1)

Labium reaching anterior third of abdominal sternite III. Head and anterior pronotal lobe dark ochraceous; posterior pronotal lobe, scutellum, clavus and corium pale ochraceous, with ochraceous punctures. Ventral surface covered with silvery hairs.

Head and anterior pronotal lobe covered with tiny decumbent silvery hairs; eyes and ocelli reddish brown, ocelli located very close to anterior margin of pronotum; antennae pale brown, with joints pale yellow; rostrum pale yellow with tip of

segment IV brown. Pronotal collar and lateral margins of posterior pronotal lobe yellow. Pronotum and scutellum very densely punctuate. Acetabulae creamy yellow; coxae ochraceous; femora, tibiae and tarsi pale yellow, fore femur slightly darker. Pro-, meso-, and metapleura pale ochraceous with posterior margins pale yellow. Clavus with three complete rows of punctures and one incomplete row. Corium with two rows of punctures parallel to claval suture; rest of corium with sparse punctures; membrane translucent. Abdominal venter pale ochraceous.

Head slightly declivent, wider than long. Width across eyes greater than width across anterior angles of pronotum. Tylus longer than juga. Lateral pronotal margins sinuate. Disk of scutellum slightly elevated. Fore femur ventrally with double ranked spines. Evaporative area occupying less than half of metapleuron; peritreme auriculate.

Female

Measurements in mm: Body length 3.6; head length 0.57; width across eyes 0.95; interocular distance 0.62; interocellar distance 0.32; postocular distance 0.02; antennal segments: I 0.22, II 0.45, III 0.37, IV 0.7; rostral segments: I 0.46, II 0.52, III 0.3, IV 0.3; pronotal length 0.87, width across humeral angles 1.32, width across anterior margin 0.72; scutellar length 0.68, width 0.68; hind leg: femur length 0.88, tibia length 0.96, tarsi length: I 0.2, II 0.07, III 0.16.




Fig. 1. Adult of *Cholula minuta* new species. The bar at the right indicates the actual size of the male. The female is slightly larger.

Male (Holotype)

Measurements in mm: Body length 3.4; head length 0.5; width across eyes 0.85; interocular distance 0.55; interocellar distance 0.32; postocular

distance 0.02; antennal segments: I 0.2, II 0.4, III 0.35, IV 0.65; rostral segments: I 0.45, II 0.48, III 0.35, IV 0.3; pronotal length 0.8, width across humeral angles 1.12, width across anterior margin 0.65; scutellar length 0.68, width 0.68; hind leg:

femur length 0.82, tibia length 0.92, tarsi length: I 0.12, II 0.07, III 0.18.

Types. Holotype, 1♂, JAMAICA: Manchester Parish, Mandeville, 24-VIII-1969, R.E. Woodruff, blacklight trap (Florida State Collection of Arthropods). Paratype. 1♀, same locality as holotype; 23-VIII-1960, J. Howard Frank, blacklight trap (R. Baranowski Collection, University of Florida).

DISCUSSION

This species is similar in coloration to *Cholula lactifera* and *C. bracteicola*, but both species are much larger than *C. minuta* sp. nov. In *C. bracteicola* the rostrum reaches abdominal sternite V, and in *C. minuta* it reaches only to anterior third of abdominal sternite III. In *C. lactifera* antennal segment III is pale ochraceous; in *C. bracteicola* the distal fourth of this segment is dark ochraceous, while in *C. minuta* all antennal segments are pale brown. The hemelytral membrane is transparent in *C. minuta* and in *C. bracteicola*, while in *C. lactifera* it has a milky appearance. *Cholula minuta* is one of the smallest species described to date, being similar in size only to *C. parvus*, but *C. minuta* is unicolorous, while *C. parvus* has a mixture of black, brown, and white coloration.

Recent sampling in Mexico has shown that several species of *Cholula* are arboreal, and are

associated with figs, so probably *C. minuta* is also associated with figs in the Caribbean. *Cholula minuta*, as well as other species in the genus, is attracted to light.

ACKNOWLEDGMENTS

I thank Harry Brailovsky (Universidad Nacional Autonoma de Mexico) for his comments on the manuscript. I especially thank James A. Slater for lending me the holotype and for encouraging me to describe this species and continue studying lygaeids. I also thank Richard M. Baranowski for lending me the paratype specimen. Financial support to visit R. Baranowski's collection and publish this paper was provided by a CONACyT grant (34238 V).

REFERENCES CITED

- BRAILOVSKY, H. 1981. Hemiptera-Heteroptera de México XXI. Notas acerca de *Cholula* Distant y descripción de nuevas especies (Lygaeidae: Rhyparochrominae: Myodochini). Folia Entomológica Mexicana 47: 51-68.
- CERVANTES PEREDO, L., AND I. R. PACHECO. 2003. Biology and description of a new species of *Cholula* (Rhyparochromidae: Myodochini) associated with a fig in Mexico. J. New York Entomol. Soc. 111(1): 41-47.
- DISTANT, W. L. 1882-1893. Biologia Centrali-Americana. Heteroptera I. London: 210-215 and 400-404.