

Cryptic Invasion of the Exotic *Bemisia tabaci* Biotype Q Occurred Widespread in Shandong Province of China

Authors: Chu, Dong, Zhang, You Jun, and Wan, Fang Hao

Source: Florida Entomologist, 93(2) : 203-207

Published By: Florida Entomological Society

URL: <https://doi.org/10.1653/024.093.0209>

BioOne Complete (complete.BioOne.org) is a full-text database of 200 subscribed and open-access titles in the biological, ecological, and environmental sciences published by nonprofit societies, associations, museums, institutions, and presses.

Your use of this PDF, the BioOne Complete website, and all posted and associated content indicates your acceptance of BioOne's Terms of Use, available at www.bioone.org/terms-of-use.

Usage of BioOne Complete content is strictly limited to personal, educational, and non - commercial use. Commercial inquiries or rights and permissions requests should be directed to the individual publisher as copyright holder.

BioOne sees sustainable scholarly publishing as an inherently collaborative enterprise connecting authors, nonprofit publishers, academic institutions, research libraries, and research funders in the common goal of maximizing access to critical research.

CRYPTIC INVASION OF THE EXOTIC *BEMISIA TABACI* BIOTYPE Q OCCURRED WIDESPREAD IN SHANDONG PROVINCE OF CHINADONG CHU^{1,2*}, YOU JUN ZHANG³ AND FANG HAO WAN⁴¹High-tech Research Center, Shandong Academy of Agricultural Sciences, and Key Laboratory for Genetic Improvement of Crop Animal and Poultry of Shandong Province, Jinan 250100, China²Key Laboratory of Crop Genetic Improvement and Biotechnology, Huanghuaihai, Ministry of Agriculture, the People's Republic of China, Jinan 250100, China. ³Institute of Vegetables and Flowers, Chinese Academy of Agricultural Sciences, Beijing 100081, China⁴State Key Laboratory for Biology of Plant Diseases and Insect Pests, Institute of Plant Protection, Chinese Academy of Agricultural Sciences, Beijing 100081, China

ABSTRACT

Bemisia tabaci (Gennadius) is an important agricultural pest worldwide. The pest is a species complex composed of numerous biotypes, among which biotypes B and Q are the 2 most invasive and widely distributed. Our previous study found that the ratio of the biotype Q has been increasing and displacement of biotypes B by Q has been occurring on cotton and eggplant in Shandong Province of China during the past several years. To determine whether biotype Q has been increasing on other hosts and possible displacement of biotypes has been occurring in the province as a whole, we further surveyed *B. tabaci* biotypes B and Q on cultivated and wild host species near cotton or eggplant fields in 7 locations of Shandong Province during 2005-2008 with cleavage amplified polymorphic sequence (CAPS) of the *mtCOI* (mitochondrial cytochrome oxidase subunit I) marker. This research showed biotype Q has been increasing on all kinds of host plants and the displacement of biotypes B by Q has been occurring in the province as a whole. The displacement mechanism should be further researched and such knowledge might guide the application of the insecticides or adjustment of the crops to effectively control the pest.

Key Words: *Bemisia tabaci*, biotype Q, mitochondrial cytochrome oxidase marker

RESUMEN

Bemisia tabaci (Gennadius) es una plaga importante a la agricultura mundial. La especie es un complejo de especies compuesta de biotipos numerosos, entre ellos los biotipos B y Q son los 2 mas invasivos y distribuidos ampliamente. Nuestro estudio anterior encontró que la proporción de biotipo Q ha ido incrementando y el desplazamiento del biotipo B por el Q ha ido ocurriendo sobre algodón y berenjena en la Provincia de Shangdon en China durante varios de los años pasados. Para determinar si el biotipo Q se ha ido incrementando sobre otros hospederos y si el posible desplazamiento de biotipos ha ido ocurriendo en toda la provincia, tambien muestreamos los biotipos B y Q de *B. tabaci* sobre especies de hospederos cultivados y silvestres cerca de campos de algodón y berenjena en 7 localidades en la Provincia de Shandong durante 2005-2008 usando el metodo escisión de la secuencia polimórfica amplificado (ESPA) del marcador subunidad I de *mtCOI* (oxidasa citocromo mitocondrial). Esta investigación mostró que el biotipo Q se ha ido incrementando sobre toda clase de plantas hospederos y el desplazamiento de biotipo B por el Q ha ido ocurriendo en toda la provincia. El mecanismo del desplazamiento debe ser investigado y este conocimiento puede guiar la aplicación de insecticidas o un ajuste de los cultivos para controlar la plaga efectivamente.

Bemisia tabaci (Gennadius) is an important agricultural pest worldwide. It damages crops through direct feeding and vectors many plant viruses. The pest has been considered as a species complex that includes many genetic groups that are morphologically indistinguishable.

Some are labeled as biotypes or host races because of differences in host range, geographical distribution, transmission ability of plant virus, and other biological characteristics. Biotype B is believed to originate from the Middle East-Northern Africa and have spread into many

countries or regions over the past 2 decades. Biotype Q may have originated in the Mediterranean countries and circumstantial data shows that this biotype has been introduced into many non-Mediterranean countries or regions during the past several years (Chu et al. 2005; Ueda 2006; Brown et al. 2007).

Bemisia tabaci outbreaks in the mid-1990s in both Southern China and Northern China and subsequent research showed that the whitefly occurring in most of China was biotype B (Wu et al. 2002). In 2003, biotype Q was found in Kunming of Yunnan Province and then subsequently found in Beijing and Henan (Chu et al. 2006). Many populations of *B. tabaci* in Shandong Province, one of the most important agricultural provinces of China, have proved to be biotype B, but biotype Q was discovered in 2006 by use of mitochondrial cytochrome oxidase subunit I (*mtCOI*) sequence (Chu et al. 2007). It is important to monitor the spreading and the density of the biotype Q because it possesses greater resistance to many insecticides than biotype B in many countries (Dennehy et al. 2005; Horowitz et al. 2005). The biotypes of *B. tabaci* on cotton and eggplant in 6 locations within Shandong Prov-

ince was determined with *mtCOI* sequences and biotype B-specific primers (Chu et al. 2010), which showed that the ratio of the biotype Q has been increasing and displacement of biotypes B by Q has been occurring during the past 4 years (2005-2008) on these two crops. To determine whether biotype Q has been increasing on other hosts and possible displacement of biotypes has been occurring in the province as a whole, we further surveyed *B. tabaci* biotypes on cultivated and wild host species near cotton or eggplant fields in 7 locations of Shandong Province during 2005-2008.

MATERIALS AND METHODS

Bemisia tabaci biotype was determined with the cleaved amplified polymorphic sequences (CAPS) of *mtCOI* amplified with new primers (C1-J-2195/R-BQ-2819). Adult whiteflies were collected from different plants including crops and weeds in 7 locations, DeZhou, JiNan, ZiBo, ShouGuang, JiNan, LiaoCheng, LinYi and ZaoZhuang in Shandong Province during 2005-2008 (Table 1). The adults were placed in tubes with 95% ethanol and stored at -20°C. Individual adults were ground and DNA was extracted. The




Fig. 1. Composition changes of *B. tabaci* biotypes B and Q in Shandong Province during 2005-2008 ■, biotype Q; ▨, biotype B.

TABLE 1. DATA ON WHITEFLY COLLECTION AND BIOTYPE DETERMINATION BASED ON THE MARKERS.

Location	Year	Host plant	Total individuals	Number of biotype		Percentage of biotype (%)		
				B	Q	B	Q	
DeZhou	2006	Cotton	31	23	8	74.2	25.8	
		Eggplant	61	34	27	55.7	44.3	
		All host plants	92	57	35	62.0	38.0	
	2007	Tomato	21	7	14	33.3	66.7	
		Cotton	71	17	54	23.9	76.1	
		Eggplant	45	7	38	15.6	84.4	
		Zucchini	37	18	19	48.6	51.4	
		All host plants	174	49	125	28.2	71.8	
	2008	Cotton	27	0	27	0.0	100.0	
		Japanese hop	30	0	30	0.0	100.0	
		Eggplant	30	0	30	0.0	100.0	
		All host plants	87	0	87	0.0	100.0	
ZiBo	2006	Cotton	45	32	13	71.1	28.9	
		Morning glory	20	0	20	0.0	100.0	
		Eggplant	25	23	2	92.0	8.0	
		All host plants	90	55	35	61.1	38.9	
	2007	Cotton	75	2	73	2.7	97.3	
		Eggplant	48	7	41	14.6	85.4	
		All host plants	123	9	114	7.3	92.7	
	2008	Japanese hop	28	0	28	0.0	100.0	
		Cotton	29	0	29	0.0	100.0	
		Eggplant	30	6	24	20.0	80.0	
		All host plants	87	6	81	6.9	93.1	
	ShouGuang	2006	Chinese cabbage	60	23	37	38.3	61.7
Rosebush			17	16	1	94.1	5.9	
Eggplant			30	1	29	3.3	96.7	
All host plants			107	40	67	37.4	62.6	
2007		Tomato	49	0	49	0.0	100.0	
		Cucumber	24	0	24	0.0	100.0	
		Pepper	25	0	25	0.0	100.0	
		Japanese hop	66	1	65	1.5	98.5	
		Cotton	98	13	85	13.3	86.7	
		Pumpkin	62	0	62	0.0	100.0	
		Eggplant	57	3	54	5.3	94.7	
		Sweet pepper	24	0	24	0.0	100.0	
		Chinese cabbage	41	1	40	2.4	97.6	
		All host plants	446	18	428	4.0	96.0	
2008		Japanese hop	29	0	29	0.0	100.0	
		Cotton	25	0	25	0.0	100.0	
		Eggplant	30	0	30	0.0	100.0	
		All host plants	84	0	84	0.0	100.0	
		JiNan	2005	Cotton	24	24	0	100.0
2006			Eggplant	28	28	0	100.0	0.0
	Cotton		31	31	0	100.0	0.0	
	All host plants		59	59	0	100.0	0.0	
2007	Cotton		47	18	29	38.3	61.7	
2008	Japanese hop		24	1	23	4.2	95.8	
	Cotton		29	0	29	0.0	100.0	
	Eggplant		19	4	15	21.1	78.9	
	All host plants		72	5	67	6.9	93.1	
LiaoCheng	2005		Cucumber	33	33	0	100.0	0.0
			Cotton	39	38	1	97.4	2.6
			All host plants	72	71	1	98.6	1.4

TABLE 1. (CONTINUED) DATA ON WHITEFLY COLLECTION AND BIOTYPE DETERMINATION BASED ON THE MARKERS.

Location	Year	Host plant	Total individuals	Number of biotype		Percentage of biotype (%)		
				B	Q	B	Q	
	2006	Winter squash	27	23	4	85.2	14.8	
		Cotton	28	13	15	46.4	53.6	
		Morning glory	11	10	1	90.9	9.1	
		Eggplant	26	13	13	50.0	50.0	
		Cucumber	24	14	10	58.3	41.7	
		All host plants	116	73	43	62.9	37.1	
	2007	Cotton	54	0	54	0.0	100.0	
		Eggplant	51	8	43	15.7	84.3	
		All host plants	105	8	97	7.6	92.4	
	2008	Japanese hop	30	1	29	3.3	96.7	
		Cotton	30	0	30	0.0	100.0	
		Eggplant	26	1	25	3.8	96.2	
		All host plants	86	2	84	2.3	97.7	
	LinYi	2006	Cotton	28	28	0	100.0	0.0
			Eggplant	29	29	0	100.0	0.0
Cucumber			8	8	0	100.0	0.0	
All host plants			65	65	0	100.0	0.0	
2007		Cucumber	13	11	2	84.6	15.4	
		Cotton	21	21	0	100.0	0.0	
		Eggplant	20	19	1	95.0	5.0	
		All host plants	54	51	3	94.4	5.6	
2008		Cotton	26	1	25	3.8	96.2	
		Eggplant	20	4	16	20.0	80.0	
		All host plants	46	5	41	10.9	89.1	
ZaoZhuang		2005	Cucumber	30	30	0	100.0	0.0
	2006	Cucumber	29	29	0	100.0	0.0	
	2008	Pepper	26	0	26	0.0	100.0	
		Japanese hop	29	0	29	0.0	100.0	
		Cotton	29	0	29	0.0	100.0	
		Eggplant	24	0	24	0.0	100.0	
		All host plants	108	0	108	0.0	100.0	

mtCOI fragment (about 620bp) was first cleaved by the restriction endonucleases *VspI* (Khasdan et al. 2005) and then the uncut fragment was cleaved by the restriction endonucleases *StuI* (Ueda 2006). All of the *mtCOI* that could be cut by *VspI* should be biotype Q and *mtCOI* cut by *StuI* should be B.

RESULTS AND DISCUSSION

Our results shown in Fig. 1 and Table 1 revealed the following: In 2005, the biotype of *B. tabaci* populations in JiNan, LiaoCheng, ZaoZhuang were determined and biotype Q was only found in LiaoCheng in very low proportion (1.4%). In 2006, biotype Q was found in DeZhou (38.0%), ZiBo (38.9%), ShouGuang (62.6%), LiaoCheng (37.1%) and was absent in JiNan, LinYi and ZaoZhuang. By 2007, biotype Q dominated in most locations, DeZhou (71.8%), ZiBo (92.7%), ShouGuang (96.0%), LiaoCheng (92.4%), and bio-

type Q also was found in JiNan (61.7%) and LinYi (5.6%). In 2008, Q biotype comprised 100.0%, 93.1%, 100.0%, 93.1%, 97.7%, 89.1% and 100.0% of the *B. tabaci* population in DeZhou, ZiBo, ShouGuang, JiNan, LiaoCheng, LinYi and ZaoZhuang, respectively.

The present results are consistent with previous research on cotton and eggplant (Chu et al. 2010). These results suggest that the changes of *B. tabaci* biotypes occurred not only on the cotton and eggplant but also on the other plants including crops and weeds in Shandong Province during the past several years.

The displacement mechanism of biotypes Q and B remains uncertain, though the increase of biotype Q in many countries may be due to application of insecticides because biotype Q possesses greater resistance to insecticides than biotype B, but ecological and economic factors should be also considered. For example, the host plants that biotype Q preferred also might mediate the competi-

tion of B and Q. Multiple introduction of biotype Q from the other provinces or regions through human activities or natural sources should not be neglected.

Overall, our present result showed that the biotypes of *B. tabaci* changed greatly and *B. tabaci* biotype Q has been increasing on all kinds of hosts during the past several years. The displacement of biotypes B by Q has been occurring in the province as a whole. The speed of the displacement of biotypes B and Q was fast and essentially a cryptic invasion (Geller et al. 1997) because the biotypes are morphologically indistinguishable. The displacement mechanism should be further researched and such knowledge might guide the application of the insecticides or adjustment of the crops to effectively control the pest. Differentiation of *B. tabaci* biotypes is important, and molecular markers are important discrimination tools.

ACKNOWLEDGMENTS

This work was funded by the Outstanding Youth Science Foundation of Shandong Province (JQ200811), National Basic Research and Development Program (2009CB119200), National Natural Science Foundation of China (30771410), Special Scientific Research Fund for Commonweal Trade of China (200803005) and Key Projects in the National Science & Technology Pillar Program in the Eleventh Five-year Plan Period (2006BAD08A18).

REFERENCES CITED

- BROWN, J. K. 2007. The *Bemisia tabaci* complex: genetic and phenotypic variability drives begomovirus spread and virus diversification. Plant Disease APS-Net Feature Article December-January 2006-07. <http://www.apsnet.org/online/feature/btabaci/>.
- CHU, D., JIANG, T., LIU, G. X., JIANG, D. F., TAO, Y. L., FAN, Z. X., ZHOU, H. X., AND BI, Y. P. 2007. Biotype status and distribution of *Bemisia tabaci* (Hemiptera: Aleyrodidae) in Shandong Province of China based on mitochondrial DNA markers. Environ. Entomol. 36: 1290-1295.
- CHU, D., WAN, F. H., ZHANG, Y. J., AND BROWN, J. K. 2010. Change in the biotype composition of *Bemisia tabaci* in Shandong Province of China From 2005 to 2008. Env. Entomol. 39. DOI: 10.1603/EN09161.
- CHU, D., ZHANG, Y. J., BROWN, J. K., CONG, B., XU, B. Y., WU, Q. J., AND ZHU, G. R. 2006. The introduction of the exotic Q biotype of *Bemisia tabaci* (Gennadius) from the Mediterranean region into China on ornamental crops. Florida Entomol. 89: 168-174.
- DENNEHY, T. J., DEGAIN, B. A., HARPOLD, V. S., BROWN, J. K., MORIN, S., FABRICK, J. A., BYRNE, F. J., AND NICHOLS, R. L. 2005. New challenges to management of whitefly resistance to insecticides in Arizona, p. 31 *In* University of Arizona Cooperative Extension, Vegetable Report. http://cals.arizona.edu/pubs/crops/az1382/az1382_2.pdf
- GELLER, J. B., WALTON, E. D., GROSHOLZ, E. D., AND RUIZ, G. M. 1997. Cryptic invasions of the crab *Carcinus* detected by molecular phylogeography. Mol. Ecol. 6: 901-906.
- HOROWITZ, A. R., KONTSEDALOV, S., KHASDAN, V., AND ISHAAYA, I. 2005. Biotypes B and Q of *Bemisia tabaci* and their relevance to neonicotinoid and pyriproxyfen resistance. Arch. Insect Biochem. Physiol. 58: 216-225.
- KHASDAN, V., LEVIN, I., ROSNER, A., MORIN, S., KONTSEDALOV, S., MASLENIN, L., AND HOROWITZ, A. R. 2005. DNA markers for identifying biotypes B and Q of *Bemisia tabaci* (Hemiptera: Aleyrodidae) and studying population dynamics. Bull. Entomol. Res. 95: 605-613.
- UEDA, S. 2006. Simple and rapid detection by mtCOI PCR-RFLP to distinguish the Q biotype of *Bemisia tabaci*. Kyushu Plant Protection Research 52: 44-48.
- WU, X. X., LI, Z. X., HU, D. X., AND SHEN, Z. R. 2002. Using RAPD-PCR to distinguish biotypes of *Bemisia tabaci* (Homoptera: Aleyrodidae) in China. Entomologia Sinica 9:1-8.