



Some Fungus-Growing Ants (Hymenoptera: Formicidae) from Northeastern Mexico

Author: Sanchez-Peña, Sergio R.

Source: Florida Entomologist, 93(4) : 501-504

Published By: Florida Entomological Society

URL: <https://doi.org/10.1653/024.093.0404>

BioOne Complete (complete.BioOne.org) is a full-text database of 200 subscribed and open-access titles in the biological, ecological, and environmental sciences published by nonprofit societies, associations, museums, institutions, and presses.

Your use of this PDF, the BioOne Complete website, and all posted and associated content indicates your acceptance of BioOne's Terms of Use, available at www.bioone.org/terms-of-use.

Usage of BioOne Complete content is strictly limited to personal, educational, and non - commercial use. Commercial inquiries or rights and permissions requests should be directed to the individual publisher as copyright holder.

BioOne sees sustainable scholarly publishing as an inherently collaborative enterprise connecting authors, nonprofit publishers, academic institutions, research libraries, and research funders in the common goal of maximizing access to critical research.

SOME FUNGUS-GROWING ANTS (HYMENOPTERA: FORMICIDAE) FROM NORTHEASTERN MEXICO

SERGIO R. SANCHEZ-PEÑA

Departamento de Parasitología, Universidad Autónoma Agraria Antonio Narro, Saltillo, Coahuila 25315, Mexico

ABSTRACT

The fungus-growing ants (Hymenoptera: Formicidae: Attini) of northeastern Mexico are poorly known. Herein new distributional records in NE Mexico and habitat observations are provided for the fungus-growing ants *Apterostigma mexicanum* Lattke, *Atta texana* (Buckley), *Cyphomyrmex rimosus* (Spinola), *Mycocepurus smithii* Forel, *Mycetosoritis hartmanni* Wheeler, *Sericomyrmex aztecus* Wheeler, *Trachymyrmex smithi* Buren, and *Trachymyrmex turrifex* Wheeler.

Key Words: fungus, symbiosis, ants, Tamaulipas, attine ants

RESUMEN

Son poco conocidas las hormigas cultivadoras de hongos (Hymenoptera: Formicidae: Attini) presentes en el noreste de México. En el presente trabajo se reportan nuevos datos de distribución y de habitat en el noreste de México, para las hormigas cultivadoras de hongos *Apterostigma mexicanum* Lattke, *Atta texana* (Buckley), *Cyphomyrmex rimosus* (Spinola), *Mycetosoritis hartmanni* Wheeler, *Mycocepurus smithii* Forel, *Sericomyrmex aztecus* Wheeler, *Trachymyrmex smithi* Buren y *Trachymyrmex turrifex* Wheeler.

Translation provided by the author.

Fungus-growing ants of the tribe Attini are important model systems in studies on behavioral ecology, coevolution, mutualism, parasitism and biogeography (Currie et al. 2003; Munkacsi et al. 2004; Mikheyev et al. 2007). Attine ants are a predominantly Neotropical group with few successful extensions into the Nearctic ecozone (Weber 1972). Species of *Cyphomyrmex*, *Atta texana* (Buckley) and particularly *Trachymyrmex septentrionalis* (McCook) are the northernmost distributed of all attine ants (Weber 1972). Northeastern (NE) Mexico (states of Coahuila, Nuevo León, San Luis Potosí, and Tamaulipas) includes the distribution limits of several attines; this region, and particularly the state of Tamaulipas encompasses the contact of the Nearctic and Neotropical realms. The attines of NE Mexico are poorly known. Along the Gulf of Mexico coastal plain, the northernmost colonies of the Mexican leaf-cutting ant, *Atta mexicana* (Smith) have been reported in Tamaulipas, in San Fernando, and in the state of Nuevo León at Sabinas Hidalgo and Cerralvo, at the piedmont of the Sierra Madre Oriental (Sánchez-Peña 2005). In Tamaulipas state, Flores-Maldonado et al. (1999) reported *Apterostigma pilosum* Mayr and *Trachymyrmex turrifex* Wheeler in Cañón del Novillo, near Ciudad Victoria. *Trachymyrmex saussurei* (Forel) a neotropical species, exists in Gómez Farías, Tamaulipas (Rabeling et al. 2007). The present paper includes distribution records and habitat observations for attine species, some of which were not expected to occur or are not frequently reported in NE Mexico.

MATERIALS AND METHODS

With the exception of some specimens of *Cyphomyrmex rimosus* Spinola from Matamoros caught with pitfall traps, all specimens were collected by hand by the author during directed searches for workers. Descriptions used for identification are listed with each species. Records are reported by state (uppercase) and municipality (municipio); more specific localities are mentioned when pertinent. Voucher specimens are deposited at the Entomological Collection of Universidad Autónoma Agraria Antonio Narro, (UAAAN), Saltillo, Coahuila, Mexico. Specimens of taxa marked with an asterisk (*) have been deposited at the Bohart Museum of Entomology, University of California at Davis (UCDC).

RESULTS AND DISCUSSION

Table 1 is a summary of collected attine ants. These are detailed below.

Apterostigma mexicanum Lattke (*). SAN LUIS POTOSI: Municipio of Aquismón. 11.IV.2008. N 21°35'59", W 99°06'12", 980 masl (meters above sea level). About 200 m above the Sótano de las Golondrinas (Pit of Swallows). *Apterostigma* includes the most basal extant attines along with *Myrmicocrypta* and *Mycocepurus* (Currie et al. 2003). Workers of *A. mexi-*

TABLE 1. ATTINE ANTS COLLECTED IN NORTHEASTERN MEXICO, THEIR LOCALITIES AND GENERAL HABITATS.

| Ant species | Locality/State | Habitat |
|--------------------------------|--|---|
| <i>Apterostigma mexicanum</i> | 200 m above Pit of Swallows/ San Luis Potosí | Clearing, former tropical rainforest |
| <i>Atta texana</i> | Jiménez/ Coahuila Múzquiz/ Coahuila | Temperate gallery forest of <i>Taxodium</i> , <i>Platanus</i> , <i>Salix</i> |
| <i>Cyphomyrmex rimosus</i> | Many localities/ Tamaulipas and Nuevo León | Springs, temperate gallery forest, mesic habitats |
| <i>Mycetosoritis hartmanni</i> | Escape, Cadereyta/Nuevo León El Encino, Llera/Tamaulipas | Under cover crop of organic farm Tropical gallery forest with <i>Taxodium</i> , <i>Ficus</i> and <i>Inga</i> |
| <i>Mycocepurus smithii</i> | Guadalupe/Nuevo León | Temperate gallery forest of <i>Platanus</i> and <i>Juglans</i> |
| <i>Sericomyrmex aztecus</i> | Tamul/San Luis Potosí Taninul, Valles/San Luis Potosí | Tropical gallery forest Medium height tropical forest with <i>Bursera</i> and <i>Ficus</i> |
| <i>Trachymyrmex smithi</i> | Zapalinamé, Saltillo/Coahuila | Xerophilous temperate oak forest |
| <i>Trachymyrmex turrifex</i> | Santa María, Pesquería/Nuevo León García/Nuevo León Matamoros/Tamaulipas | Garden; desert scrub area with <i>Parkinsonia texana</i> Desert scrub with <i>P. texana</i> Garden in urban area |

canum were collected on soil in extensive clearing in medium-tall rainforest that included the trees *Brosimum alicastrum* Sw. (ojite, ramón), *Bursera simaruba* (L.) Sarg. (chaká, gumbo-limbo), *Ficus cotinifolia* H.B.K. (higuerón, strangler fig) near the transition to cloud forest. Apparently previously known from 5 specimens, all from tropical forest in Veracruz state, near Córdoba (Lattke 1997).

Atta texana (Buckley) (*). COAHUILA: Municipio of Jiménez, town of Jiménez. 22.IX.2006. 29°03'04"N, 100°40'11"W, 240 masl. Municipio of Múzquiz, town of Múzquiz, Sabinas River. 5.VII.2008. 27°58'09", W 101°34'53", 490 masl. At Jiménez town, the Texas leaf-cutting ant was collected within 200 m from the Rio Grande (Rio Bravo) river, and also 15 km to the south, down the river. In both sites *A. texana* lives on the deeper soil flats ("vegas") along the river, in woods and clearings with pecans [(*Carya illinoensis* (Wangenh.) K. Koch)], walnut (*Juglans* sp.), and huisache, (*Acacia farnesiana* (L.) Willd.). Mature colonies form nest mounds 5-10 m in diameter. At Múzquiz, *A. texana* was in a riparian forest of baldecypress (*Taxodium* cf. *mucronatum* Ten.), willow (*Salix nigra* Marshall) and sycamore (*Platanus* sp.). The Múzquiz locality is near the southwestern distribution limit for *A. texana*. There are few reports of this ant in Mexico, and no recent records of the reported populations of *A. texana* along the Gulf of Mexico, in the states of Tamaulipas, Veracruz, and Tabasco (Smith 1963). There is an unexpected, extremely disjunct record from Tehuacán, central Mexico (Ríos-Casanova et al.

2004). Coronado-Padilla et al. (1972) reported it from Allende, Coahuila (about 100 km NE from Múzquiz). The distribution of *Atta* spp. is quite fragmented in Northeastern Mexico; the factors responsible are possibly lack of moisture and soil type.

Cyphomyrmex rimosus (Spinola). Widespread morphospecies, known from Texas (San Marcos) and southern USA to Argentina (Wheeler 1907; Longino 2008) thus coordinates are not given here. Collected in mesic microhabitats, mainly springs and streams between 10 and 500 masl. NUEVO LEON: municipios of Cerralvo, at the spring; Guadalupe: La Pastora State Park, 10.VII.2006; Montemorelos: Rancho Marrufo, km 6 Montemorelos-General Terán highway, in wooded area along fencerow, 8.XII.2007; Pesquería: Santa María la Floreña, along tree-covered irrigation channel from Río Pesquería, 24.VIII.2008; Sabinas Hidalgo, at the spring, 18.VII.2006. TAMAULIPAS: municipios of Gómez Farías, San Pedrito town, in garden, 25.VIII.2008; Matamoros, Ejido (community) El Longoreño, gallery forest and grassy areas near the Rio Grande and ponds, 31.VIII.2007.

Mycetosoritis hartmanni Wheeler. NUEVO LEON: Municipio of Cadereyta, Rancho El Escape, Ejido Jerónimo Treviño. 8.XII.2007. N 25°32'25", W 99°54'17", 504 masl. TAMAULIPAS: Municipio of Llera, El Encino-La Libertad road, near Sabinas River (Río Sabinas), 23.IX.2006. N 23°08'09", W 99°08'53", 110 masl. The Nuevo León locality is an irrigated, organic vegetable farm at a semiarid site, on alluvial soils

near the Santa Catarina river. There, several small plots (<1 ha) were grown to cover crops (Fabaceae), providing full shade. *Mycetosoritis hartmanni* workers were foraging under the cover crop, at the northern edge of one plot, at noon. The Tamaulipas location is disturbed tropical gallery forest with the trees *T. mucronatum*, *Ficus* sp., and *Inga vera* Willd., about 20 km north of Gómez Farías. *Mycetosoritis hartmanni* is an infrequently collected species (Mackay 1998; Longino 2009). In Mexico, this ant is known from the humid, medium-height tropical forest at Gómez Farías, Tamaulipas (Jusino-Atresino & Phillips 1992). The type locality is warm-temperate (Austin, Texas) (Wheeler 1907). Longino (2009) lists *M. hartmanni* from south Texas (Laguna Atascosa Wildlife Refuge) in coastal, low shrubby vegetation. *Mycetosoritis hartmanni* has not been reported living in the ephemeral habitat of annual agricultural crops.

Mycocepurus smithii Forel. NUEVO LEON: Municipio of Guadalupe, La Pastora State Park. 10.VII.2006 and 28.XII.2006. N 25°39'53", W 100°15'15", 500 masl. This 200-ha park includes a remnant patch of old forest with interior live oaks (*Quercus fusiformis* Small) and anacua (*Erethia anacua* (Terán & Berland.) I. M. Johnst.) and gallery forest of sycamores (*Platanus* sp.), walnut (*Juglans* sp.) and bald cypress (*T. mucronatum*), on intermittent creeks running into the La Silla river. This park has been engulfed by the metropolitan area of Monterrey since at least 20 years ago. An aggregation of nest entrances of *Mycocepurus* colonies were in the gallery forest, on the flat surface of sandy banks along shallow dry creeks. Workers were observed at 5:00 PM on 10.VII.2006 at 30 m from the dry creek, foraging in the forest litter. Workers of *C. rimosus* were present also. On 28.XII.2006, foraging was observed at 13:00 h, after night temperatures of 5–10°C; the species apparently does not become inactive in winter here, as the attine *Trachymyrmex septentrionalis* does further north (Weber 1972). Local *Mycocepurus* appears to be mainly riparian, foraging in the same areas (moist microhabitats) as the ponerine ants *Pachycondyla villosa* (F.) and *Odontomachus* cf. *laticeps* Roger. Ants of open and disturbed habitats (*Pogonomyrmex barbatus* (Smith), *Solenopsis geminata* (Fabricius) and *Dorymyrmex* sp.) nested within 20 meters of *M. smithii*, in clearings. Rosas-Mejía et al. (2008) reported *Mycocepurus* sp. as an urban ant in Ciudad Victoria, 300 km south of Monterrey. Jusino-Atresino & Phillips (1992) collected *M. smithii* from the subdeciduous tropical forest of Gómez Farías, Tamaulipas, about 400 km south of Monterrey. Also reported from more distant tropical sites in Mexico to the south (states of Jalisco, Nayarit, San Luis Potosí, Tabasco, and Veracruz) (Mackay et al. 2004; Vázquez-Bolaños 2007), in deciduous and subdeciduous tropical forests, oak-

pine forest, and secondary vegetation in urban areas (Vázquez-Bolaños 2007). The report herein is the northernmost for the species; it is in the Nearctic ecozone at almost the same latitude of Brownsville, Texas, and in the Rio Grande Basin as well. This locality is apparently the most xeric (annual average precipitation 551 mm) and temperate (SAGARPA 2002) known for *Mycocepurus* in most of its range. Ants of the genera *Mycetosoritis* and *Mycocepurus* are small, cryptic, and infrequently collected (Mackay 1998).

Sericomyrmex aztecus Forel (*). SAN LUIS POTOSÍ: Municipio of Aquismón. 10.IV.2008. N 21°48'09", W 99°10'49", 200 m masl. Rio Santa María, Tamul. Undercover of tropical gallery forest including trees of *B. simaruba* and *F. cotinifolia*. Municipio of Ciudad Valles, N 21°56'11", W 98°53'23", 100 masl. Near gardens of hotel at Taninul. This is a mesic microhabitat in secondary low and medium-height tropical forest with *B. simaruba* and *F. cotinifolia*. The northernmost species of *Sericomyrmex* is *S. aztecus* from Mexico. Longino (2008) collected *S. aztecus* in Chiapas, near the Guatemala border. The Taninul, Ciudad Valles location reported herein, about 40 km from the state of Tamaulipas, is probably the northernmost record for the genus. Both Wheeler (1925) and Longino (2008) consider that Mesoamerican (or most) species of *Sericomyrmex* are very similar, and probably describe intraspecific variation.

Trachymyrmex smithi Buren. COAHUILA: Municipio and city of Saltillo. 19.VII.2005. N 25°21'36", W 100°58'42", 1680 masl. Lomas de Lourdes, Sierra de Zapalinamé. Workers foraging after summer rains on reddish-soil hills, above SE Saltillo. The habitat is temperate, an ecotone of xerophilous brush and forest with oaks (*Quercus laeta* Liebm. and the endemic *Q. saltillensis* Trel.), madrone (*Arbutus xalapensis* Kunth), and pistache (*Pistacia mexicana* H.B.K.). The type locality for *T. smithi* is La Rosa, General Cepeda, Coahuila, about 50 km W, in rocky Chihuahuan desert habitats (Buren 1944). Mackay & Mackay (2002) list it from Chihuahua state, Mexico. From the records, this species appears to occur in high, temperate deserts where summer temperatures are usually below 33°C.

Trachymyrmex turrifex Wheeler. NUEVO LEON: Municipio of Pesquería, town of Santa María la Floreña, 24.VIII.2008. N 25°44'06", W 99°49'44", 320 masl. Nest in shaded sandy soil in garden; natural vegetation is thorny brush with honey mesquite (*Prosopis glandulosa* Torr.), huisache, palo verde (*Parkinsonia texana* (A. Gray) Watson), and desert hackberry (*Celtis pallida* Torr.). Municipio of García. 17.IX.2009. N 25°42'38", W 100°32'53", 840 masl. On sandy soil in thorny brush as above. TAMAULIPAS: Municipio of Matamoros. N 25°51'25", W 97°30'31", 10 masl. 22.VII.2009. Under trees in urban garden

in sandy soil; nest aggregations (6 nests/10 m²). Known from matorral at Vallecillo, Nuevo León (40 km S of Laredo, Texas) (Buren (1944)). This is possibly the most common *Trachymyrmex* species in the lower, hotter arid areas of NE Mexico.

ACKNOWLEDGMENTS

W. Mackay, P. Ward, and C. Rabeling kindly verified the identification of specimens. W. Mackay and 2 reviewers also improved the manuscript. Thanks to those persons responsible for the wonderful web-available identification materials for ants. Supported by Universidad Autónoma Agraria Antonio Narro (Dirección de Investigación, item 01-04-0206-39).

REFERENCES CITED

- BUREN, W. F. 1944. A new fungus-growing ant from Mexico. *Psyche* 51(1-2): 5-7.
- CORONADO-PADILLA, R., MORALES, A., AND ESPINOSA, E. 1972. Distribución geográfica de las especies de hormigas arrieras existentes en la República Mexicana. *Folia Entomol. Mexicana* 23/24: 95-96.
- CURRIE, C. R., WONG, B., STUART, A. E., SCHULTZ, T. R., REHNER, S. A., MUELLER, U. G., SUNG, G. H., SPATAFORA, J. W., AND STRAUS, N. A. 2003. Ancient tripartite coevolution in the attine ant-microbe symbiosis. *Science* 299: 386-388.
- FLORES-MALDONADO, K. Y., PHILLIPS S. A., AND SANCHEZ-RAMOS, G. 1999. The myrmecofauna (Hymenoptera: Formicidae) along an altitudinal gradient in the Sierra Madre Oriental of northeastern Mexico. *Southwest. Nat.* 44(4): 457-461.
- JUSINO-ATRESINO, R., AND PHILLIPS, JR., S. A. 1998. Myrmecofauna en la reserva ecológica de la biosfera "El Cielo", Tamaulipas, México. *Biotam* 4: 41-54.
- LATTKE, J. E. 1997. Revisión del género *Apterostigma* Mayr (Hymenoptera: Formicidae). *Arq. Zool. (São Paulo)* 34: 121-221.
- LONGINO, J. T. 2009. Ants of Costa Rica. <http://academic.evergreen.edu/projects/ants/AntsofCostaRicaA.html>. Accessed 5 December 2009.
- MACKAY, W. P. 1998. Dos especies nuevas de hormigas de la tribu Attini de Costa Rica y México: *Mycetosoritis vinsoni* y *Mycoccephurus curvispinosus* (Hymenoptera: Formicidae). *Rev. Biol. Trop.* 46(2): 421-426.
- MACKAY W. P., AND MACKAY, E. 2002. The Ants of New Mexico. The Edwin Mellen Press, Lewiston, NY.
- MACKAY, W. P., MAES, J-M, ROJAS-FERNANDEZ, P., AND LUNA, G. 2004. The ants of North and Central America: the genus *Mycoccephurus* (Hymenoptera: Formicidae). 7pp. *J. Insect Science*, 4:27.
- MIKHEYEV, A. S., MUELLER, U. G., AND BOOMSMA, J. J. 2007. Population genetic signatures of diffuse coevolution between leaf-cutting ants and their cultivar fungi. *Mol. Ecol.* 16(1): 209-216.
- MUNKACSI, A. B., PAN, J. J., VILLESSEN, P., MUELLER, U. G., BLACKWELL, M., AND MCLAUGHLIN, D. J. 2004. Convergent coevolution in the domestication of coral mushrooms by fungus-growing ants. *Proc. Royal Soc., London B* 1776-1782.
- RABELING, C., COVER, S. P., MUELLER, U. G., AND JOHNSON, R. A. 2007. A review of the North American species of the fungus-gardening ant genus *Trachymyrmex* (Hymenoptera: Formicidae). *Zootaxa* 1664: 1-54.
- RÍOS-CASANOVA, L., A., VALIENTE-BANUET, A., AND RICO-GRAY, V. 2004. Las hormigas del Valle de Tehuacán (Hymenoptera: Formicidae): una comparación con otras zonas áridas de México. *Acta Zool. Mexicana* 20(1): 37-54.
- ROSAS-MEJÍA, M., HORTA-VEGA, J., FLORES-MALDONADO, K. Y., AND CORREA-SANDOVAL, A. 2008. Formicidae de la zona urbana de Ciudad Victoria, Tamaulipas, Mexico. *Memorias Congreso de la Soc. Mexicana Entomol. (Estrada-Venegas, E., Equihua-Martinez A., Padilla-Ramírez, J. and Mendoza-Estrada, A. Eds.). Colegio de Posgraduados, Texcoco, Mexico.*
- SAGARPA. 2002. Anuario Estadístico. Secretaría de Agricultura, Ganadería, Desarrollo Rural, Pesca y Alimentación. Mexico, D.F.
- SANCHEZ-PENA, S. R. 2005. Essays on Organismal Aspects of the Fungus-Growing Ant Symbiosis: Ecology, Experimental Symbiont Switches and Fitness of *Atta*, and A New Theory on The Origin of Ant Fungiculture. Ph. D. Dissertation, The University of Texas at Austin.
- SMITH, M. R. 1963. Notes on the leaf-cutting ants, *Atta* spp., of the United States and Mexico (Hymenoptera: Formicidae). *Proc. Entomol. Soc. Washington* 65: 299-302.
- WEBER, N. A. 1972. Gardening ants, the attines. *Mem. American Phil. Soc.* 92. Philadelphia, PA.
- WHEELER, W. M. 1907. The fungus-growing ants of North America. *Bull. American Mus. Nat. Hist.* 23: 669-807.
- WHEELER, W. M. 1925. A new guest ant and other new Formicidae from Barro Colorado Island, Panama. *Biol. Bull.* 49: 150-181.