

Natural Enemies Managing the Invasion of the Fig Whitefly, *Singhiella simplex* (Hemiptera: Aleyrodidae), Infesting a *Ficus benjamina* Hedge

Authors: Avery, Pasco B., Mannion, Catharine M., Powell, Charles A., McKenzie, Cindy L., and Osborne, Lance S.

Source: Florida Entomologist, 94(3) : 696-698

Published By: Florida Entomological Society

URL: <https://doi.org/10.1653/024.094.0338>

BioOne Complete (complete.BioOne.org) is a full-text database of 200 subscribed and open-access titles in the biological, ecological, and environmental sciences published by nonprofit societies, associations, museums, institutions, and presses.

Your use of this PDF, the BioOne Complete website, and all posted and associated content indicates your acceptance of BioOne's Terms of Use, available at www.bioone.org/terms-of-use.

Usage of BioOne Complete content is strictly limited to personal, educational, and non - commercial use. Commercial inquiries or rights and permissions requests should be directed to the individual publisher as copyright holder.

BioOne sees sustainable scholarly publishing as an inherently collaborative enterprise connecting authors, nonprofit publishers, academic institutions, research libraries, and research funders in the common goal of maximizing access to critical research.

NATURAL ENEMIES MANAGING THE INVASION OF THE FIG WHITEFLY, *SINGHIELLA SIMPLEX* (HEMIPTERA: ALEYRODIDAE), INFESTING A *FICUS BENJAMINA* HEDGE

PASCO B. AVERY¹, CATHARINE M. MANNION², CHARLES A. POWELL¹, CINDY L. MCKENZIE³ AND LANCE S. OSBORNE⁴

¹University of Florida, Institute of Food and Agricultural Sciences, Indian River Research and Education Center, 2199 South Rock Road, Fort Pierce, FL 34945

²Tropical Research and Education Center, University of Florida, Homestead, FL, 33031

³USDA, ARS, U.S. Horticultural Research Laboratory, Subtropical Insect Research Unit, 2001 S. Rock Rd., Ft. Pierce, FL 34945

⁴Mid Florida Research and Education Center, University of Florida, Apopka, FL 32703

The fig whitefly, *Singhiella simplex* (Singh), a recent adventive species native of Burma, China and India (Singh 1931) has become a major pest in Florida (Hodges 2007) feeding on and defoliating *Ficus* shrubs and trees. This pest, first discovered in Miami-Dade County in 2007, is spreading throughout Florida; and recently it was found in Saint Lucie County, FL (personal observation, Avery 2009). Literature on the biology of the fig whitefly is sparse and most references are extension documents (Mannion et al. 2008; Caldwell 2009; Mannion 2010). The life cycle may be similar to that of the other *Singhiella* species that are present in Florida, (*Singhiella citrifolii* (Morgan), with at least three generations per year (Hodges 2007). Leaves of *Ficus* that turn yellow prior to defoliation are one of the most obvious symptoms of a fig whitefly infestation. The fig whitefly is most commonly found infesting weeping fig (*Ficus benjamina*), but may eventually damage other species of *Ficus* as well (Mannion et al. 2008).

In the landscape, several natural enemies have been observed attacking this whitefly, which may play an important role in long term control. Awareness of these natural enemies is very important in making pesticide application decisions so as not to adversely affect them. Commonly observed natural enemies include ladybird beetle predators, *Harmonia axyridis* (Pallas), *Olla-v-nigrum* (Mulsant), *Exochomus children* Mulsant, *Chilocorus nigritis* (F.), *Curinus coeruleus* (Mulsant); parasitoids, *Encarsia protransvena* Viggiani, *Amitus bennetti* Viggiani & Evans; and lacewings, *Chrysopa* spp. (Mannion 2010). Moreover, entomopathogenic fungi may also play a role in managing this pest (Elliot et al. 2000; Torres-Barragán et al. 2004). However, no entomopathogenic fungus isolated from the fig whitefly in Florida has been reported to date.

During the fall of 2009, a fig whitefly invasion occurred on a *F. benjamina* hedge at a residence in Ft. Pierce, Florida. We used this opportunity to: 1) identify and inventory the natural enemies present, and 2) determine their effectiveness for

managing the fig whitefly over the Sep to Nov time period.

The layout for the study was a complete block design with four plots in a linear design. Each plot measured ~5 m of a *F. benjamina* hedge (~10-15 plants), which ran along a concrete block wall (~1.2 m tall) located at a residence in Ft. Pierce, Florida. Each hedge segment (northern 27°22'50.94 N × 80°22'00.22 W and southern 27°22'48.96 N × 80°22'00.25 W) was divided by a concrete driveway; and the northern side was more severely affected by leaf drop than the southern. Randomly chosen leaf samples (10 per plot) were detached from the hedge on a weekly basis for 7 wk. Each sample was placed into individual re-sealable plastic bags and brought back to the lab for examination under a binocular microscope (40X) to count the number of live, dead, or parasitized whitefly nymphs. Parasitism was recognized either by observing the development of the parasitoid inside the translucent nymphal case, or by observing a blackened (melanized) nymphal case with or without an exit hole. A total of 40 disks were punched with a #5 cork borer (50.3 mm² diam) near the center of each leaf on one side of the midrib. The abaxial and adaxial sides of 1 disk per leaf were used for observation and counting the number of nymphs of each category mentioned above. The total weekly mean (±SEM) number of nymphs, number parasitized, and percent parasitism on the leaf disks were determined for 49 d. Based on photos of the parasitized nymphs (Mannion et al. 2008; Mannion 2010), the parasitoids appear to be *E. protransvena*; however, *Encarsia* species identification was not confirmed.

Disks were then placed on moist filter paper in a Petri dish, covered, sealed with Parafilm™, and incubated at 25°C for 14 d to allow for the development of mycosis, and to determine percent mortality due to fungal pathogens. After incubation, 5 mycosed nymphs/disk randomly chosen were removed from the leaf disk using a pin and placed on water agar (20 per plate) for isolation and identification of the fungal pathogen. Percent nymphs infected with a fungal species were determined for 0, 14, and 49 d post initial observation. Agar plates

were sealed with Parafilm™, and incubated at 25°C under a 16 h photophase for at least 1 week. The fungi were re-isolated from the mycosed nymphs and grown on PDA plates for identification. Voucher fungal *in vitro* culture isolates were sent to Svetlana Gouli at the University of Vermont, Dr. Richard Humber at the USDA, Ithaca, New York and to Dr. Rob Samson’s research team at the CBS-KNAW Fungal Diversity Centre in The Netherlands for identification and deposition.

The insect pests observed included the fig whitefly, *S. simplex*, another whitefly, *Tetraleurodes fici* Quaintance & Baker (one parasitized with an exit hole) and the weeping ficus thrips, *Gynaikothrips uzeli* Zimmerman (Table 1). The parasitoids, *Encarsia* species (all appeared to be *E. protransvena*) were observed after parasitization, and within the melanized nymphal case with or without an exit hole. Adult ladybird beetles, *C. coeruleus*, *H. axyridis* and eggs and larvae of the green lacewing, *Chrysopa* species were observed on the leaves. The natural enzootic pathogenic fungi isolated were *Isaria fumosorosea* Wize, *Paecilomyces lilacinus* Thom (Samson), and *Lecanicillium*, *Fusarium*, and *Aspergillus* species.

The total percent nymphal mortality per leaf disk due to pathogenic fungi varied overtime. *I. fumosorosea* and *Lecanicillium*, *Aspergillus* and *Fusarium* species were isolated from 5, 5, 48 and 50% of the dead nymphs at d 0, respectively. After 14 d, 85 and 15% of the nymphs were infected with *Aspergillus* sp. and *P. lilacinus*, respectively. At d 49, *Aspergillus* and *Fusarium* species were isolated 65% and 35% of the time, respectively. The fungal species isolated from the mycosed nymphs were assumed to have caused the mortality of the insect; however, Koch’s postulate was not confirmed.

The total weekly mean (±SEM) number of nymphs, number parasitized, and percent parasitism observed on both sides of the leaf disks over 49 d were 57.7 ± 9.91, 5.9 ± 3.72, and 7.3 ± 3.62, respectively. The proportion of nymphal mortality on a weekly basis due to parasitism was ~10% up to 35 d post initial observation; after that, no parasitized nymphs (second- fourth instars) were observed on leaf disks.

Gerling et al. (2001) indicated that all *Encarsia* species parasitize and emerge from the fourth instar of their whitefly host, but attack mainly the second- fourth instar hosts. Therefore, the low number of older (third- fourth instars) nymphs available to be parasitized after 35 d, may have resulted from leaf drop. Throughout this field pilot study based on leaf disk samples, only ~10% mortality was attributed to parasitization by *Encarsia* species, ~90% was due to other natural causes; i.e. enzootic entomopathogenic fungi, predation, etc. This finding warrants confirmation through in-depth research.

This is the first record of Hypocreales fungi; *I. fumosorosea*, *P. lilacinus*, and *Aspergillus*, *Lecanicillium*, and *Fusarium* species being isolated

TABLE 1. INSECT PESTS AND NATURAL ENEMIES OBSERVED ON A RESIDENTIAL *FICUS BENJAMINA* HEDGE IN FT. PIERCE, FLORIDA BETWEEN SEP-NOV 2009.

Category	Order	Family	Scientific name	Common Name	Observations
Insect pests	Hemiptera	Aleyrodidae	<i>Singhiella simplex</i>	fig whitefly	feeding on leaves
	Hemiptera	Aleyrodidae	<i>Tetraleurodes fici</i>	whitefly	feeding on leaves
	Thysanoptera	Phlaeothripidae	<i>Gynaikothrips uzeli</i>	weeping ficus thrip	in leaf fold galls
Natural Enemies	Hymenoptera	Aphelinidae	<i>Encarsia</i> sp.*	parasitoid	parasitized fig whitefly nymphs
	Coleoptera	Coccinellidae	<i>Harmonia axyridis</i>	Multicolored Asian lady beetle	adults roaming on leaves
	Coleoptera	Coccinellidae	<i>Curinus coeruleus</i>	Metallic blue lady beetle	adults roaming on leaves
	Neuroptera	Chrysopidae	<i>Chrysopa</i> sp.	green lacewing	eggs and larvae on a leaves
	Hypocreales	Cordycipitaceae	<i>Isaria fumosorosea</i>	entomopathogenic fungi	leaf surface and on fig whitefly
	Hypocreales	Clavicipitaceae	<i>Paecilomyces lilacinus</i>	entomopathogenic fungi	leaf surface and on fig whitefly
	Hypocreales	Cordycipitaceae	<i>Lecanicillium</i> sp.	entomopathogenic fungi	leaf surface and on fig whitefly
	Hypocreales	Nectriaceae	<i>Fusarium</i> sp.	filamentous fungi	leaf surface and on fig whitefly
	Euriales	Trichocomaceae	<i>Aspergillus</i> sp.	green mold	leaf surface and on fig whitefly

*All appeared to be *E. protransvena*.

from dead fig whitefly nymphs on *F. benjamina*. *Aspergillus* and *Fusarium* species, not commonly known as entomopathogenic, are now being tested as potential biocontrol agents for controlling whitefly species (Panyasiri et al. 2007). The other fungal species, *I. fumosorosea*, *P. lilacinus* and *Lecanicillium* species are commonly used as biopesticides (de Faria & Wraight 2007) for controlling whitefly (Avery et al. 2008; Cabanillas & Jones 2009; Shinde et al. 2010), except *P. lilacinus* which is nematophagous. However, recently *P. lilacinus* has demonstrated potential for controlling whitefly species (Gökçe & Er 2005; Fiedler & Sosnowska 2007).

Based on this study, it would be recommended that the endemic population of parasitoids and enzootic entomopathogenic fungi be considered as part of a multi-trophic ecosystem, which may be adversely impacted by the application of any pesticide. Therefore, it is important to assess the long term ecological impact that a pesticide application against the fig whitefly may have on the natural enemies in the landscape ecosystem.

SUMMARY

Based on leaf disk samples taken in this field pilot study, only ~10% of the fig whitefly mortality was attributed to parasitization by *Encarsia* species, ~90% was due to other natural causes; i.e. enzootic entomopathogenic fungi, predation, etc. This is the first record of the Hypocreales fungi, *I. fumosorosea*, *P. lilacinus*, and *Aspergillus*, *Lecanicillium*, and *Fusarium* species being isolated from dead *S. simplex* nymphs on *F. benjamina*. It is important to assess the long term impacts pesticide applications made for managing the fig whitefly may have on the ecosystem, and especially the ecological impact on the natural enemies which include enzootic entomopathogenic fungi.

ACKNOWLEDGMENTS

Many thanks go to John Prokop for his labor and technical assistance, and Anna Sara Hill, for the fig whitefly insectary work, at the USDA-ARS, U.S. Horticultural Research Laboratory; to Eliza Duane and Phyllis Hebert for their technical assistance at IRREC. We thank Svetlana Gouli at the University of Vermont, Invertebrate Pathology and Microbial Pest Control Laboratory in Burlington, Vermont, Dr. Richard Humber at the USDA-ARS Collection of Entomopathogenic Fungal Cultures (ARSEF), in Ithaca, New York and Dr. Rob Samson's research team at the CBS-KNAW Fungal Diversity Centre, Utrecht, The Netherlands for identification of the fungal isolates. We are grateful for funding through the USDA's Tropical and Subtropical Agriculture Research (T-STAR) Program, American Floral Endowment, and the USDA-ARS Floriculture and Nursery Research Initiative.

REFERENCES CITED

- EVERY, P. B., FAULL, J., AND SIMMONDS, M. S. J. 2008. Effects of *Paecilomyces fumosoroseus* and *Encarsia formosa* on the control of the greenhouse whitefly: preliminary assessment of a compatibility study. *BioControl* 53: 303-316.
- CABANILLAS, H. E., AND JONES, W. A. 2009. Pathogenicity of *Isaria* sp. (Hypocreales: Clavicipitaceae) against the sweet potato whitefly B biotype, *Bemisia tabaci* (Hemiptera: Aleyrodidae). *Crop Prot.* 28: 333-337.
- CALDWELL, D. 2009. Fig whitefly is defoliating weeping fig trees and hedges. University of Florida, IFAS Extension Collier County Bulletin, May, 2009.
- DE FARIA, M. R., AND WRAIGHT, S. P. 2007. Mycoinsecticides and mycoacaricides: A comprehensive list with worldwide coverage and international classification of formulation types. *Biol. Control* 43: 237-256.
- ELLIOT, S. L., SABELIS, M. W., JANSSEN, A., VAN DER GEEST, L. P. S., BEERLING, E. A. M., AND FRANSEN, J. 2000. Can plants use entomopathogens as bodyguards? *Ecol. Lett.* 3: 228-235.
- FIEDLER, Z., AND SOSNOWSKA, D. 2007. Nematophagous fungus *Paecilomyces lilacinus* (Thom) Samson is also a biocontrol agent for control of greenhouse insects and mite pests. *BioControl* 52: 547-558.
- GERLING, D., ALOMAR, O., AND ARNO, J. 2001. Biological control of *Bemisia tabaci* using predators and parasitoids. *Crop Prot.* 20: 779-799.
- GÖKÇE, A., AND ER, M. K. 2005. Pathogenicity of *Paecilomyces* spp. to the glasshouse whitefly, *Trialeurodes vaporariorum*, with some observations on the fungal infection process. *Turk. J. Agric. For.* 29: 331-339.
- HODGES, G. 2007. The Fig Whitefly *Singhiella simplex* (Singh) (Hemiptera: Aleyrodidae): A New Exotic Whitefly Found on *Ficus* Species in South Florida. Pest Alert DPI-FDACS <http://www.freshfromflorida.com/pi/enpp/ento/Singhiella%20simplex.html>
- MANNION, C. 2010. Whiteflies in the landscape. University of Florida, IFAS Extension Bulletin, Nov. 2010. <http://trec.ifas.ufl.edu/mannion/pdfs/Whitefly%20in%20the%20Landscape%20%20Nov2010.pdf>
- MANNION, C., OSBORNE, L., HUNSBERGER, A., MAYER, H., AND HODGES, G. 2008. *Ficus* whitefly: A new pest in South Florida. University of Florida, IFAS Extension Bulletin, Aug., 2008.
- PANYASIRI, C., ATTATHOM, T., AND POEHLING, H. M. 2007. Pathogenicity of entomopathogenic fungi-potential candidates to control insect pests on tomato under protected cultivation in Thailand. *J. Plant Dis. Prot.* 114: 278-287.
- SHINDE, S. V., PUROHIT, M. S., SABALPARA, A. N., AND PATEL, M. B. 2010. First report of entomopathogenic fungus *Lecanicillium lecanii* (Zimm.) Zare and Games on sugarcane whitefly *Aleurolobus barodensis* (Maskell) from Gujarat. *Biosci. Trends* 3: 1.
- SINGH, K. 1931. A contribution towards our knowledge of the Aleyrodidae (Whiteflies) of India. *Mem. Dep. Agric. India* 12: 1-98.
- TORRES-BARRAGÁN, A., ANAYA, A. L., ALATORRE, R., AND TORIELLO, C. 2004. Entomopathogenic fungi from 'El Eden' Ecological Reserve, Quintana Roo, Mexico. *Mycopathologia* 158: 61-71.