

## **Fruit Flies and Lance Flies (Diptera: Tephritoidea) and Their Host Plants in a Conservation Unit of the Cerrado Biome in Tocantins, Brazil**

Authors: Bomfim, Darcy A. Do, Gislotti, Laura J., and Uchôa, Manoel A.

Source: Florida Entomologist, 97(3) : 1139-1147

Published By: Florida Entomological Society

URL: <https://doi.org/10.1653/024.097.0321>

---

BioOne Complete ([complete.BioOne.org](https://complete.BioOne.org)) is a full-text database of 200 subscribed and open-access titles in the biological, ecological, and environmental sciences published by nonprofit societies, associations, museums, institutions, and presses.

Your use of this PDF, the BioOne Complete website, and all posted and associated content indicates your acceptance of BioOne's Terms of Use, available at [www.bioone.org/terms-of-use](https://www.bioone.org/terms-of-use).

Usage of BioOne Complete content is strictly limited to personal, educational, and non - commercial use. Commercial inquiries or rights and permissions requests should be directed to the individual publisher as copyright holder.

---

BioOne sees sustainable scholarly publishing as an inherently collaborative enterprise connecting authors, nonprofit publishers, academic institutions, research libraries, and research funders in the common goal of maximizing access to critical research.

# FRUIT FLIES AND LANCE FLIES (DIPTERA: TEPHRITOIDEA) AND THEIR HOST PLANTS IN A CONSERVATION UNIT OF THE CERRADO BIOME IN TOCANTINS, BRAZIL

DARCY A. DO BOMFIM<sup>1</sup>, LAURA J. GISLOTT<sup>2</sup> AND MANOEL A. UCHÔA<sup>1,3,\*</sup>

<sup>1</sup>Programa de Pós-Graduação em Entomologia e Conservação da Biodiversidade, Universidade Federal da Grande Dourados (UFGD), Caixa Postal 241, Avenida Guaicurus Km 12, 79804-970 Dourados-MS, Brazil  
darcybombfim@hotmail.com

<sup>2</sup>UNICAMP, Campinas, SP, Brazil; lauragislotti@gmail.com

<sup>3</sup>Researcher Fellow of National Council for Scientific and Technological Development (CNPq) of Brazil

\*Corresponding author; E-mail: uchoa.manoel@gmail.com

## ABSTRACT

Fruit flies (Tephritidae) and lance flies (Lonchaeidae) are specialized in exploring fruits as resource, and the native Angiospermae represent potential hosts for their frugivorous larvae. The aim of this study was to investigate the relationship of fruit fly and lance fly species with native host fruits in the Parque Estadual do Lajeado, Palmas, Tocantins, Brazil. The survey was carried out from May 2011 to Apr 2013, with sampling of fruits to obtain third instar larvae, pupae and adults. We sampled fruits of 18 plant species of which 7 were infested by fruit flies or lance flies, and from which emerged 888 adults of *Neosilba* spp. (Lonchaeidae), and 81 adults of *Anastrepha* (Tephritidae). *Neosilba zadolicha* emerged from fruit of *Siparuna camporum*, *Dyospiros hispida*, *Annona coreacea*, *Mouriri pusa*, *Cheiloclinium cognatum* and *Salacia crassifolia*. *Neosilba bifida*, *N. glaberrima*, and *Neosilba* **n. sp.1**, emerged from fruits of *D. hispida*. The species of *Anastrepha* obtained were: *A. serpentina* and *A. nr. mucronota* reared from *S. crassifolia* and *C. cognatum* fruits; *A. obliqua* from *S. crassifolia*, *A. sororcula* from *Myrcia guianensis*, *A. turpiniae*, and *A. zenildae* from *M. pusa*. Of the seven host fruit species, four were infested by Tephritidae, and six by Lonchaeidae. These are the first records for: *A. sororcula* in fruits of *Myrcia guianensis*, *A. nr. mucronota* in *Salacia crassifolia* and *Cheiloclinium cognatum*, *A. serpentina* in *Cheiloclinium cognatum*, and *A. turpiniae* and *A. zenildae* in *Mouriri pusa*. The associations of fruit flies and lance flies with host fruits is locally and temporally dependent on the flora, since the species of flies require available fruit species. In this survey we observed the highly polyphagous *A. obliqua*, that in this survey to colonize only one species of fruit.

Key Words: biodiversity, cerrado of northern Brazil, frugivory, new records, phytophagy

## RESUMO

As moscas das frutas (Tephritidae e Lonchaeidae) são um grupo especializado de insetos em explorar frutas como recurso, e as Angiospermas nativas representam potenciais hospedeiros para suas larvas frugívoras. O objetivo deste estudo foi investigar a associação de espécies de moscas de frutas com plantas hospedeiras nativas no Parque Estadual do Lajeado, em Palmas, Tocantins, Brasil. A pesquisa foi realizada a partir de maio de 2011 a abril de 2013, com amostragem de frutas para obter larvas de terceiro instar, pupas e adultos. Foram amostrados frutos em 18 espécies de plantas, sendo sete infestadas por moscas das frutas, das quais emergiram 888 espécimes de *Neosilba* spp. (Lonchaeidae) e 81 adultos de *Anastrepha* (Tephritidae). *Neosilba zadolicha* emergiu de frutos de *Siparuna camporum*, *Dyospiros hispida*, *Annona coreacea*, *Mouriri pusa*, *Cheiloclinium cognatum* e *Salacia crassifolia*. *Neosilba bifida*, *N. glaberrima* e *Neosilba* **sp.n.1** emergiram de frutos de *D. hispida*. As espécies de *Anastrepha* obtidas foram: *A. serpentina* e *A. pr. mucronota* criadas de frutos de *S. crassifolia* e *C. cognatum*; *A. obliqua* de *S. crassifolia*, *A. sororcula* de *Myrcia guianensis*, *A. turpiniae* e *A. zenildae* de *M. pusa*. Das sete espécies frutíferas hospedeiras, quatro foram infestadas por Tephritidae e seis por Lonchaeidae. Estes são os primeiros registros para: *A. sororcula* em frutos de *Myrcia guianensis*, *A. pr. mucronota* em *Salacia crassifolia* e *Cheiloclinium cognatum*, *A. serpentina* em *Cheiloclinium cognatum*, *A. turpiniae* e *A. zenildae* em *Mouriri pusa*. As associações de moscas das frutas com as frutíferas hospedeiras são localmente e temporalmente dependentes da flora, uma vez que as espécies de moscas das

frutas precisam de frutos disponíveis, como observado para a espécie altamente polífaga *A. obliqua*, que neste estudo infestou apenas uma espécie de frutífera.

Palavras Chave: Biodiversidade, Cerrado do Norte do Brasil, frugivoria, fitofagia, novos registros.

Many studies have been conducted to document biodiversity, e.g., Basset et al. (2000), Hawkins & Pausas (2004), and Lewinsohn et al. (2005). Brazil is a country with high biological diversity, and in the Amazon region most species of insects are still unknown (Strikis et al. 2012).

Tephritoid flies have great adaptability to different regions worldwide, and represent the main group of fleshy fruit consumers (White & Elson-Harris 1992). The use of host plants by individual phytophagous species is generally restricted to phylogenetically close groups, often a single family or genus (Strong et al. 1984; Bernays & Chapman 1994), but some species of fruit flies and lance flies are highly polyphagous. Both native and introduced fruits are potential host plants for larvae of frugivorous Tephritidae and Lonchaeidae (Uchôa 2012). Availability of fruits, the high biodiversity and endemism in the Cerrado biome, make these fruit trees potential sites for infestation by fruit flies, since the various fruiting patterns of these host plants assures that fruit is produced throughout the year.

Studies of the associations of species of Tephritidae and their host plants are important to understand their patterns of diversity, ecology and evolution (Aluja et al. 2003), to comprehend their natural history, and to plan strategies to manage fruit fly pests in orchards (Uramoto et al. 2004).

Larvae of *Neosilba* spp. (Lonchaeidae) develop in a wide variety of structures and plant species, including: sprouts, fruits, flowers and buds. Large populations can be found associated with mature fruits (Uchôa 2012). Current studies on *Neosilba* species include taxonomy, biology and ecology (Galeano-Olaya & Canal 2012). These native species of Lonchaeidae under human disturbance may come to use exotic cultivated plant as their hosts. Such disturbances can favor the host range expansion of phytophagous insect species beyond their primary native hosts to include cultivated exotic species (Selivon 2000).

Until the late twentieth century there were no host plant records for species of fruit flies in the State of Tocantins (Zucchi 2008). Bomfim et al. (2007a, b) published the first records. However, this information is still very rudimentary and requires additional studies to more fully elucidate the relationships between fruit flies and lance flies and their hosts, especially in natural environments where there are opportunities to observe additional native host plants.

Protected areas present one of the best strategies to conserve the natural heritage. In these

areas natural flora and fauna are preserved, as well as the ecological processes that govern ecosystems, ensuring the maintenance of the stock of biodiversity (Seplan 2005).

The Cerrado is a tropical biome with high plant diversity and endemism, where many unknown interactions between insects and their host plants should occur. Thus, considering the lack of knowledge about frugivorous larvae of Tephritoidea and their association with native host fruit species from Cerrado in the Amazon region (especially in the state of Tocantins), this study aimed to investigate species of fruit flies and lance flies and their association with host fruits in the Conservation Unit of the Parque Estadual do Lajeado, at Palmas, Tocantins, Brazil.

#### MATERIALS AND METHODS

The study was carried out in the Parque Estadual do Lajeado (PEL), an Integral Protection Conservation Unit in the municipality of Palmas Tocantins, Brazil, bounded by the coordinates S 10° 00' 24" -W 48° 17' 03", S 10° 00' 13" and -W 48° 15' 45", S 10° 11' 50" and -W 48° 12' 56", S 10° 11' 25" and -W 48° 10' 37", S 10° 09' 14" and -W 48° 09' 54". According to Seplan (2005) the area has average temperatures ranging between 25° C and 28° C; the rainfall is concentrated in a 5-6 month period, starting in Oct or Nov and lasting until Mar, with relative humidity ranging from 0 to 40%, and accumulated rainfall of about 1,700 mm per yr. Sampling was carried out from May 2011 to Apr 2013. The search for host fruits occurred biweekly throughout the sampling period, and the collection of fruits occurred according to their availability in the field. All the fruits were collected directly from the trees.

The fruits were kept in a rearing room at the Universidade Federal do Tocantins (UFT), where they were counted and weighed. Then, these fruits were arranged on fiber networks fixed in a wood platform and arranged in plastic trays containing a water slide, where they remained for 30 days to collect the frugivorous third instar larvae of Tephritoidea as they emerged from the fruit. Every 12 h the larvae were removed from the trays, placed in transparent cups (300 mL) containing sterilized sand as a pupation substrate, where the pupae remained until adult emergence occurred. The adults were reared for 3 days on a solution of honey and water (1:5 v:v), to allow full development of the color of their bodies and wings after emergence.

The species of *Neosilba* (Lonchaeidae) were identified by Dr. Laura Gislotti, Universidade Estadual de Campinas (UNICAMP), Campinas-SP, Brazil, according to McAlpine & Steyskal (1982). Voucher specimens were deposited in the Entomological Collection of UNICAMP.

The tephritids were identified in the Universidade Federal da Grande Dourados (UFGD), Dourados-MS, Brazil, using specific identification keys (Stone 1942; Steyskal 1977; Zucchi 2000; Norrbom 1997; Norrbom & Korytkowski 2009). After identification, the Tephritidae were deposited in the Coleção Entomológica do Museu da Biodiversidade (MuBio-UFGD). Dried herbarium specimens (exsiccatae) of the identified host plants were deposited in the Herbarium of Tocantins (HTO) of the Universidade Federal do Tocantins (UFT), at Porto Nacional, Tocantins, Brazil.

To analyze the species richness of fruit fly and lance fly species accumulation curves were plotted using the software EstimateS 8.2 (Colwell 2009) with 100 randomizations. Each point of the curve corresponds to the average of the accumulated species richness in the 100 curves. Each species of fruit collected was considered a sample, totaling 18 samples. The species richness was cal-

culated by the estimators Jackknife 2, and Chao 2. The similarity between species of fruit flies and lance flies regarding exploitation of their host fruits was calculated through cluster analysis using the Bray-Curtis distance by the software Biodiversity Pro 2 (McAleece et al. 1997).

## RESULTS

In total 2,682 fruits of 18 species of native plants were sampled. Seven plant species were infested by the fruit fly and lance fly species. These included 888 *Neosilba* (Lonchaeidae) adults recovered, which consisted of the species *N. zado-lichia*, *N. pendula*, *N. bifida*, *N. glaberrima*, and *Neosilba* n.sp.1, and 81 adults of *Anastrepha* spp. (Tephritidae): *A. serpentina*, *A. obliqua*, *A. nr. mucronota*, *A. sororcula*, *A. turpiniae* and *A. zenildae* (Table 1).

Four of the 7 host species were infested by Tephritidae and 6 by Lonchaeidae. *Anastrepha serpentina* was recovered from *Salacia crassifolia* and *Cheiloclinium cognatum* (Hippocrateaceae), *A. obliqua* from *S. crassifolia*; *A. nr. mucronota* from *S. crassifolia* and *C. cognatum*; *A. sororcula* from *Myrcia guianensis* (Myrtaceae); and *A. tur-*

TABLE 1. SPECIES OF FRUIT FLIES AND LANCE FLIES (DIPTERA: TEPHRITOIDEA) AND THEIR NATIVE HOST FRUITS IN THE CONSERVATION UNIT PARQUE ESTADUAL DO LAJEADO, TO, BRAZIL (MAY 2011 TO APR 2013).

Fruit Fly Species	Fruit Tree Species	No. adults	Herbarium of Tocantins (HTO) record No.
<i>A. serpentina</i>	<i>Salacia crassifolia</i> (Mart. ex Schult.) G. Don*	15	6,312 HTO
	<i>Cheiloclinium cognatum</i> (Miers) A. C. Sm.****	10	8,427 HTO
<i>A. obliqua</i>	<i>Salacia crassifolia</i> (Mart. ex Schult.) G. Don**	10	6,312 HTO
<i>A. nr.mucronota</i>	<i>Salacia crassifolia</i> (Mart. ex Schult.) G. Don****	1	6,312 HTO
	<i>Cheiloclinium cognatum</i> (Miers) A. C. Sm.****	1	8,427 HTO
<i>A. sororcula</i>	<i>Myrcia guianensis</i> (Aubl.) DC.****	12	10,513 HTO
<i>A. turpiniae</i>	<i>Mouriri pusa</i> Gardner****	31	4,575 HTO
<i>A. zenildae</i>	<i>Mouriri pusa</i> Gardner****	1	4,575 HTO
<i>Neosilba zado-lichia</i>	<i>Siparuna camporum</i> (Tul.) A.DC.**	6	3,429 HTO
	<i>Diospyros hispida</i> A.DC.**	826	10,565 HTO
	<i>Annona coriacea</i> (Mart.)**	36	5,096 HTO
	<i>Mouriri pusa</i> Gardner**	1	4,575 HTO
	<i>Salacia crassifolia</i> (Mart. ex Schult.) G. Don**	14	6,312 HTO
	<i>Cheiloclinium cognatum</i> (Miers) A. C. Sm.**	1	8,427 HTO
<i>N. pendula</i>	<i>Salacia crassifolia</i> (Mart. ex Schult.) G. Don	1	6,312 HTO
<i>Neosilba bifida</i> ***	<i>Diospyros hispida</i> A.DC.	1	10,565 HTO
<i>Neosilba glaberrima</i>	<i>Diospyros hispida</i> A.DC.**	1	10,565 HTO
<i>Neosilba n. sp.1</i>	<i>Diospyros hispida</i> A.DC.	2	10,565 HTO

\*First record in the host for Tocantins;

\*\*First record in the host in northern Brazil;

\*\*\*First record of this lance fly species in northern Brazil;

\*\*\*\*First record of these fruit fly species in this host.

*piniae* and *A. zenildae* from *Mouriri pusa* (Memecylaceae).

Five species of Lonchaeidae were recovered as follows: *Neosilba zadolicha* from fruits of *Siparuna camporum* (Siparunaceae), *Diospyros hispida* (Ebenaceae), *Annona coriacea* (Anonaceae), *M. pusa*, *S. crassifolia* and *C. cognatum*; *N. pendula* from fruit of *S. crassifolia*; and *N. bifida*, *N. glaberrima*, and *Neosilba* n. sp.1 from *D. hispida* fruits. The greatest number of species of fruits (6) were infested by *N. zadolicha*, the most polyphagous and abundant species (Table 1).

The season of greatest fruit production was from Oct to Mar. *Salacia crassifolia* and *C. cognatum* were the only host fruits simultaneously infested by larvae of *Neosilba* spp. and *Anastrepha* spp.

The host fruits with the highest rates of infestation, taking in account all Tephritoidea, were: *S. crassifolia*, *C. cognatum* (Hippocrateaceae), and *D. hispida* (Ebenaceae). The infestation rate by larvae of *Anastrepha* species (Tephritidae) was highest in the following host fruits: *C. cognatum* (32.32 larvae/kg), *S. crassifolia* (30.01 larvae/kg), and *Myrcia guianensis* (8.59 larvae/kg). Fruits of *M. guianensis* were 100% infested by *Anastrepha sororcula*. For the species of *Neosilba* (Lonchaeidae) the greatest rate of infestation occurred in *M. pusa* (149.35 larvae/kg), *D. hispida* (123.10 larvae/kg), and *A. coriacea* (36.73 larvae/kg). One hundred percent of fruits of *S. camporum*, *D. hispida*, and *A. coriacea* were infested by species of *Neosilba* (Table 2).

The species accumulation curves were upward, and the curves did not indicate stabilization (Fig. 1). Cluster analysis generated 4 groups based on the use of host fruits: 2 of them formed by *Anastrepha* species, 1 by *Neosilba* species, and a fourth group by *A. obliqua* and *N. pendula* (Fig. 2).

## DISCUSSION

The species accumulation curves (Fig. 1) suggest the possibility of finding more species of fruit fly and lance fly species in the fruit species surveyed, if the sampling had been continued. The species richness of frugivorous flies could have been enhanced if we had increased the number of fruit samples from the fruit species evaluated. The estimators Jackknife 2 predicted a species richness of 24, and the Chao 2 estimated a maximum of 27 species for the samples actually made; these estimates are higher than the number of species recovered from fruits sampled in this inventory ( $n = 10$ ) (Fig. 1).

The similarity in the use of host plant taxa between the different frugivorous fly species (cluster analysis) in this study showed that species of *Neosilba* were more similar to each other than with the species of *Anastrepha*, with the exception of the similarity between *N. pendula* and *A.*

*obliqua* (Fig. 2). Interestingly the latter 2 species are widely polyphagous, as pointed out by Uchôa (2012), but herein they attacked a single species of fruit (*S. crassifolia*, Table 1). This may have occurred due the effect of a fire that burned the vegetation (Bomfim & Uchôa unpublished data), probably reducing the populations of the 2 fly species.

The choice of the host plant by insects is not the result of a simple behavior, but represents a dynamic hierarchy of various components, including co-evolutionary aspects (Connell 1980; Aluja & Mangan 2008). The use of host plants may vary at the individual level, in response to changes in the internal physiological conditions of the fly (e.g., the load of eggs) caused by a shortage of preferred hosts or prior experience with a host. In the absence of their preferred host plant, a polyphagous insect may exhibit hierarchies of preference among those currently available. The usage pattern of host plants can vary according to the relative abundance of potential alternate host(s). It is expected that the diversity of fruit trees in the natural forest increases the likelihood of finding monophagous species of fruit flies (Selivon 2000; Aluja & Mangan 2008). Therefore, the association with the host fruit is a local phenomenon and depends on host availability in the ecosystem, as in the cases of *Anastrepha obliqua* and *N. pendula* (Araujo & Zucchi 2002); a single species of fruit was infested by the latter 2 species in this survey, despite both species being polyphagous species.

Sixteen interactions between species of fruit flies and lance flies and their host plants sampled in this study were discovered and reported here. This is the consequence of the floristic richness in the Cerrado Biome, and the lack of studies on trophic interactions between Tephritoidea and host plants in the Neotropics. According to Zucchi (2008), the hosts of about 48.70% of *Anastrepha* species reported in Brazil are still unknown.

Prior to this study, host plants were known in the State of Tocantins for the following nine species of fruit flies: *A. coronilli*, *A. fraterculus*, *A. nr. mucronota*, *A. obliqua*, *A. sororcula*, *A. striata*, *A. turpiniae*, *A. zenildae*, and *C. capitata* (Bomfim et al. 2007b). This research has added new associations with hosts for: *A. nr. mucronota*, *A. serpentina*, *A. sororcula*, *A. turpiniae*, and *A. zenildae*. Also this research has associated 5 species of *Neosilba* with their host fruits in northern Brazil (Table 1). These results highlight the importance of inventories of frugivorous tephritoids in native host plants, because fruit production by the native plants is interspersed throughout the yr, which provides conditions for the survival of fruit flies. The existence of a wide range of hosts for these flies in Tropical America, with fruit production in different seasons, provide conditions for maintenance of high populations of frugivorous Tephritoidea.



TABLE 2. LEVELS OF INFESTATION AND RESOURCE PARTITIONING BY FRUIT FLY AND LANCE FLY SPECIES (DIPTERA: TEPHRITIDAE AND LONCHAEIDAE) IN SOME NATIVE HOST FRUIT SPECIES FROM THE CONSERVATION UNIT PARQUE ESTADUAL DO LAJEADO, TO, BRAZIL (MAY 2011 TO APR 2013).

Species of host fruits	No. of Larvae	No. of Fruits	Biomass of fruit (g)	Larvae/fruit	Larvae/kg of fruit	<i>Anastrepha</i> spp. (%)	<i>Neosilba</i> spp. (%)
<i>Salacia crassifolia</i> (Hippocrateaceae)	114	101	3799	1.13	30.01	71	29
<i>Cheiloclinium cognatum</i> (Hypocrateaceae)	85	115	2630	0.74	32.32	92	08
<i>Myrcia guianensis</i> (Myrtaceae)	28	144	326	0.19	8.59	100	00
<i>Siparuna camporum</i> (Siparunaceae)	18	1952	5957	0.01	3.02	00	100
<i>Diospyros hispida</i> (Ebenaceae)	1101	256	8944	4.30	123.10	00	100
<i>Annona coriacea</i> (Anonaceae)	90	5	2450	18.00	36.73	00	100
<i>Mouriri pusa</i> (Memecylaceae)	115	109	770	1.06	149.35	99	01
<i>Alibertia edulis</i> Rich (Rubiaceae)	0	41	1482	0	0	0	0
<i>Anacardium occidentale</i> L. (Anacardiaceae)	0	82	2420	0	0	0	0
<i>Brosimum gaudichaudii</i> Tréc. (Moraceae)	0	116	940	0	0	0	0
<i>Caryocar brasiliense</i> Camb. (Caryocaraceae)	0	84	7405	0	0	0	0
<i>Couepia grandiflora</i> Benth. (Chrysobalanaceae)	0	31	362	0	0	0	0
<i>Hancornia speciosa</i> Gomes (Apocynaceae)	0	8	210	0	0	0	0
<i>Mauritia flexuosa</i> L.f. (Arecaceae)	0	72	3744	0	0	0	0
<i>Byrsonima</i> sp. (Malpighiaceae)	0	140	2289	0	0	0	0
<i>Leandra</i> sp. (Melastomataceae)	0	132	330	0	0	0	0
<i>Passiflora</i> sp. (Passifloraceae)	0	6	241	0	0	0	0
<i>Psidium</i> sp. (Myrtaceae)	0	17	280	0	0	0	0
Total	1,551	3,411	44,579	—	—	—	—




Fig. 1. Accumulation curves for fruit fly and lance fly species (Tephritidae and Lonchaeidae) in the host fruits calculated using Sobs, Chao 2 and Jack 2 estimators, sampled in the Conservation Unit of Parque Estadual do Lajeado, Palmas, to, Brazil (May 2011 to Apr 2013).

Research on fruit flies in other natural reserves in Brazil has revealed important aspects of diversity and trophic interactions with host plants and parasitoids. In a study in the Adolpho Ducke Reserve in Amazonas State, 12 species of *Anastrepha* were found, of which eight were new species (Costa 2005). In Conservation Unit Vale do Rio Doce (Atlantic Forest), State of Espírito Santo, 21 species of *Anastrepha* were sampled, and 5 of these were new records for Espírito Santo, and one was a new record for Brazil. In that study 5 new species were found (Uramoto et al. 2008). So, each survey carried out in natural environments in the Neotropical Region has increased new information on Tephritoidea diversity, their host plants and natural enemies.

#### New Host Fruits of Tephritidae

*Anastrepha serpentina* was reared for the first time from fruits of *Cheiloclinium cognatum*; *A. nr. mucronota* from *Salacia crassifolia* and *C. cognatum*; *A. turpiniae* and *A. zenildae* from *Mouriri pusa* fruits. Here we also reported the first record for *A. sororcula* in fruits of *M. guianensis*, a small wild fruit with a diam of 5 mm.

*A. turpiniae* and *A. zenildae* from *Mouriri pusa* fruits. Here we also reported the first record for *A. sororcula* in fruits of *M. guianensis*, a small wild fruit with a diam of 5 mm.

#### New host fruit of Lonchaeidae

All records of host plants for the species of *Neosilba* in this survey are new associations. The species of *Neosilba* were more abundant and colonized more species of fruit plants than did the species of *Anastrepha*. Other studies have also highlighted the importance of some Lonchaeidae species as primary fruit infesting species, and that several species of *Dasiops*, and *Neosilba* are voracious omnivores (Uchôa et al. 2002; Nicácio & Uchôa 2011; Uchôa 2012).

In all states of the northern region of Brazil there are reports of the occurrence of *Neosilba* species, i.e., Amazonas, Amapá, Acre, Pará, Rondônia, Roraima (Strikis et al. 2011), and Tocantins (Bomfim et al. 2007a, b.). Recent studies indicate that some species of *Neosilba* colonize




Fig. 2. Analysis of similarity based in the use of host fruits by species of fruit and lance flies (Tephritidae and Lonchaeidae) in the Conservation Unit of Parque Estadual do Lajeado, Palmas-to, Brazil (May 2011 to Apr 2013).

a greater number of host fruit species than the species of *Anastrepha* (Uchôa et al. 2002; Uchôa 2012), and this demonstrates the importance and the need for investigations on the Lonchaeidae in the Neotropical Region.

In this research *Neosilba zadolicha* infested a greater diversity of host fruits compared to all other obtained species of frugivorous flies. This polyphagous species has already been reported from 17 host species (Strikis et al. 2011; Uchôa 2012). The occurrences reported herein of *N. zadolicha* in the host plants: *S. camporum*, *D. hispida*, *A. coriacea*, *M. pusa*, *S. crassifolia* and *C. cognatum*, are new records. *Neosilba pendula* and *N. glaberrima* were individually recorded each in a single host (*Salacia crassifolia* and *Diospyros hispida*, respectively - Table 1). However, both species are considered polyphagous (Uchôa 2012). Araújo & Zucchi (2002) reported the polyphagy of *N. pendula*, and that this species indiscriminately infested native and exotic fruits.

Due to high levels of infestation in native and exotic fruits, lonchaeids are currently considered of economic importance in Brazil (Uchôa et al. 2002; Uchôa 2012; Strikis et al. 2011; Strikis et al. 2012; Gislotti & Prado 2012). In this work,

the larvae of some species of *Neosilba* colonized higher number of fruit species than *Anastrepha*. This pattern was also observed before in areas of the Cerrado and the Pantanal (Uchôa et al. 2002; Uchôa & Nicácio 2010). The lonchaeids have been ignored in most studies of frugivorous flies in Brazil, possibly due to lack of taxonomic knowledge.

The fruits of *Diospyros hispida* were colonized only by the species of *Neosilba* (Table 2), suggesting a probable specificity of lonchaeids for this host plant. According to Aluja & Mangan (2008), an alternative explanation for unexpected patterns in the usage of host plants by insects in the field could involve associative learning, and in that case, there is no need for changes in larval characteristics.

#### Resource Partitioning by Fruit Flies

Sharing of resources between species of *Neosilba* and those of *Anastrepha* was here observed, both groups (*Anastrepha* species and *Neosilba* species) infested *S. crassifolia* and *C. cognatum*. A similar pattern was observed by Uchôa et al. (2002) who found infestations by species of these



2 genera in *Citrus* spp. (Rutaceae). Braga-Filho et al. (2001) also found that fruits of *S. crassifolia* were natural hosts for species of *Anastrepha* and *Neosilba*, with concomitant infestation. According to Connell (1980), niches occupied by competitors in ecological communities are shaped by mutual co-evolution, which allows many species to coexist.

#### ACKNOWLEDGMENTS

The authors are grateful to CAPES-Coordenação de Aperfeiçoamento de Pessoal de Nível Superior by scholarship for DAB, and to CNPq-Conselho Nacional de Desenvolvimento Científico e Tecnológico by a grant based on research productivity to MAU.

#### REFERENCES CITED

- ALUJA, M., AND MANGAN, R. L. 2008. Fruit fly (Diptera: Tephritidae) host status determination: Critical conceptual, methodological, and regulatory considerations. *Ann. Rev. Entomol.* 53: 473-502.
- ALUJA, M., RULL, J., SIVINSKI, J., NORRBOM, A. L., WHARTON, R. A., MACIAS-ORDÓÑEZ, R., DIAZ-FLEISCHER, F., AND LÓPEZ, M. 2003. Fruit flies of the genus *Anastrepha* (Diptera: Tephritidae) and associated native parasitoids (Hymenoptera) in the tropical rainforest Biosphere Reserve of Montes Azules, Chiapas, Mexico. *Environ. Entomol.* 32: 1377-1385.
- ARAUJO, E. L., AND ZUCCHI, R. A. 2002. Hospedeiros e níveis de infestação de *Neosilba pendula* (Bezzi) (Diptera: Lonchaeidae) na região de Mossoró/Assu, RN. *Arq. Inst. Biol.* 69: 91-94.
- BASSET, B. Y., NOVOTNY, V., MILLER, S. E., AND PYLE, R. 2000. Quantifying biodiversity: Experience with parataxonomists and digital photography in Papua New Guinea and Guyana. *BioScience* 50: 899-908.
- BERNAYS, E. A., AND CHAPMAN, R. F. 1994. Host-plant specialization by phytophagous insects. Chapman & Hall, London. 312 pp.
- BOMFIM, D. A., UCHÔA, M. A., AND BRAGANÇA, M. A. L. 2007a. Biodiversidade de moscas-das-frutas (Diptera, Tephritoidea) em matas nativas e pomares domésticos de dois municípios do estado do Tocantins, Brasil. *Rev. Brasileira Entomol.* 51: 217-233.
- BOMFIM, D. A., UCHÔA, M. A., AND BRAGANÇA, M. A. L. 2007b. Hosts and parasitoids of fruit flies (Diptera: Tephritoidea) in the State of Tocantins, Brazil. *Neotrop. Entomol.* 36: 984-986.
- BRAGA-FILHO, J. R., VELOSO, V. R. S., NAVES, R. V., AND FERREIRA, G. A. 2001. Entomofauna associada aos frutos do bacupari, *Salacia crassifolia* (MART.) Peyr, nos cerrados do Brasil Central. *Pesq. Agrop. Trop.* 31: 47-54.
- CONNELL, J. H. 1980. Diversity and the coevolution of competitors, or the ghost of competition past. *Oikos* 35: 131-138.
- COLWELL, R. K. 2009. EstimateS: statistical estimation of species richness and shared species from samples. Version 8.2. User's guide and application published at: <http://purl.oclc.org/estimates>. Accessed 15-IV-2014.
- COSTA, S. G. M. 2005. Himenópteros parasitóides de larvas frugívoras (Diptera: Tephritoidea) na reserva florestal Adolpho Ducke, Manaus, Amazonas, Brasil. Dissertação de Mestrado. Instituto Nacional de Pesquisas da Amazônia, Universidade Federal do Amazonas, Manaus. 102 pp.
- GALEANO-OLAYA, P. E., AND CANAL, N. A. 2012. New species of *Neosilba* McAlpine (Diptera: Lonchaeidae) and new records from Colombia. *Pap. Avul. Zool. MZUSP* 52: 361-385.
- GISLOTI, L. J., AND PRADO, A. P. 2012. Parasitism of *Neosilba perezi* (Diptera: Tephritidae) larvae by a braconid, *Phaenocarpa neosilba* (Hymenoptera: Braconidae: Alysiinae). *Florida Entomol.* 95: 900-904.
- HAWKINS, B. A., AND PAUSAS, J. G. 2004. Does plant richness influence animal richness?: The mammals of Catalonia (NE Spain). *Divers. Distrib.* 10: 247-252.
- LEWINSOHN, T. M., NOVOTNY, V., AND BASSET, Y. 2005. Insects on plants: Diversity of herbivore assemblages revisited. *Annu. Rev. Ecol. Evol. Syst.* 36: 597-620.
- MCALLECE, N., LAMBSHEAD, P. J. D., PATERSON, G. L. J., AND GAGE, J. G. 1997. Biodiversity professional. Beta-Version. London, The Nat. Hist. Mus. and the Scottish Assoc. for Marine Sci. <http://biodiversity-pro.software.informer.com/2.0/> Accessed 5-IX-2013.
- MCALPINE, J. F., AND STEYSKAL, G. C. 1982. A revision of *Neosilba* McAlpine with a key to the world genera of Lonchaeidae (Diptera). *Canadian Entomol.* 114: 105-137.
- NICÁCIO, J., AND UCHÔA, M. A. 2011. Diversity of frugivorous flies (Diptera: Tephritidae and Lonchaeidae) and their relationship with host plants (Angiospermae) in environments of South Pantanal Region, Brazil. *Florida Entomol.* 94: 443-46.
- NORRBOM, A. L. 1997. Revision of the *Anastrepha benjamini* species group and the *A. pallidipennis* complex (Diptera: Tephritidae). *Insecta Mundi* 11: 141-157.
- NORRBOM, A. L., AND KORYTKOWSKI, C. A. 2009. A revision of the *Anastrepha robusta* species group (Diptera: Tephritidae). *Zootaxa* 2182: 1-91.
- SELIVON, D. 2000. Biologia e padrões de especiação, pp. 25-29 *In* A. Malavasi and R. A. Zucchi [eds.], *Moscas-das-frutas de importância econômica no Brasil: conhecimento básico e aplicado*. Editora Holos, Ribeirão Preto, 327 pp.
- SEPLAN-SECRETARIA DO MEIO AMBIENTE E PLANEJAMENTO. 2005. Plano de manejo do parque Estadual do Lajeado. Goiania. 286p. <<http://www.gesto.to.gov.br/uc/50/documentos/>>. Accessed 7-IV-2011.
- STEYSKAL, G. 1977. Pictorial key to species of the genus *Anastrepha* (Diptera: Tephritidae). *Entomol. Soc. Washington, Washington, DC*. 35 pp.
- STONE, A. 1942. The fruit flies of the genus *Anastrepha*. U. S. Dept. Agric. Misc. Publ. Washington, DC, USDA. 112 pp.
- STRIKIS, P. C., DEUS, E. G., SILVA, R. A., PEREIRA, J. D. B., JESUS, C. R., AND MARSARO JÚNIOR, A. L. 2011. Conhecimento sobre Lonchaeidae na Amazônia brasileira, p. 205-215 *In* Silva, R. A., Lemos, W. P., and Zucchi, R. A. (eds), *Moscas-das-Frutas na Amazônia brasileira: Diversidade, hospedeiros e inimigos naturais*. Embrapa, Amapá, 299 pp.
- STRIKIS, P. C., MARSARO JÚNIOR, A. L., ADAIME, R., AND LIMA, C. R. 2012. First report of infestation of cassava fruit, *Manihot esculenta*, by *Neosilba perezi* (Romero & Ruppell) (Lonchaeidae) in Brazil. *Brazilian J. Biol.* 72: 631-632.

- STRONG, D. R. J., LAWTON, J. H., AND SOUTHWOOD, T. R. E. 1984. Insects on plants: Community patterns and mechanisms. Oxford: Blackwell. 313 pp.
- UCHÔA, M. A. 2012. Fruit Flies (Diptera: Tephritoidea): Biology, host plants, natural enemies, and the implications to their natural control. pp. 271-300 *In* M. L. Larramendy and S. Soloneski, S. [eds.], Integrated Pest Management and Pest Control: Current and future tactics. InTech. Rijeka, Croatia. 659 pp.
- UCHÔA, M. A. AND NICÁCIO, J. 2010. New records of neotropical fruit flies (Tephritidae), lance flies (Lonchaeidae) (Diptera: Tephritoidea), and their host plants in the South Pantanal and adjacent areas, Brazil. *Ann. Entomol. Soc. America* 103: 723-733.
- UCHÔA, M. A., OLIVEIRA, I., MOLINA, R. M. S., AND ZUCCHI, R. A. 2002. Species diversity of frugivorous flies (Diptera: Tephritoidea) from hosts in the Cerrado of the state of Mato Grosso do Sul, Brazil. *Neotrop. Entomol.* 31: 515-524.
- URAMOTO, K., MARTINS, D. S., AND ZUCCHI, R. A. 2008. Fruit flies (Diptera, Tephritidae) and their association with native host plants in a remnant area of the highly endangered Atlantic Rain Forest in the State of Espírito Santo, Brazil. *Bull. Entomol. Res.* 98: 457-466.
- URAMOTO, K., WALDER, J. M. M., AND ZUCCHI, R. A. 2004. Biodiversidade de moscas-das-frutas do gênero *Anastrepha* (Diptera, Tephritoidea) no campus da ESALQ - USP, Piracicaba, São Paulo. *Rev. Brasileira Entomol.* 48: 409-414.
- WHITE, I. M., ELSON-HARRIS, M. M. 1992. Fruit flies of economic significance: Their identification and bio-nomics. CAB International, Wallingford. 601 pp.
- ZUCCHI, R. A. 2000. Taxonomia, pp. 13-24 *In* A. Malavasi and R. A. Zucchi [eds.], *Moscas-das-frutas de importância econômica no Brasil: conhecimento básico e aplicado*. Editora Holos, Ribeirão Preto, 327 pp.
- ZUCCHI, R. A. 2008. Fruit flies in Brazil - *Anastrepha* species their host plants and parasitoids. Available at [www.lea.esalq.usp.br/anastrepha/](http://www.lea.esalq.usp.br/anastrepha/), updated on 20-III-2014. Accessed on 21-IV-2014.