

Mosquito Surveillance in Mexico: The Use of Ovitrap for *Aedes aegypti*, *Ae. albopictus*, and Non-Target Species

Authors: Ortega-Morales, Aldo I., Moreno-García, Miguel, González-Acosta, Cassandra, and Correa-Morales, Fabián

Source: Florida Entomologist, 101(4) : 623-626

Published By: Florida Entomological Society

URL: <https://doi.org/10.1653/024.101.0425>

BioOne Complete (complete.BioOne.org) is a full-text database of 200 subscribed and open-access titles in the biological, ecological, and environmental sciences published by nonprofit societies, associations, museums, institutions, and presses.

Your use of this PDF, the BioOne Complete website, and all posted and associated content indicates your acceptance of BioOne's Terms of Use, available at www.bioone.org/terms-of-use.

Usage of BioOne Complete content is strictly limited to personal, educational, and non-commercial use. Commercial inquiries or rights and permissions requests should be directed to the individual publisher as copyright holder.

BioOne sees sustainable scholarly publishing as an inherently collaborative enterprise connecting authors, nonprofit publishers, academic institutions, research libraries, and research funders in the common goal of maximizing access to critical research.

Mosquito surveillance in Mexico: the use of ovitraps for *Aedes aegypti*, *Ae. albopictus*, and non-target species

Aldo I. Ortega-Morales¹, Miguel Moreno-García^{2,3,*}, Cassandra González-Acosta³, and Fabián Correa-Morales³

Abstract

Vector-borne diseases are of great public health concern in Mexico. Timely surveillance and control measures require methods to quickly detect mosquito population fluctuations. Surveillance is important also to detect the establishment of uncommon mosquito species. Ovitrap have been a convenient way to detect the presence of female mosquitoes. For these reasons, the Mexican Ministry of Health adopted a mosquito surveillance program using ovitraps in the country. In this study, we present the checklist of 12 target and non-target species that have been collected nationwide in ovitraps since 2013. We include distributional and ecological notes as well as the medical importance of each species. Our results showed that ovitraps can be used to estimate mosquito diversity in urban and semi-urban areas.

Key Words: mosquito control; diversity; collecting method; mosquito-borne diseases

Resumen

Las enfermedades transmitidas por vectores son una gran preocupación para la salud pública en México. Las medidas oportunas de vigilancia y control requieren métodos para detectar rápidamente las fluctuaciones de la población de mosquitos. La vigilancia también es importante para detectar el establecimiento de especies de mosquitos poco comunes. Las ovitrampas han sido una forma conveniente de detectar la presencia de mosquitos hembras. Por estas razones, el Ministerio de Salud de México adoptó un programa de vigilancia de mosquitos usando ovitrampas en el país. En este estudio presentamos la lista de verificación de doce especies blanco y no blanco que se han recolectado a nivel nacional en ovitrampas desde el 2013. Incluimos notas de distribución y ecológicas, así como la importancia médica de cada especie. Nuestros resultados mostraron que las ovitrampas se pueden usar para estimar la diversidad de mosquitos en áreas urbanas y semiurbanas.

Palabras Clave: control de mosquitos; diversidad; método de recolección; enfermedades transmitidas por mosquitos

Vector-borne diseases, such as dengue, chikungunya, and Zika are of great public health concern in Mexico. During 2014 to 2017, the Mexican Ministry of Health reported 91,836 confirmed cases of dengue; 13,564 cases of chikungunya; and 11,865 cases of Zika (DGE 2017). The consistently high number of vector-borne disease cases has made mosquito vector control a primary priority in the country. Control programs include governmental planned strategies and community participation such as “National days of elimination of breeding,” “Clean backyard,” and “Stored water management” campaigns.

In Mexico, *Aedes aegypti* (L.) and *Aedes albopictus* Skuse (both Diptera: Culicidae) are the main vectors of vector-borne diseases. Previous studies have reported *Ae. aegypti* present in 28 states of Mexico (Ibáñez-Bernal & Martínez-Campos 1994a; Mora-Covarrubias et al. 2010; Ávila-Rodríguez et al. 2012; Espinoza-Gómez et al. 2013; Kuri-Morales et al. 2017). In addition, *Ae. albopictus* has been reported in 12 states of the country (Francy et al. 1990; Ibáñez-Bernal & Martínez-Campos 1994b; Pesina et al. 2001; Flisser et al. 2002; Martínez & Estrada 2003; Villegas-Trejo et al. 2010; Salomón-Grajales et al. 2012;

Torres-Avenidaño et al. 2015; Ortega-Morales et al. 2016; Ortega-Morales & Siller-Rodríguez 2016).

For both vectors, timely surveillance and control measures in endemic or high-risk locations for disease transmission requires sensitive methods to quickly detect changes in mosquito abundance. Surveillance is important also to detect the establishment of uncommon or invasive species in regions where they have not been reported (Hernandez-Avila et al. 2013; Chaverri et al. 2018). Urban and semi-urban larval habitats of *Ae. aegypti* and *Ae. albopictus* usually are comprised of a variety of containers. Natural containers include water-filled tree holes, bamboo internodes, and bromeliad axils (Ceretti-Junior et al. 2014; Docile et al. 2017). *Aedes aegypti* prefer artificial containers such as small cisterns, discarded tires, flower vases, buckets, and cans (Cheong 1967; Yee et al. 2010; Estallo et al. 2011).

Given the biological specificity for aquatic habitats, the collection of eggs and larvae using ovitraps has been a convenient way for detecting the presence of adult female mosquitoes. Ovitrap do not require electrical power sources, additional carbon dioxide (Snetselaar et al. 2014), or daily monitoring. These traps are safe, reusable, and inex-

¹Departamento de Parasitología, Universidad Autónoma Agraria Antonio Narro Unidad Laguna, Periférico Raúl López Sánchez y carretera a Santa Fe, C.P. 27300, Torreón, Coahuila, Mexico; E-mail: agrortega@hotmail.com (A. I. O. M.)

²Unidad de Bioensayos-Centro Regional de Control de Vectores Panchimalco, Servicios de Salud de Morelos, Emiliano Zapata 95, Jojutla, C.P. 62900, Morelos, Mexico; E-mail: miguelmoga2000@yahoo.com.mx (M. M. G.)

³Centro Nacional de Programas Preventivos y Control de Enfermedades, Benjamín Franklin 132, Col. Escandón, C.P. 11800, Mexico City, Mexico; E-mail: fabiancorrea@msn.com (F. C. M.), cgonzalez_vectores@hotmail.com (C. G. A.)

*Corresponding author; E-mail: miguelmoga2000@yahoo.com.mx

pensive (CDC 2017). For these reasons, ovitraps are one of the most practical tools for surveillance of container-inhabiting *Aedes* (Wu et al. 2013; Codeço et al. 2015; Chaverri et al. 2018).

Material and Methods

In 2013, the Mexican Ministry of Health implemented a mosquito surveillance program using ovitraps. The ovitraps used by the national surveillance program are 1 L black colored plastic cups, filled with tap water, and lined with a strip of filter paper along the water margin where female *Aedes* species lay their eggs. Every year, 250,000 ovitraps are placed in 712 semi-urban and urban locations (with or without active presence of dengue, Zika, or chikungunya cases) covering all 32 states of the country (including Mexico City). Ovitrap are placed outdoors either at randomly selected houses (after obtaining informed consent) or nonresidential areas (CENAPRECE 2017). Ovitrap are monitored weekly, when the water is changed, and the paper strip from each positive ovitrap is collected by public health technicians and transported to the laboratory. Eggs are hatched and reared to the fourth instar, and identified using the morphological identification keys of Schick (1970), and Clark-Gil & Darsie (1983). In this study, we present a checklist of mosquito species that have been collected nationwide in ovitraps since 2013. We include distributional and ecological notes, as well as the medical importance of each species.

Results and Discussion

Aedes (Georgecraigius) epactius Dyar & Knab (Diptera: Culicidae): *Aedes epactius* was the most commonly collected species in ovitraps from Mexico City, and less frequently in the states of Morelos and Puebla. This is a common mosquito species in arid and semi-arid regions of Mexico, including urban and suburban areas. In the Mexico Valley area (including the states surrounding Mexico City: Estado de Mexico, Hidalgo, Morelos, Puebla, Queretaro, and Tlaxcala), *Ae. epactius* generally appears in spring and early summer. Populations usually decrease when the rainy season starts. *Aedes epactius* also is known as the rock-hole mosquito where its natural developmental habitats are water filled rock cavities (Zavortink 1972). In Mexico, immature stages of this species have been found in a wide variety of artificial containers such as buckets, discarded tires, flower vases, and watering canals for cattle. Commonly, this mosquito prefers clean water with limited leaf material. Adults are crepuscular and are most active at dusk. Females are highly anthropophagous and usually rest indoors in dark, warm places. Although *Ae. epactius* is highly anthropophagous, its role as a possible vector of human disease is currently unknown.

Aedes (Protomacleaya) gabriel Schick (Diptera: Culicidae): This species was collected only from ovitraps in the subtropical forest of Sierra Madre del Sur in southern Morelos. Although *Ae. gabriel* is historically considered endemic to Mexico, it has only been reported in the states of Jalisco, Morelos, and Zacatecas (Schick 1970; Henimeann & Belkin 1977). This species has been poorly studied, so the true abundance of *Ae. gabriel* is unknown in the regions where it occurs. This is especially true during the rainy season when rainwater filled tree holes and rock holes become larval habitats. Adults are diurnal and females have been collected when they attempt to feed on humans. The medical importance of *Ae. gabriel* is unknown.

Aedes (Protomacleaya) podographicus Dyar & Knab (Diptera: Culicidae): *Aedes podographicus* was collected in ovitraps from the states of Chiapas and Queretaro, constituting the first record in Queretaro. This is one of the most common mosquito species in the tropi-

cal regions of Mexico, and the most common species of the subgenus *Protomacleaya*. Larval habitats usually include natural water-holding habitats such as tree holes. However, *Ae. podographicus* also has been found in artificial containers such as discarded tires and flower vases, with murky water and abundant organic material at the bottom of the aquatic habitat site (Schick 1970). Adults are diurnal and females are active at ground level and in the tree canopy. They can feed on humans and are persistent biters even when the host is moving. Although *Ae. podographicus* has not been incriminated as a vector of human disease, its preferential feeding on humans and monkeys could have potential importance in the transmission of diseases such as yellow fever. For surveillance of presence of this disease in southeastern Mexico, we placed adult mosquito traps in the tree canopy using spider monkeys (*Ateles geoffroyi* Kuhl [Atelidae]) as bait. Large numbers of *Ae. podographicus* were collected together with *Sabethes chloropterus* (Humboldt) and *Haemagogus* spp. (both Diptera: Culicidae). However, no yellow fever virus infection was detected in any of the mosquitoes collected in those traps.

Aedes (Stegomyia) aegypti (L.): This species is the most medically important mosquito in Mexico, as it is the main vector of dengue, Zika, and chikungunya. The national mosquito surveillance program is focused mainly on this species. In the wild, immature stages of *Ae. aegypti* have been found in tree holes. In urban and semi-urban locations, it is commonly found in artificial containers such as discarded tires, buckets, flower vases, and any other domestic container able to collect rain water. Adults are diurnal and usually feed and rest in indoor environments.

Aedes (Stegomyia) albopictus (Skuse): In this study, *Ae. albopictus* was collected from ovitraps in the states of Chiapas, Hidalgo, Mexico City (A.I.O.-M. unpublished data), Morelos, Nuevo León, San Luis Potosí, and Veracruz. Previously, this species has been collected commonly from ovitraps from the states of Chiapas, Coahuila, Hidalgo, Morelos, Nuevo León, Quintana Roo, San Luis Potosí, Sinaloa, Tamaulipas, and Veracruz (Ortega-Morales & Siller-Rodríguez 2016). In our current study, *Ae. albopictus* was collected in Mexico City, confirming its presence in Mexico's capital. Because this species has been incriminated as an arbovirus vector, *Ae. albopictus* is considered the second most medically important mosquito species in Mexico. Immature stages of *Ae. albopictus* are found commonly in a variety of artificial and natural containers with clean water, such as tree holes and bamboo internodes. This species is more common during the rainy season in tropical and subtropical regions of the states. Adults are diurnal and females can feed on a wide variety of hosts including humans. The recent record of *Ae. albopictus* in Mexico City is worrisome for public health officials because the Metropolitan Area of Mexico City is one of the world's most populated cities. In 2010, INEGI (Instituto Nacional de Estadística y Geografía) recorded 22 million people living in the Metropolitan Area of Mexico City. The presence of *Ae. aegypti* and *Ae. albopictus* make the Metropolitan Area of Mexico City an area at high risk of vector-borne disease outbreaks.

Haemagogus equinus Theobald (Diptera: Culicidae): In Mexico, 4 species of the genus *Haemagogus* has been reported, with *Hg. equinus* the most common. This mosquito was collected from ovitraps placed in the states of Chiapas, Hidalgo, and Morelos. This is the first record of *Hg. equinus* in Morelos State. This species occurs in the tropical forests in southern Mexico, and in the northeastern state of Tamaulipas, reaching their northernmost distribution in southeastern Texas, USA. In the wild, *Hg. equinus* develops mostly in tree holes but the species has been collected in artificial containers (discarded tires and buckets), always with murky water and organic material. The adults are active during the d, and females feed on humans along the shaded forest floor.

Culex (Anodiopora) restrictor Dyar & Knab (Diptera: Culicidae): *Culex restrictor* was collected in ovitraps placed in the state of Tabasco. Previously, this species has been reported from southeastern Mexico to the northeastern state of Tamaulipas (Ortega-Morales et al. 2015). *Cx. restrictor* commonly develops in tree holes and bamboo internodes but it also has been collected from artificial containers (such as discarded tires). Nothing is known about the biology of the adult stages or medical importance. In southeastern Mexico, adults have been collected using CDC light traps baited with octenol (A.I.O-M. unpublished data).

Culex coronator sensu lato Dyar & Knab (Diptera: Culicidae): In Mexico, the *Culex coronator* complex includes the species *Cx. coronator* Dyar & Knab, *Cx. ousqua* Dyar, and *Cx. usquatus* Dyar. We were unable to accurately distinguish which species from the complex was collected because identification relies on morphometric analysis of male genitalia and we only reared progeny to the larval stage. *Culex coronator s.l.* was collected from ovitraps in the state of Chiapas, where *Cx. coronator s.s.* and *Cx. ousqua* have been previously recorded (Casas-Martínez et al. 2012). Immature stages of *Cx. coronator s.l.* inhabit a variety of aquatic sites. In Mexico, members of this group have been collected in natural aquatic sites such as ponds and swamps, always with abundant emerging vegetation, and in the partially shaded margins of streams. They also have been collected in artificial containers, such as buckets and cattle drinking canals filled with clean water. Adults are most active at dusk and nighttime; females feed on birds and a variety of mammals (Molaei et al. 2006). Although many viruses have been isolated from *Cx. coronator s.s.*, the species is not considered a vector of human diseases.

Culex mollis Dyar & Knab (Diptera: Culicidae): *Culex mollis* was collected from ovitraps placed in the state of Tabasco. This species also is considered an uncommon mosquito. In historical records for Mexico, *Cx. mollis* has been recorded in only a few instances, and then from the tropical regions of southeastern Mexico (Díaz-Nájera & Vargas 1973). The immature stages of *Cx. mollis* are found commonly in tree holes with murky water and abundant organic material. The biology of the adult stage, feeding preference, and medical importance is unknown.

Culex quinquefasciatus Say (Diptera: Culicidae): This species was collected from ovitraps placed in the state of Tabasco. This is probably one of the most common mosquito species in Mexico. *Culex quinquefasciatus* has been recorded in all states of the country and its populations are stable throughout the year. This species uses a wide variety of natural and artificial habitats with abundant organic material. Larvae can be found in open ponds, ditches, and less frequently in axils of plants, discarded tires, flower vases, buckets, cattle drinking canals, and abandoned swimming pools. Adults are nocturnal and active from dusk to sunrise. Females are opportunistic and feed on a variety of hosts including birds, mammals, and humans. *Culex quinquefasciatus* has been incriminated as an important vector of St. Louis Encephalitis and West Nile Virus.

Limatus durhamii Theobald (Diptera: Culicidae): This species was collected in ovitraps placed in forests of the states of Campeche (Casas-Martínez et al. 2012), Chiapas, and Tabasco (A.I. O-M. personal observation). Generally, the immature stages of this mosquito have been found in sites such as bamboo internodes, fruit casings, axils of plants, and crab shells. However, in Mexico this species is most frequently found in artificial containers (buckets, discarded tires, and flower vases), always with clean water and low organic material. Adults are diurnal and females are persistent biters that commonly feed on humans. Although a number of viruses, including the yellow fever virus, have been detected in *Li. durhamii*, it is not considered an important vector of human disease.

Wyeomyia guatemala Dyar & Knab (Diptera: Culicidae): This species was collected from ovitraps placed in the state of Chiapas. Typically, *Wy. guatemala* lays eggs in bromeliad axils. Historical records of *Wyeomyia* have reported the presence of larvae in bromeliad axils in tropical regions in southeastern Mexico (Ortega-Morales et al. 2010). The adults are diurnal and live in the shade of tropical rain forests. Females fly near ground level and feed on humans. The medical importance of *Wy. guatemala* is unknown.

Acknowledgments

We thank the control personnel from the Ministry of Health of the 32 states of Mexico for their assistance with field work. We thank all the staff from each Mexican Specialized Bioassay Unit. We are extremely grateful to Audrey Lenhart for her comments and proofreading this paper for the use of English.

References Cited

- Ávila-Rodríguez V, Pérez-Muñoz R, Márquez-Hernández C, Castañeda-Gaytán G, Nava-Cambreros U. 2012. Diversidad de mosquitos Culícidos en cinco municipios de Durango, México. *Entomologia Mexicana* pp. 878–882. www.entomologia.socmexent.org/revista/entomologia/2013/EMF/878-882.pdf (last accessed 18 Jul 2018).
- Casas Martínez M, Orozco Bonilla A, Bond-Compeán JG. 2012. Diversidad y distribución geográfica de las especies de culícidos de importancia médica en la región centro-occidental de México. Instituto Nacional de Salud Pública. Centro Regional de Investigación en Salud Pública. Informe final SNIB-CONABIO, proyecto No. FE009. Mexico City, Distrito Federal, Mexico.
- CDC (Centers for Disease Control and Prevention). 2017. Surveillance and Control of *Aedes aegypti* and *Aedes albopictus* in the United States. Centers for Disease Control and Prevention, Atlanta, Georgia, USA. <https://www.cdc.gov/chikungunya/pdfs/Surveillance-and-Control-of-Aedes-aegypti-and-Aedes-albopictus-US.pdf> (last accessed 27 Apr 2018).
- CENAPRECE (Centro Nacional de Programas Preventivos y Control de Enfermedades). 2017. *Guía metodológica para la vigilancia entomológica con ovitrampas*. Centro Nacional de Programas Preventivos y Control de Enfermedades Mexico City, Mexico City, Distrito Federal, Mexico. https://www.gob.mx/cms/uploads/attachment/file/37865/guia_vigilancia_entomologica_ovitrampas.pdf (last accessed 23 Mar 2018).
- Ceretti-Junior W, Medeiros-Sousa AR, Multini LC, Urbinatti PR, Vendrami DP, Natal D, Marques S, Fernandes A, Ogata H, Marrelli MT. 2014. Immature mosquitoes in bamboo internodes in municipal parks, city of Sao Paulo, Brazil. *Journal of the American Mosquito Control Association* 30: 268–274.
- Chaverri LG, Dillenbeck C, Lewis D, Rivera C, Romero LM, Chaves LF. 2018. Mosquito species (Diptera: Culicidae) diversity from ovitraps in a Mesoamerican tropical rainforest. *Journal of Medical Entomology* 55: 646–653.
- Cheong WH. 1967. Preferred *Aedes aegypti* larval habitats in urban areas. *Bulletin of the World Health Organization* 36: 586–589.
- Clark-Gil S, Darsie R. 1983. The mosquitoes of Guatemala, distribution and bionomics. *Mosquito Systematics* 15: 151–284.
- Codeço CT, Lima AWS, Araújo SC, Lima JBP, Maciel-de-Freitas R, Honório NA, Galardo AKR, Braga IA, Coelho GE, Valle D. 2015. Surveillance of *Aedes aegypti*: comparison of house index with four alternative traps. *PLoS Neglected Tropical Diseases* 9: e0003475. doi: 10.1371/journal.pntd.0003475
- DGE (Dirección General de Epidemiología). 2017. Boletín Epidemiológico Sistema Nacional de Vigilancia Epidemiológica Sistema Único de Información. Mexico City. Dirección General de Epidemiología. <https://www.gob.mx/salud/acciones-y-programas/direccion-general-de-epidemiologia-boletin-epidemiologico> (last accessed 23 Mar 2018).
- Díaz-Nájera A, Vargas L. 1973. Mosquitos mexicanos, distribución geográfica actualizada. *Revista de Investigación en Salud Pública* 22: 111–125.
- Docile TN, Figueiro R, Honorio NA, Baptista DF, Pereira G, Santos JAA dos, Codeço CT. 2017. Frequency of *Aedes* sp. Linnaeus (Diptera: Culicidae)

- and associated entomofauna in bromeliads from a forest patch within a densely urbanized area. *Neotropical Entomology* 46: 613–621.
- Espinoza-Gómez F, Arredondo-Jiménez JI, Maldonado-Rodríguez A, Pérez-Rentería C, Newton-Sánchez ÓA, Chávez-Flores E, Gómez-Ibarra E. 2013. Distribución geográfica de mosquitos adultos (Diptera: Culicidae) en áreas selváticas de Colima, México. *Revista Mexicana de Biodiversidad* 84: 685–689.
- Estallo EL, Ludueña-Almeida FF, Visintin AM, Scavuzzo CM, Intorini MV, Zaidenberg M, Almirón WR. 2011. Prevention of dengue outbreaks through *Aedes aegypti* oviposition activity forecasting method. *Vector Borne Zoonotic Diseases* 11: 543–549.
- Flisser A, Velasco-Villa A, Martínez-Campos C, González-Domínguez F, Briseño-García B, García-Suárez R, Caballero-Servín A, Hernández-Monroy I, García-Lozano H, Gutiérrez-Cogco L, Rodríguez-Angeles G, López-Martínez I, Galindo-Virgen S, Vázquez-Campuzano R, Balandrano-Campos S, Guzmán-Bracho C, Olivo-Díaz A, Rosa JL de la, Magos C, Escobar-Gutiérrez A, Correa D. 2002. Infectious diseases in Mexico. A survey from 1995–2000. *Archives of Medical Research* 33: 343–350.
- Francy DB, Moore CG, Eliason DA. 1990. Past, present and future of *Aedes albopictus* in the United States. *Journal of the American Mosquito Control Association* 6: 127–132.
- Heinemann S, Belkin J. 1977. Collection records of the project “Mosquitoes of Middle America” 9. Mexico (MEX, MF, MT, MX). *Mosquito Systematics* 9: 483–535.
- Hernandez-Avila JE, Rodriguez M-H, Santos-Luna R, Sanchez-Castaneda V, Roman-Perez S, Rios-Salgado VH, Salas-Sarmiento JA. 2013. Nationwide, web-based, geographic information system for the integrated surveillance and control of dengue fever in Mexico. *PLoS One* 8: e70231. doi.org/10.1371/journal.pone.0070231
- Ibanez-Bernal S, Martinez-Campos C. 1994a. Clave para la identificación de larvas de mosquitos comunes en las áreas urbanas y suburbanas de la República Mexicana (Diptera: Culicidae). *Folia Entomológica Mexicana* 92: 43–73.
- Ibáñez-Bernal S, Martínez-Campos C. 1994b. *Aedes albopictus* in Mexico. *Journal of the American Mosquito Control Association* 10: 231–232.
- Kuri-Morales P, Correa-Morales F, González-Acosta C, Sánchez-Tejeda G, Dávalos-Becerril E, Fernanda Juárez-Franco M, Díaz-Quiñonez A, Huerta-Jiménez H, Mejía-Guevara MD, Moreno-García M, González-Roldán JF. 2017. First report of *Stegomyia aegypti* (= *Aedes aegypti*) in Mexico City, Mexico. *Medical and Veterinary Entomology* 31: 240–242.
- Martínez MC, Estrada JLT. 2003. First evidence of *Aedes albopictus* (Skuse) in southern Chiapas, Mexico. *Emerging Infectious Diseases* 9: 606–607.
- Molaei G, Andreadis TG, Armstrong PM, Anderson JF, Vossbrinck CR. 2006. Host feeding patterns of *Culex* mosquitoes and West Nile Virus transmission, northeastern United States. *Emerging Infectious Diseases* 12: 468–474.
- Mora-Covarrubias A de la, Jiménez-Vega F, Treviño-Aguilar SM. 2010. Distribución geoespacial y detección del virus del dengue en mosquitos *Aedes (Stegomyia) aegypti* de Ciudad Juárez, Chihuahua, México. *Salud Pública de México* 2: 127–133.
- Ortega-Morales AI, Bond G, Méndez-López R, Garza-Hernández JA, Hernández-Triana LM, Casas-Martínez M. 2018. First record of invasive mosquito *Aedes albopictus* in Tabasco and Yucatan, Mexico. *Journal of the American Mosquito Control Association* 34: 120–123.
- Ortega-Morales AI, Cueto-Medina SM, Siller-Rodríguez QK. 2016. First record of the Asian tiger mosquito *Aedes albopictus* in Hidalgo State, Mexico. *Journal of the American Mosquito Control Association* 32: 234–236.
- Ortega Morales AI, Mis-Avila P, Elizondo-Quiroga A, Harbach RE, Siller-Rodríguez QK, Fernández-Salas I. 2010. The mosquitoes of Quintana Roo State, Mexico (Diptera: Culicidae). *Acta Zoológica Mexicana* 26: 33–46.
- Ortega-Morales AI, Siller-Rodríguez QK. 2016. First record of *Aedes albopictus* (Diptera: Culicidae) in San Luis Potosi, Mexico. *Journal of Vector Ecology* 41: 314–315.
- Ortega-Morales AI, Zavortink TJ, Huerta-Jiménez H, Sánchez-Ramos FJ, Valdés Perezgagsa MT, Reyes-Villanueva G, Siller-Rodríguez QK, Fernández-Salas I. 2015. Mosquito records from Mexico: the mosquitoes (Diptera: Culicidae) of Tamaulipas State. *Journal of Medical Entomology* 52: 171–184.
- Pesina HO, Hernandez RM, Valdez Rodriguez MA. 2001. *Aedes albopictus* in Allende City, Nuevo León, Mexico. *Journal of the American Mosquito Control Association* 17: 260–261.
- Salomón-Grajales J, Lugo-Moguel GV, Tinal-Gordillo VR, Cruz-Velázquez J de la, Beaty BJ, Eisen L, Lozano-Fuentes S, Moore CG, García-Rejón JE. 2012. *Aedes albopictus* mosquitoes, Yucatan Peninsula, Mexico. *Emerging Infectious Disease* 18: 525–527.
- Schick RX. 1970. Mosquito Studies (Diptera, Culicidae). XX - The Terrens Group of *Aedes* (Finlaya). *Contributions of the American Entomological Institute* 5: 1–158.
- Snetselaar J, Andriessen R, Suer RA, Osinga AJ, Knols BG, Farenhorst M. 2014. Development and evaluation of a novel contamination device that targets multiple life-stages of *Aedes aegypti*. *Parasite & Vectors* 7: 200. doi.org/10.1186/1756-3305-7-200
- Torres-Avendaño JI, Castillo-Ureta H, Torres-Montoya EH, Meza-Carrillo E, Lopez-Mendoza RL, Vazquez-Martinez MG, Rendon-Maldonado JG. 2015. First record of *Aedes albopictus* in Sinaloa, Mexico. *Journal of the American Mosquito Control Association* 31: 164–166.
- Villegas-Trejo A, Manrique-Saide P, Che-Mendoza A, Cruz-Canto W, Fernández MG, González-Acosta C, Dzul-Manzanilla F, Huerta H, Arredondo-Jiménez JI. 2010. First report of *Aedes albopictus* and other mosquito species in Morelos, Mexico. *Journal of the American Mosquito Control Association* 26: 321–323.
- Wu H-H, Wang C-Y, Teng H-J, Lin C, Lu L-C, Jian S-W, Chang N-T, Wen T-H, Wu J-W, Liu D-P, Lin L-J, Norris DE, Wu H-S. 2013. A dengue vector surveillance by human population-stratified ovitrap survey for *Aedes* (Diptera: Culicidae) adult and egg collections in high dengue-risk areas of Taiwan. *Journal of Medical Entomology* 50: 261–269.
- Yee DA, Kneitel JM, Juliano SA. 2010. Environmental correlates of abundances of mosquito species and stages in discarded vehicle tires. *Journal of Medical Entomology* 47: 53–62.
- Zavortink TJ. 1972. The New World species formerly placed in *Aedes* (Finlaya). *Contributions of the American Entomological Institute* 8: 1–206.