

A Newly Detected Exotic Ambrosia Beetle in Argentina: *Euwallacea interjectus* (Coleoptera: Curculionidae: Scolytinae)

Authors: Landi, Lucas, Braccini, Celina Laura, Knížek, Milos, Pereyra, Vanina Antonella, and Marvaldi, Adriana Elena

Source: Florida Entomologist, 102(1) : 240-242

Published By: Florida Entomological Society

URL: <https://doi.org/10.1653/024.102.0141>

BioOne Complete (complete.BioOne.org) is a full-text database of 200 subscribed and open-access titles in the biological, ecological, and environmental sciences published by nonprofit societies, associations, museums, institutions, and presses.

Your use of this PDF, the BioOne Complete website, and all posted and associated content indicates your acceptance of BioOne's Terms of Use, available at www.bioone.org/terms-of-use.

Usage of BioOne Complete content is strictly limited to personal, educational, and non - commercial use. Commercial inquiries or rights and permissions requests should be directed to the individual publisher as copyright holder.

BioOne sees sustainable scholarly publishing as an inherently collaborative enterprise connecting authors, nonprofit publishers, academic institutions, research libraries, and research funders in the common goal of maximizing access to critical research.

A newly detected exotic ambrosia beetle in Argentina: *Euwallacea interjectus* (Coleoptera: Curculionidae: Scolytinae)

Lucas Landi^{1,*}, Celina Laura Braccini¹, Milos Knížek², Vanina Antonella Pereyra³, and Adriana Elena Marvaldi⁴

In Argentina, poplar (*Populus* spp.: Salicaceae) cultivation has been developing since the late 19th century, particularly in the lower Paraná Delta region. The poplar-forested area in the country was estimated to be around 14,500 has by 2012 (Signorelli & Gaute 2012). Poplars provide lumber for various uses and are important in the manufacture of reconstituted board products. Also, they represent the basis for the pulp and paper industry. Although some poplar species may have fairly extensive geographic ranges, they are restricted in occurrence and often exhibit their best development in riparian areas. Following a significant drought event in Dec 2008, some scolytine species (Coleoptera: Curculionidae: Scolytinae) have appeared regularly in poplar forests in the region (Landi et al. 2017). Today, the rate of introduction and establishment of non-native insect species is increasing worldwide as a result of human activity, mainly due to the transport of goods in international trade (Brockerhoff et al. 2006). In the presence of suitable conditions, the establishment of exotic species may be favored, and may become invasive by attacking living trees (Liebhold et al. 1995).

Among the ambrosia-feeding scolytines, the Xyleborini includes approximately 1,200 species in 30 genera that are distributed in forests throughout the world (Cognato et al. 2011). Besides their xylomycectophagy, the Xyleborini are biologically defined by extreme inbreeding and the presence of a haplodiploid genetic system wherein haploid males are flightless and often dwarfed (Hulcr et al. 2007).

Euwallacea (Hopkins) (Coleoptera: Curculionidae) is a large, cosmopolitan genus that contains over 50 species, mostly native from Asia (Wood & Bright 1992). Only a few species are introduced in the Americas (e.g., *Euwallacea fornicatus* Eichhoff, *Euwallacea validus* Eichhoff, and *Euwallacea interjectus* Blandford) (Rabaglia et al. 2006; Cognato et al. 2015), none of which is reported to occur presently in Argentina.

The purpose of this article is to communicate the detection of the Asian ambrosia beetle *E. interjectus* in Argentina and South America for the first time. Both traditional morphology-based taxonomy and molecular diagnosis through DNA barcoding were combined in order to identify the species.

In Argentina, *E. interjectus* was first found in Apr 2009 from a 10-yr-old poplar plantation (34.139332°S, 58.795332°W). Healthy trees of *Populus deltoides* W. Bartram cv 'Deltoides' (Salicaceae) attracted

attention due to the presence of sap exuding from beetle entrance holes. They were cut into 30 cm logs and taken to the laboratory, where they were maintained in order to capture the adult beetles as they emerged; the beetles were preserved in 96% ethanol. Afterwards, the logs were chopped, immature stages were collected, and wood damage (i.e., galleries, wood stained by symbiotic fungi) was confirmed. Later, between Nov 2012 and Feb 2013, 12-unit multifunnel traps were located in poplar plantations as part of a surveillance program. Traps were baited with ethanol lures (low rate of release), and wet collection cups filled with propylene glycol (150 mL) were used as the killing and preserving agent. Consequently, a second collection was achieved. In both cases, adults were deposited at the Instituto Nacional de Tecnología Agropecuaria (Instituto de Recursos Biológicos, Buenos Aires, Argentina), and samples were selected for both morphological and molecular study.

Taxonomic identification was based on structural characters observed under stereomicroscope, and photographs of diagnostic characters were taken by means of scanning electron microscopy. As Cognato et al. (2015) pointed out, *E. interjectus* shares very similar morphological traits with *E. validus*. In general, these species can be distinguished by the shape of the declivity, the punctures, and placement of the tubercles on the second declivital interstriae. In *E. interjectus*, the declivity is gradually sloped from base to apex, punctures of the declivital striae are shallow, giving the declivity a smooth appearance, and tubercles on the second declivital interstriae extend from the base to the apex (Fig. 1).

Sequences of cytochrome oxidase subunit I (COI) were acquired from 2 of the specimens sampled from poplar plantations in Argentina in order to verify the species identification with molecular evidence. The primers used allowed us to amplify and sequence almost an entire COI fragment that includes both the 5' region, the standardized DNA "barcode," as well as the 3' region of COI, which also is widely used for species identification purposes. The protocols and conditions for DNA isolation, PCR amplification, and sequencing of the COI gene were conducted as in Pereyra et al. (2016). Sequences then were checked and compared with others available in GenBank through the Basic Local Alignment Search Tool

¹Instituto Nacional de Tecnología Agropecuaria (INTA), Instituto de Recursos Biológicos, De los Reseros y Dr. Nicolás Repetto s/n, CP 1686, Hurlingham, Buenos Aires, Argentina; E-mail: landi.lucas@inta.gob.ar (L. L.); braccini.celina@inta.gob.ar (C. L. B.)

²Forestry and Game Management Research Institute, Jíloviště - Strnady, Praha 5 - Zbraslav, CZ 156 04, Czechia; E-mail: knizek@vulhm.cz (M. K.)

³Laboratorio de Entomología, Instituto Argentino de Investigaciones de las Zonas Áridas (IADIZA - CCT - CONICET Mendoza), Av. Ruiz Leal s/n, Parque Gral. San Martín - Mendoza CC 507 (5500), Mendoza, Argentina; E-mail: vanipereyra@gmail.com (V. A. P.)

⁴División Entomología, Facultad de Ciencias Naturales y Museo, Universidad Nacional de La Plata, Consejo Nacional de Investigaciones Científicas y Técnicas (CONICET), Paseo del Bosque s/n, 1900 La Plata, Buenos Aires, Argentina; E-mail: marvaldi@fcnym.unlp.edu.ar (A. E. M.)

*Corresponding author; E-mail: landi.lucas@inta.gob.ar, marvaldi@fcnym.unlp.edu.ar




Fig. 1. *Euwallacea interjectus* female. (A) Lateral view, (B) Dorsal view, (C) Elytral declivity gradually sloped to the apex, showing tubercles on the second declivital interstriae. Scale bars: (A, B) 500 μ m, (C) 200 μ m.

(BLAST). BLAST searches gave significant alignments with *E. interjectus*, with 100% - 99% identity values with specimens from Japan and USA sequenced in previous studies (e.g., Cognato et al. 2015). Voucher specimens are deposited in the Entomology collection of the Museo de Ciencias Naturales de La Plata (La Plata, Buenos Aires, Argentina). The new sequences of *E. interjectus* from Argentina are deposited in GenBank under accession numbers MH105073 and MH105074.

The occurrence of genus *Euwallacea* in South America has been detected at least once during the current decade. In 2011, SAG (Servicio Agrícola Ganadero de Chile) intercepted *E. validus* in a ship with containers filled with stamped wooden packing coming from the US. The

presence of *E. interjectus* in poplar cultivations of Argentina represents a new find and is cause for concern due to the potential of this species to become invasive and spread rapidly (Cognato et al. 2015). The geographical expansion of the species is probably in progress, favored by both the shipment of timber products globally and the characteristics of the mating system of many Scolytinae (Kirkendall & Faccoli 2010; Rassati et al. 2018). Potential risks associated with this species include not only direct wood damage but also the spread of different species of fungi (O'Donnell et al. 2015). *Euwallacea interjectus* is among the relatively few species that attack living trees. Although it has been detected only affecting stressed plants so far, under suitable conditions it may attack healthy trees (Knížek & Beaver 2004). Consequently, *E. interjectus* could become economically relevant if it damages healthy stands of poplar monocultures in the Paraná Delta region. The genus *Euwallacea* has been reported to attack a significant number of hosts even in riparian areas (Boland 2016). Previously, *E. interjectus* had not been observed affecting poplar trees outside its natural range of distribution (Wood & Bright 1992). This article constitutes the first report of the species not only in South America and Argentina, but also affecting cultivated poplars. *Euwallacea interjectus* is now abundant in the Delta of Paraná River region, and several incidences of mass attack on live water-stressed poplars have been observed. Thus, there is an urgent need for study and monitoring of this exotic species as it may become a serious problem.

We acknowledge Dr. Osvaldo Di Iorio for first calling attention to this species as a new detection in Argentina. We thank Dra. Patricia Fernández and Dr. Alain Roques for encouragement to contact different specialists. The help of Dra. Andrea Vega during the first scanning electron microscopy session is gratefully appreciated. This work was supported by the National Agency of Promotion of Science (ANPCyT, Argentina, PICTs 2011-2573 and 2016-2798 to A.E.M.), and by the National Institute of Agricultural Technology (INTA, Argentina, PNFOR 2212 1104072 and PRET BANOR 1271411). A portion of the research of M. Knížek was supported by the Ministry of Agriculture of the Czech Republic, institutional support MZE-RO0118.

Summary

An exotic species of ambrosia beetle (Coleoptera: Curculionidae: Scolytinae) of Asian origin has been detected recently in Argentina. *Euwallacea interjectus* (Blandford) has been collected infesting cultivated poplars in the Paraná Delta region. Identification was based on morphology and on DNA sequences of the COI gene. This finding adds to the increasing number of recent introductions of exotic ambrosia beetles into South America.

Key Words: Scolytinae; Xyleborini; DNA; new record; geographical distribution; poplars

Sumario

Recientemente se detectó por primera vez en Argentina una especie exótica de escarabajo de Ambrosía (Coleoptera: Curculionidae: Scolytinae) de origen asiático. *Euwallacea interjectus* (Blandford) fue colectada afectando cultivos de álamo en la región del Delta del Río Paraná. Su identificación se basó en la morfología y en las secuencias de ADN del gen COI. Este hallazgo se suma al número cada vez mayor de introducciones recientes de escarabajos de Ambrosía exóticos en Sudamérica.

Palabras Clave: Scolytinae; Xyleborini; ADN; nuevo registro; distribución geográfica; álamos

References Cited

- Boland JM. 2016. The impact of an invasive ambrosia beetle on the riparian habitats of the Tijuana River Valley, California. *PeerJ* 4: e2141. doi: 10.7717/peerj.2141
- Brockhoff EG, Bain J, Kimberley JM, Knížek M. 2006. Interception frequency of bark and ambrosia beetles (Coleoptera: Scolytinae) and relationship with establishment in New Zealand and worldwide. *Canadian Journal of Forest Research* 36: 289–298.
- Cognato AI, Hulcr J, Dole SA, Jordal BH. 2011. Phylogeny of haplo-diploid, fungus-growing ambrosia beetles (Curculionidae: Scolytinae: Xyleborini) inferred from molecular and morphological data. *Zoologica Scripta* 40: 174–186.
- Cognato AI, Hoebeke ER, Kajimura H, Smith SM. 2015. History of the exotic ambrosia beetles *Euwallacea interjectus* and *Euwallacea validus* (Coleoptera: Curculionidae: Xyleborini) in the United States. *Journal of Economic Entomology* 108: 1129–1135.
- Hulcr J, Dole SA, Beaver RA, Cognato AI. 2007. Cladistic review of generic taxonomic characters in Xyleborina (Coleoptera: Curculionidae: Scolytinae). *Systematic Entomology* 32: 568–584.
- Kirkendall L, Faccoli M. 2010. Bark beetles and pinehole borers (Curculionidae, Scolytinae, Platypodinae) alien to Europe. *ZooKeys* 56: 227–251.
- Knížek M, Beaver R. 2004. Taxonomy and systematics of bark and ambrosia beetles, pp. 41–54. In Lieutier F, Day KR, Battisti A, Grégoire JC, Evans HF [eds.], *Bark and Wood Boring Insects in Living Trees in Europe, a Synthesis*. Kluwer Academic Publishers, Dordrecht, The Netherlands and Boston, Massachusetts, USA.
- Landi L, Gómez D, Braccini CL, Pereyra VA, Smith SM, Marvaldi AE. 2017. Morphological and molecular identification of the invasive *Xylosandrus crassiusculus* (Coleoptera: Curculionidae: Scolytinae) and its South American range extending into Argentina and Uruguay. *Annals of the Entomological Society of America* 110: 1–6.
- Liebholt AM, MacDonald WL, Bergdahl D, Mastro VC. 1995. Invasion by exotic forest pests: a threat to forest ecosystems. *Forest Science* 30: 1–49.
- O'Donnell K, Sink S, Libeskind-Hadas R, Hulcr J, Kasson MT, Ploetz RC, Konkol JL, Ploetz JN, Carrillo D, Campbell A, Duncan RE, Liyanage PN, Eskalen A, Na F, Geiser DM, Bateman C, Freeman S, Mendel Z, Sharon M, Aoki T, Cossé AA, Rooney AP. 2015. Discordant phylogenies suggest repeated host shifts in the *Fusarium–Euwallacea* ambrosia beetle mutualism. *Fungal Genetics and Biology* 82: 277–290.
- Pereyra VA, Gomez C, La Manna L, Roux G, Lanteri AA, Vallejos N, Marvaldi AE. 2016. Introduction and establishment of *Pissodes castaneus* (Coleoptera: Curculionidae) in the Andean Patagonia of Argentina. *Journal of Economic Entomology* 109: 222–231.
- Rabaglia RJ, Dole SA, Cognato AI. 2006. Review of American Xyleborina (Coleoptera: Curculionidae: Scolytinae) occurring north of Mexico, with an illustrated key. *Annals of the Entomological Society of America* 99: 1034–1056.
- Rassati D, Haack RA, Knížek M, Faccoli M. 2018. National trade can drive range expansion of bark- and wood-boring beetles. *Journal of Economic Entomology* 111: 260–268.
- SAG (Servicio Agrícola Ganadero de Chile). 2018. <http://www.sag.gob.cl> (last accessed 16 Jan 2018).
- Signorelli A, Gaute M. 2012. Plantaciones forestales en las islas del Delta del Paraná. <http://deltaforestal.blogspot.com.ar/2012/01/plantaciones-forestales-en-las-islas.html> (last accessed 16 Jan 2018).
- Wood SL, Bright DE. 1992. A catalog of Scolytidae and Platypodidae (Coleoptera), part 2. Taxonomic index Volume A. *Great Basin Naturalist Memoirs* 13: 1–833.