

**Protopolybia Exigua (Hymenoptera: Vespidae) Nesting on Citrus Grandis (Rutaceae)**

Authors: Bruno Pandelo Brügger, Antonio José Vinha Zanuncio, Carlos Frederico Wilcken, Marcus Alvarenga Soares, and José Cola Zanuncio

Source: Florida Entomologist, 102(1) : 262-263

Published By: Florida Entomological Society

URL: <https://doi.org/10.1653/024.102.0149>

---

BioOne Complete ([complete.BioOne.org](http://complete.BioOne.org)) is a full-text database of 200 subscribed and open-access titles in the biological, ecological, and environmental sciences published by nonprofit societies, associations, museums, institutions, and presses.

Your use of this PDF, the BioOne Complete website, and all posted and associated content indicates your acceptance of BioOne's Terms of Use, available at [www.bioone.org/terms-of-use](http://www.bioone.org/terms-of-use).

Usage of BioOne Complete content is strictly limited to personal, educational, and non-commercial use. Commercial inquiries or rights and permissions requests should be directed to the individual publisher as copyright holder.

---

BioOne sees sustainable scholarly publishing as an inherently collaborative enterprise connecting authors, nonprofit publishers, academic institutions, research libraries, and research funders in the common goal of maximizing access to critical research.

# *Protopolybia exigua* (Hymenoptera: Vespidae) nesting on *Citrus grandis* (Rutaceae)

Bruno Pandelo Brügger<sup>1,\*</sup>, Antonio José Vinha Zanuncio<sup>2</sup>, Carlos Frederico Wilcken<sup>3</sup>, Marcus Alvarenga Soares<sup>4</sup>, and José Cola Zanuncio<sup>1</sup>

*Protopolybia exigua* (Saussure) (Hymenoptera: Vespidae) has been reported in Brazil from the states of Acre, Amazonas, Bahia, Ceará, Espírito Santo, Goiás, Maranhão, Minas Gerais, Mato Grosso, Pará, Pernambuco, Rio de Janeiro, Rio Grande do Sul, Santa Catarina, and São Paulo (Richards 1978). Social wasps are predators (Donovan 2003; De Souza et al. 2012) that feed on insects, mainly defoliating Lepidoptera (Richter 2000). They contribute to insect population reduction, and represent an important natural enemy group for biological control (Prezoto & Braga 2013). In addition, these insects are used in environmental impact as-

essment and evaluation of forest conservation (Dos Santos et al. 2016; Brügger et al. 2011, 2017). Although known from native (forest) environments, this species is not well known in cultivated agroecosystems. The objective of this communication is to report the first case of nesting of *P. exigua* on citrus plants.

Two nests were collected with an entomological net in Divinópolis, Minas Gerais State, Brazil (20.149904°S, 44.895827°W) on pomelo (*Citrus grandis* [L.] (Rutaceae) in Jul of 2017. One of the *P. exigua* nests found under pomelo leaves (Fig. 1) was active and another was aban-



Fig. 1. *Protopolybia exigua* (Hymenoptera: Vespidae) nest under *Citrus grandis* (Rutaceae) leaf.

<sup>1</sup>Departamento de Entomologia/BIOAGRO, Universidade Federal de Viçosa, 36570-900, Viçosa, Minas Gerais, Brazil; E-mail: brunopb2002@yahoo.com.br (B. P. B.); zanuncio@ufv.br (J. C. Z.)

<sup>2</sup>Departamento de Engenharia Florestal, Universidade Federal de Viçosa, 36570-900 Viçosa, Minas Gerais, Brazil; E-mail: ajvzanuncio@ufu.br (A. J. V. Z.)

<sup>3</sup>Departamento de Proteção Vegetal, Universidade Estadual Paulista (UNESP, FCA), Botucatu, Botucatu, São Paulo, Brazil; E-mail: cwilcken@fca.unesp.br (C. F. W.)

<sup>4</sup>Programa de Pós-Graduação em Produção Vegetal, Universidade Federal dos Vales Jequitinhonha e Mucuri (UFVJM), 39100-000, Diamantina, Minas Gerais, Brazil; E-mail: marcussoares@yahoo.com.br (M. A. S.)

\*Corresponding author; E-mail: brunopb2002@yahoo.com.br

done. These insects were killed in ether vapor and preserved in 70% ethanol for identification. The numbers of pedicels, brood, and adults in the colony were determined.

The active colony of *P. exigua* had 12 pedicels, 600 brood cells, and 15 adult wasps. The number of pedicels, brood, and adults of this wasp in this colony were higher than those of this wasp collected in Pedregulho (southeastern Brazil), which was a younger nest, possessing 1 pedicel and 307 brood cells with 46 eggs, 28 workers, 37 intermediaries, and 30 queens (Noll et al. 1996). The high cell count but low numbers of active wasps in the *P. exigua* nest in this study indicate a declining stage. The life expectancy of queens of this wasp is up to 1 yr, but their nests are abandoned after about 6 mo due to the invasion by parasitoids of *P. exigua* wasp larvae, as reported for *Pachysomoides* sp. (Hymenoptera: Ichneumonidae: Cryptinae), and *Megaselia scalaris* (Loew) (Diptera: Phoridae) in nests of *Mischocyttarus cassununga* (von Ihering) (Hymenoptera: Vespidae) (Soares et al. 2006). In addition, the study was conducted in a cold, dry season, when foraging activity of *P. exigua* is lower, and few nests of this species are active (Ribeiro-Júnior et al. 2006).

The pedicel number of the *P. exigua* colony is common for the genus *Protopolybia*, whose species build nests supported by a central pedicel or several smaller ones (Wenzel 1998). The nests of these wasps are found underneath or between leaves, in the first case with a fragile and whitish wrapping, and in the second, supported by 1 leaf and several others glued to each other by oral secretion, functioning as an envelope (Somavilla et al. 2012).

The identification of nesting habits of social wasps in agroecosystems can be used to justify transferring or maintaining their colonies for biological control; moreover, there were no previous reports of *P. exigua* nesting on *C. grandis* plants.

We express our thanks to the Brazilian agencies “Conselho Nacional de Desenvolvimento Científico e Tecnológico (CNPq), Coordenação de Aperfeiçoamento de Pessoal de Nível Superior (CAPES/PELD), Fundação de Amparo à Pesquisa do Estado de Minas Gerais (FAPEMIG),” and “Programa Cooperativo sobre Proteção Florestal/ PROTEF” of the “Instituto de Pesquisas e Estudos Florestais/IPEF” for scholarships and financial support. We also are grateful to Andressa Vinha Zanuncio of the “Universidade Federal de São João Del Rey” for providing the nests studied.

## Summary

Social wasps are predators. Two nests of *Protopolybia exigua* (Saussure) (Hymenoptera: Vespidae) were collected on pomelo (*Citrus grandis* [L.] (Rutaceae) leaves, 1 active and another abandoned. The colonies of *P. exigua* were located under pomelo leaves, providing protection against adverse environmental conditions. The active nest had 12 pedicels, 600 brood cells, and 15 adult wasps. The knowledge of nesting habits of wasps in agroecosystems favors the management of these insects for biological control.

Key Words: biological control; foraging; grapefruit

## Sumario

Vespas sociais são predadores. Dois ninhos de *Protopolybia exigua* (Saussure, 1854) (Hymenoptera: Vespidae) foram coletados em folhas de grapefruit (*Citrus grandis* [L.] (Rutaceae), um ativo e outro abandonado. As colônias de *P. exigua* estavam localizadas sob folhas de pomelo, proporcionando proteção contra condições ambientais adversas. O ninho ativo tinha 12 pedicelos, 600 células de cria e 15 vespas adultas. O conhecimento dos hábitos de nidificação de vespas em agroecossistemas favorece o manejo desses insetos para o controle biológico.

Palavras Chave: controle biológico; forrageamento; pomelo

## References Cited

- Brügger BP, Araújo LSS, De Souza AR, Prezoto F. 2011. Social wasps (*Synoeca cyanea*) damaging *Psidium* sp. (Myrtaceae) fruits in Minas Gerais State, Brazil. *Sociobiology* 57: 533–535.
- Brügger BP, Castro BMC, Prezoto F, Serrão JE, Zanuncio JC. 2017. Feeding by the social wasp *Polybia scutellaris* (Hymenoptera: Vespidae) on *Syzygium jambos* (Myrtaceae) fruits in Minas Gerais, Brazil. *Florida Entomologist* 100: 172–173.
- De Souza AR, Venâncio DFA, Prezoto F, Zanuncio JC. 2012. Social wasps (Hymenoptera: Vespidae) nesting in eucalyptus plantations in Minas Gerais, Brazil. *Florida Entomologist* 95: 1000–1002.
- Donovan BJ. 2003. Potential manageable exploitation of social wasps, *Vespula* spp. (Hymenoptera: Vespidae), as generalist predators of insect pests. *International Journal of Pest Management* 49: 281–285.
- Dos Santos EF, Noll FB, Brandão CRF. 2016. Structural organization of the social paper wasp (Hymenoptera: Polistinae) assemblage along a latitudinal gradient in the Atlantic Rainforest: correlating fauna partitioning to biodiversity centers. *Journal of Insect Conservation* 20: 597–609.
- Noll FB, Mateus S, Zucchi R. 1996. Morphological caste differences in Neotropical swarm-founding Polistinae wasps. V - *Protopolybia exigua* (Hymenoptera: Vespidae). *Journal of the New York Entomological Society* 104: 62–69.
- Prezoto F, Braga N. 2013. Predation of *Zaprinus indianus* (Diptera: Drosophilidae) by the social wasp *Synoeca cyanea* (Hymenoptera: Vespidae). *Florida Entomologist* 96: 670–672.
- Ribeiro-Júnior C, Guimaraes DL, Elisei T, Prezoto F. 2006. Foraging activity rhythm of the Neotropical swarm-founding wasp *Protopolybia exigua* (Hymenoptera: Vespidae, Epiponini) in different seasons of the year. *Sociobiology* 47: 115–123.
- Richards OW. 1978. *The Social Wasps of the Americas, Excluding the Vespinae*. British Museum (Natural History), London. London, United Kingdom.
- Richter M. 2000. Social wasp (Hymenoptera: Vespidae) foraging behavior. *Annual Review of Entomology* 45: 121–150.
- Soares MA, Gutierrez CT, Zanuncio JC, Bellini LL, Prezotto F, Serrão JE. 2006. *Pachysomoides* sp. (Hymenoptera: Ichneumonidae: Cryptinae) and *Megaselia scalaris* (Diptera: Phoridae) parasitoids of *Mischocyttarus cassununga* (Hymenoptera: Vespidae) in Viçosa, Minas Gerais State, Brazil. *Sociobiology* 48: 673–680.
- Somavilla A, Oliveira MLD, Silveira OT. 2012. Identification guide for nests of social wasps (Hymenoptera: Vespidae: Polistinae) in reserva Ducke, Manaus, Amazonas, Brazil. *Revista Brasileira de Entomologia* 56: 405–414.
- Wenzel JW. 1998. A generic key to the nests of hornets, yellowjackets, and paper wasps worldwide (Vespidae: Vespinae, Polistinae). *American Museum Novitates* 3224: 1–39.