

**Response to Enantiomers of (Z3Z9)-6,7-Epoxy-Octadecadiene, Sex Pheromone Component of *Ectropis obliqua* Prout (Lepidoptera: Geometridae): Electroantennagram Test, Field Trapping, and in Silico Study**

Authors: Feng Guo, Jie Yu, Yunqiu Yang, and Xiaochun Wan

Source: Florida Entomologist, 102(3) : 549-554

Published By: Florida Entomological Society

URL: <https://doi.org/10.1653/024.102.0318>

---

BioOne Complete ([complete.BioOne.org](http://complete.BioOne.org)) is a full-text database of 200 subscribed and open-access titles in the biological, ecological, and environmental sciences published by nonprofit societies, associations, museums, institutions, and presses.

Your use of this PDF, the BioOne Complete website, and all posted and associated content indicates your acceptance of BioOne's Terms of Use, available at [www.bioone.org/terms-o-use](http://www.bioone.org/terms-o-use).

Usage of BioOne Complete content is strictly limited to personal, educational, and non - commercial use. Commercial inquiries or rights and permissions requests should be directed to the individual publisher as copyright holder.

---

BioOne sees sustainable scholarly publishing as an inherently collaborative enterprise connecting authors, nonprofit publishers, academic institutions, research libraries, and research funders in the common goal of maximizing access to critical research.

# Response to enantiomers of (Z3Z9)-6,7-epoxy-octadecadiene, sex pheromone component of *Ectropis obliqua* Prout (Lepidoptera: Geometridae): electroantennogram test, field trapping, and in silico study

Feng Guo<sup>1,2</sup>, Jie Yu<sup>3</sup>, Yunqiu Yang<sup>1</sup>, and Xiaochun Wan<sup>1,\*</sup>

## Abstract

Limited information on sex pheromone recognition by tea moths, *Ectropis obliqua* Prout (Lepidoptera: Geometridae), exists for this economically important pest of tea (*Camellia sinensis* L.; Theaceae). Pheromone binding proteins (PBPs), a sub-family of odorant-binding proteins, control transportation of pheromone molecules that may contribute to the discrimination of sex pheromone components. It has been reported previously that *EobIPBP1* gene (a pheromone binding protein of *E. obliqua*) is highly expressed in antennae of the male moth. Based on this information, a reliable model of *EobIPBP1* was constructed by homology modeling using the enantiomers of Z3Z9-6,7-epo-18:Hy docked into the hydrophobic cavity of the model. Docking results suggested similar binding affinities of this enantiomer to *EobIPBP1*. However, electroantennogram and field trapping experiments of *E. obliqua* males revealed that response to Z3Z9-(6S,7R)-epo-18:Hy was significantly greater than the opposite configuration, and suggested enantiomeric discrimination could occur on sex pheromone receptors of this species of tea moth.

Key Words: enantiomers; homology modeling; molecular docking; discrimination; *EobIPBP1*

## Resumen

Información limitada existe sobre el reconocimiento de la feromona sexual por la polilla del té, *Ectropis obliqua* Prout (Lepidoptera: Geometridae), una plaga del té de importancia económica (*Camellia sinensis* L.; Theaceae). Las proteínas de unión de feromonas (PBP), una subfamilia de proteínas de unión a odorantes, controlan el transporte de moléculas de feromonas que pueden contribuir a la discriminación de los componentes de las feromonas sexuales. Se ha informado previamente que el gen *EobIPBP1* (una proteína de unión a la feromona de *E. obliqua*) se expresa altamente en las antenas de la polilla macho. Basándose en esta información, se construyó un modelo confiable de *EobIPBP1* mediante un modelo de homología utilizando los enantiómeros de Z3Z9-6,7-epo-18:Hy acoplado en la cavidad hidrófoba del modelo. Los resultados de acoplamiento sugirieron afinidades de unión similares de este enantiómero a *EobIPBP1*. Sin embargo, el electroantennograma y los experimentos de captura de campo de machos de *E. obliqua* revelaron que la respuesta a Z3Z9-(6S, 7R)-epo-18:Hy fue significativamente mayor que la configuración opuesta y sugirió que podría existir discriminación enantiomérica en los receptores de feromonas sexuales de esta especie de polillas del té.

Palabras Clave: enantiómeros; modelo de homología; acoplamiento molecular; discriminación; *EobIPBP1*

The tea geometrid, *Ectropis obliqua* Prout (Lepidoptera: Geometridae) is a serious defoliator of tea bushes, *Camellia sinensis* L. (Theaceae), in southeastern China. Outbreak populations of this phytophagous pest can completely defoliate a tea bush, thereby causing severe reduction of tea production with considerable economic damage (Hazariika et al. 2009; Zhang et al. 2014). Heavy application of synthetic pesticides are still widely used to control this pest, which can result in undesirable residues on tea leaves and soil causing adverse health and environmental effects (Hazariika et al. 2001; Ye et al. 2014). The use of mating disruption strategies that employ area-wide application of sex pheromones to manage *E. obliqua* populations has become a promising alternative to pesticide applications (Hazariika et al. 2009). Attempts to identify sex pheromones of *E. obliqua* began 30 yr ago, but the components still have not been accurately identified (Li et al.

1988; Yao et al. 1991; Liu et al. 1994). Yang et al. (2015) examined the bioactive compounds extracted from female sex pheromone glands of *E. obliqua* and identified them as Z3Z9-6,7-epo-18:Hy (major component) and Z3Z6Z9-18:Hy (secondary component) (Fig. 1A). These researchers also reported on the behavioral responses of male moths of this species to these components in wind-tunnel and field bioassays.

Insect sex pheromones are hydrophobic molecules, so odorant binding proteins are needed to transport these molecules across the hydrophilic antennal lymph to olfactory receptor neurons located in male sensilla (Gómez et al. 2008). Odorant binding proteins are a group of small, soluble proteins in the sensillum lymph and are regarded as a first-layer filter to differentiate odorant molecules. Based on amino acid sequences and structural characteristics, lepidopteran odorant binding proteins can be divided into several groups, including pheromone

<sup>1</sup>State Key Laboratory of Tea Plant Biology and Utilization, Anhui Agricultural University, 130 Changjiang West Road, Hefei, Anhui 230036, P.R. China; E-mail: fengguo@ahau.edu.cn (F. G.), yyq\_lyh@126.com (Y. Y.), xcwan@ahau.edu.cn (X. C. W.)

<sup>2</sup>School of Life Science, Anhui Agricultural University, Hefei, Anhui 230036, P.R. China; E-mail: jjieyu@ahau.edu.cn (J. Y.)

<sup>3</sup>Department of Applied Chemistry, Anhui Agricultural University, Hefei, Anhui 230036, P.R. China

\* Corresponding author; Email: xcwan@ahau.edu.cn




Fig. 1. Sex pheromone components of *Ectropis obliqua* Prout. (A) sex pheromone components; (B) enantiomers of Z3Z9-6,7-epo-18:Hy.

binding proteins (PBPs) and general odorant binding proteins (Pelosi & Maida 1995). Pheromone binding proteins mainly bind to insect sex pheromones, and are thought to be involved in pheromone reception processes (Sun et al. 2013a; Jin et al. 2014) while general odorant binding proteins are believed to detect plant volatiles and sex pheromones (Liu et al. 2015a). Although several crystalline structures of pheromone binding proteins have been characterized, cloning, expression, and identification of their component parts have proven to be expensive and time consuming. Homology modeling, based on comparison of target and templates, provides a promising alternative to predict protein structures (Venthur et al. 2014). The modeled structures of pheromone binding proteins have been reported in several Lepidoptera insects such as *Bombyx mori* L. (Lepidoptera: Bombycidae) (Gräter et al. 2006), *Lymantria dispar* L. (Lepidoptera: Lymantriidae) (Honson et al. 2003), *Spodoptera litura* F. (Lepidoptera: Noctuidae) (Liu et al. 2012), and so on. Furthermore, the structures of pheromone binding proteins, prediction of binding site, clarification of binding mode, and interactions of protein-ligands can be performed by molecular docking and molecular dynamics (Mutis et al. 2014).

Recently, 2 enantiomers of the major sex pheromone components produced by female *E. obliqua*: Z3Z9-(6R,7S)-epo-18:Hy (1) and Z3Z9-(6S,7R)-epo-18:Hy (2) (Fig. 1B) have been synthesized successfully in a previous study by the current authors including electroantennogram responses (Yu et al. 2017). Moreover identification, characterization of odorant binding proteins, and their expression patterns from *E. obliqua* also have been accomplished (Sun et al. 2017). The *Eob*/PBP1 protein previously has been found to be highly expressed in antennae of male moths, and has provided us an opportunity to understand the sex pheromone recognition in this species (Ma et al. 2016a). We report here on results from electroantennograms and field trap collections to examine the attractiveness of Z3Z9-6,7-epo-18:Hy enantiomers of female *E. obliqua* to males. Furthermore, a homology modeling method was used to construct structures of *Eob*/PBP1 protein and molecular docking used to investigate the binding affinities of the different sex pheromone enantiomers.

## Materials and Methods

### INSECTS

Moths were collected by sweep net from Dong-zhi County (30.1000°N, 117.0200°E), Anhui Province, People's Republic of China, and identified as *E. obliqua* (Yang et al. 2015). Insects were maintained in a plastic jar (240 mL) covered with nylon mesh and reared at 25 °C, 60% to 70% RH, and a 14:10 (L:D) photoperiod. Eggs collected from

females were very carefully transferred to another jar, and maintained in dry conditions until hatching. Hatched larvae were reared on fresh tea leaves under ambient environmental conditions in our laboratory. After pupation, pupae were sexed and maintained on moist sand (10:1 sand:water) until emergence. Adults that emerged were maintained separately in 240 mL plastic jars (Fukang Plastic Co. Ltd., Dongguan, Guangdong, China) with 10% honey soaked in cotton ad libitum.

### CHEMICALS

Z3Z6Z9-18:Hy (a triene, chemical purity > 98%) was prepared from linolenic acid as described previously by Liu et al. (1994). A mixture of 3 corresponding racemic monoepoxydienes of this triene was prepared by oxidation with *m*-chloroperbenzoic acid. Flash chromatography was performed on silica gel to obtain racemic Z3Z9-6,7-epo-18:Hy containing hexane:benzene (5:1). Enantiomers of Z3Z9-6,7-epo-18:Hy were synthesized with chemical purity > 95% and *ee* > 80% from *cis*-2-butene-1,4-diol as previously reported by Yu et al. (2017).

### ELECTROANTENNAGRAM MEASUREMENT

Electroantennogram recordings were conducted with an EAG 2014 system (Syntech Co., Hilversum, Netherlands) consisting of a probe/micromanipulator (MP-15), a data acquisition interface box (serial IDAC-2), and a stimulus air controller (CS-55). Before electroantennogram measurements, the system was cleaned using an active charcoal purified stream of air. Male *E. obliqua* antennae were freshly excised from 1-d-old individuals, and mounted onto an antenna holder in the electroantennogram probe electrode (Spectro 360, Parker Laboratory, Princeton, New Jersey). A continuous air stream (flow rate, 2 L per min) created by the stimulus air controller was allowed to flow toward the antenna. Epoxide racemate and enantiomers of Z3Z9-6,7-epo-18:Hy were dissolved in *n*-hexane to produce 1 mg per mL solutions. It has been reported that (*Z*)-3-hexenol can elicit stable electroantennogram responses to antenna of male *E. obliqua* (Sun et al. 2014). Therefore, 10 μL (*Z*)-3-hexenol solution (0.005 μg per μL) was used as the reference stimuli and 10 μL of *n*-hexane used as the solvent control (Luo et al. 2017). Ten μL of each test solution was placed on a small strip of qualitative filter paper (0.5 × 2.5 cm, fast-speed type; Sinopharm Chemical Reagent Co., Ltd., Shanghai, China) and allowed to evaporate for 30 s. At this time, the strip was introduced into a Pasteur pipette (Jiechen Instruments Co. Ltd., Shanghai, China) (15 cm long) with its tip inserted into the tubing (8 mm in diam) up to 8 cm before the exit of the continuous air stream. The base of the pipette was connected to the pulsed air stream from the stimulus air controller. Stimuli with the test chemicals were provided to the insect antennae by puffing

pulsed air for 0.3 s (flow rate, 2 L per min) at the base of the pipette. Each recording was repeated 6 times with different antennae in the following sequence: n-hexane, reference compound, test compound (3 times), reference compound, and n-hexane. Control-adjusted electroantennogram responses of the stimuli were presented as proportional responses relative to the reference stimuli (Zhang et al. 2014). Mean data were analyzed using a 1-way analysis of variance (ANOVA) followed by LSD multiple comparison test. Differences were considered significant at  $P < 0.05$ .

## FIELD EXPERIMENTS

Field trapping experiments were conducted in a tea orchard in Xian-Ning (29.8700°N, 114.2800°E), Hubei, China, during May 2016. A pheromone-baited water pan trap was used in all field experiments (Fig. 2). Traps were baited with synthetic compounds in a 4:6 ratio of triene; 6,7-epoxide enantiomers were injected into a green rubber septa (6 mm OD; Pherobio Technology Co., Ltd., Beijing, China) that was suspended at 1.3 m from ground surface and spaced 10 m apart. Pheromone compounds were dissolved in distilled hexane to produce 1 mg per mL solutions and 10  $\mu$ L of total volume injected into a blank septa. Septa injected with distilled hexane were used as a control. All septa were stored at 4 °C until use. Each treatment had 8 replicates, and trap positions within each replicate were randomized. Trap catches were checked the next d. Statistical significance of the differences between treatments was assessed by 1-way ANOVA, followed by Duncan multiple range test ( $P < 0.05$ ).

## ALIGNMENT AND HOMOLOGY MODELING

A BLAST search (Altschul et al. 1990) of the amino acid sequence of *Eob*/PBP1 was conducted against the current Protein Data Bank database (<https://www.rcsb.org/>) to find structural templates. In this case, the crystalline structure of *Bmor*PBP (Protein Data Bank ID: 1DQE) at a resolution of 1.8 Å was chosen as the template for *Eob*/PBP1. Multiple sequence alignment was generated using the DNAMAN 8 software program (Lynnon Biosoft, San Ramon, California, USA). Using sequence alignments and the software template, Modeller 9.18 (Šali & Blundell 1993) was used to construct a 3-dimensional structure of *Eob*/PBP1 using a comparative protein modeling method (i.e., homology modeling). Ten candidates were generated by Modeller and the structure with the lowest DOPE (Discrete Optimized Protein Energy) score was chosen as a solution. To refine the protein, the relax module of program Rosetta was used to give final structure of the target protein (Das & Baker 2008).

## MOLECULAR DOCKING

The structures of female *E. obliqua* sex pheromone components, Z3Z9-6,7-epo-18:Hy (6S,7R- and 6R,7S- enantiomer), were created and optimized using chemBio3D Ultra 13 (PerkinElmer, Billerica, Massachusetts, USA). Molecules were applied with MM2 forcefield and saved in a Protein Data Bank format. The Protein Data Bank files were uploaded to AutoDockTools (The Scripps Research Institute, La Jolla, California, USA) and saved as PDBQT format for docking. The AutoDockTools



Fig. 2. Field trial of *Ectropis obliqua* Prout. (A) Water pan trap; (B) male moths captured.

software was used also to assign polar hydrogens and add Gasteiger charges to the protein, then the structure saved in PDBQT file format. After preparation of protein and ligands, structures were docked into the model of *Eob*/PBP1 using the AutoDock Vina 1.1.2 software (The Scripps Research Institute, La Jolla, California, USA). A docking grid with a size of 18 Å × 18 Å × 18 Å was used, and the grid center was set at 7.332 (X-axis), 22.972 (Y-axis), 3.629 (Z-axis). Free energy scores for ligands were calculated, and the lowest energy of the ligand-protein complexes was selected for analysis.


## Results

In the laboratory electroantennogram bioassays, male *E. obliqua* antennal response to 6S,7R-enantiomer was at the same level as epoxide racemate, but significantly greater than responses to the 6R,7S-enantiomer ( $P < 0.05$ ) (Fig. 3). The results suggest that (6R,7S)-enantiomer may not be the effective configuration.


In the field trapping experiment, the number of *E. obliqua* males captured in water pan traps was significantly greater for the 6S,7R-enantiomer (blend A) compared with 6R,7S-enantiomer (blend B) and control ( $P < 0.05$ ) (Fig. 4).

Electroantennogram and field trapping results suggested a chiral discrimination of 1 or more particular proteins at the molecular level. In order to understand the interaction between sex pheromone molecules and the protein, structural information of *Eob*/PBP1 was needed. Currently, there are no NMR or X-ray data for *Eob*/PBP1. Therefore, based on the templates and our alignment results, the 3D model of *Eob*/PBP1 was successfully generated. After refinement, the Verify Score of models by Profiles-3D was 65.13 (theoretical score 66.0542 and 29.7244) for *Eob*/PBP1, implying that overall quality of the predicted *Eob*/PBP1 structure was reliable. In addition, the Ramachandran plot of *Eob*/PBP1 revealed that 95.8% of the residues were in the favored region, 4.2% of the residues were in the allowed region, and none were in the disallowed region (Fig. 5C).

Results of the predicted 3D structure showed that *Eob*/PBP1 is compact and a stable globular protein that contains 6  $\alpha$ -helical domains:  $\alpha 1$  (residues Glu6–Glu26),  $\alpha 2$  (residues Ile32–Asn38),  $\alpha 3$  (residues Pro49–




**Fig. 3.** Electroantennogram responses of male *Ectropis obliqua* Prout antenna to racemate and enantiomers of Z3Z9-6,7-epo-18:Hy. Data are mean  $\pm$  SD ( $n = 6$ ) of male *E. obliqua* antennal responses by 2 enantiomers and racemate of Z3Z9-6,7-epo-18:Hy: Z3Z9-6S,7R-epoxy-18:Hy, Z3Z9-6R,7S-epoxy-18:Hy, racemate.




**Fig. 4.** Pan trap catches of male *Ectropis obliqua* Prout baited with binary blends of racemic and enantiomers of Z3Z9-6,7-epo-18:Hy with triene in Xian-Ning County, China. (A) Z3Z9-6S,7R-epoxy-18:H (6  $\mu$ g) + Z3Z6Z9-18:H (4  $\mu$ g); (B) Z3Z9-6R,7S-epoxy-18:H (6  $\mu$ g) + Z3Z6Z9-18:H (4  $\mu$ g); (C) blank lure (control). Data are mean  $\pm$  SD ( $n = 8$ ) of male *E. obliqua* trap catches in Xian-Ning County, China, May 2016 by 3 different combinations compared with water pan trap.

Leu62),  $\alpha 4$  (residues His76–His82),  $\alpha 5$  (residues Asp86–Ser103), and  $\alpha 6$  (residues Arg109–Leu127) (Fig. 5B). All 6 antiparallel  $\alpha$ -helices form a compact hydrophobic binding pocket. Although there still are several hydrophilic amino acids inside the binding cavity, Thr117 and Arg121 interacted with the epoxy ring of ligands. Not surprisingly, 3 pairs of disulphide bridges were found in the structure that was confirmed to be a common feature of insect pheromone binding proteins.

We found no discrimination between *Eob*/PBP1 and ligands in our docking experiments, although the results showed that orientation and conformation of Z3Z9-6,7-epo-18:Hy enantiomers were clear. Two enantiomers were anchored in the same binding cavity with opposite epoxy ring orientation (Fig. 6C), surrounded by hydrophobic amino acids. No



**Fig. 5.** Modeled 3D structure and validation of *Eob*/PBP1. (A) Sequence alignment of *Eob*/PBP1 and template 1DQE\_A.  $\alpha$ -helices are displayed as squiggles. Identical residues are highlighted in white letters with deep blue background. (B) Overall structure of the *Eob*/PBP1. Three disulfide bonds are in red. N-terminus, C-terminus, and  $\alpha$ -helices are labeled. Two potential key residues: Thr117 and Arg 121 are in orange. (C) Ramachandran plot of *Eob*/PBP1.



**Fig. 6.** Results of docking Z3Z9-6S,7R-epoxy-18:H and Z3Z9-6R,7S-epoxy-18:H into predicted model of *Eob/PBP1*. (A) Orientation of Z3Z9-6R,7S-epoxy-18:H in protein *Eob/PBP1*. Supposed key residues are in red and labeled. (B) Orientation of Z3Z9-6S,7R-epoxy-18:H in protein *Eob/PBP1*. Supposed key residues are in red and labeled. (C) Alignment of 2 enantiomers in different binding conformation. (D) The distance between oxygen in epoxy ring (Z3Z9-6S,7R-epoxy-18:H) and hydroxyl hydrogen in Thr 117 is labeled in yellow dotted line.

expected H-bonds were formed between Thr 117, Arg 121, and ligands (Figs. 6A, B). Van der Waals and hydrophobic interactions were important linkages between *Eob/PBP1* and epoxide ligands. When Z3Z9-6S,7R-epoxy-18:H was docked into the protein, the distance of oxygen in the epoxy ring and hydroxyl hydrogen in Thr 117 was found to be 4.5 Å (Fig. 6D). Moreover, interaction energies between ligands and *Eob/PBP1* confirm that Z3Z9-6S,7R-epoxy-18:H and Z3Z9-6R,7S-epoxy-18:H have similar binding affinity, though Z3Z9-6R,7S-epoxy-18:H has a lower energy value and is predicted to bind better than the opposite configuration (Table 1).

## Discussion

We found that the 6R,7S-enantiomer consistently produced a lower response and captured fewer male *E. obliqua*; this was caused by the presence of a “wrong” enantiomer. In this case, enantio-enriched but not enantio-pure Sharpless asymmetric epoxidation had occurred. Previously, epoxidation has been reported to account for 5% or more “incorrect” enantiomer synthesis (Katsuki & Sharpless 1980). We believe that a lower response of the 6R,7S-enantiomer could have been caused by the presence of trace amounts of 6S,7R-enantiomer. Mori (2007) investigated the stereochemistry-bioactivity relationships among pheromones and classified them into 10 categories. Based on our electroantennogram and field trapping results with that of Kenji’s investigation (Mori 2007), we could assume that 6S,7R-enantiomer is bioactive, and its opposite enantiomer does not inhibit or promote a response in male *E. obliqua*. This is the most common relationship with about 60% of chiral pheromones belonging to this group. Pheromones in this category can be used as their racemates in practical applica-

**Table 1.** Binding affinities of different enantiomers.

Ligands	Affinity(kcal per mol)
Z3Z9-6R,7S-epoxy-18:H	-6.3
Z3Z9-6S,7R-epoxy-18:H	-6.7

tions (Mori 2007). Recently, Z3,Z9-6,7-epoxy-18:H and Z3,Z6,Z9-18:H also has been identified as a component of sex pheromone in *Ectropis grisescens* Warren (Lepidoptera: Geometridae) (Ma et al. 2016b). In a related study, Z3,Z9-6,7-epoxy-19:H was identified as a key substance which caused pre-mating isolation of the 2 sibling moths: *E. grisescens* and *E. obliqua* (Luo et al. 2017). All of the reports were based on the racemate of Z3,Z9-6,7-epoxy-18:H. Considering the important role chirality plays, it is possible that these 2 sibling moths use different bioactive enantiomers as components of their sex pheromones that serve as an avenue of reproductive isolation.

In 2 recent silico studies by Sun et al. (2013b) and Liu et al. (2015b), male moths of *E. obliqua*, *Eob/PBP1* was proven to be specifically expressed (and dramatically detected) in their antennae, and shared a similar tissue expression pattern with pheromone binding protein genes in other moths. These results suggested a clear implication of male *E. obliqua* attractiveness in response to female-produced sex pheromones of this species (Sun et al. 2017). Therefore, a predicted 3D model could be built and used to investigate the binding mode of the different epoxide enantiomers.

In conclusion, silico interaction between ligands and *Eob/PBP1* should be confirmed by further X-ray crystallography studies. We believe that the knowledge we generated regarding the molecular interactions of *Eob/PBP1* supports the use of these sex pheromones for pest management in the field. Complementary experimental assays, such as competitive binding assays and site-directed mutagenesis, could validate the results obtained in our experiments to facilitate further research into the olfaction mechanism of *E. obliqua* males. Such investigations would serve to support alternative pest control strategies through design and development of novel pheromone analogs, as well as inhibitors of pheromone degradation.

## Acknowledgments

This study was supported by the National Natural Science Foundation of China (31200490), National Modern Agriculture Technology System (CARS-23), and Anhui Major Demonstration Project for Leading Talent Team on Tea Chemistry and Health. The senior author expresses his appreciation for the time spent at Mie University, Japan, with best wishes to Hirokata Katsuzaki and his team.

## References Cited

- Altschul SF, Gish W, Miller W, Myers EW, Lipman DJ. 1990. Basic local alignment search tool. *Journal of Molecular Biology* 215: 403–410.
- Das R, Baker D. 2008. Macromolecular modeling with rosetta. *Annual Review of Biochemistry* 77: 363–382.
- Gómez VRC, Carrasco JV. 2008. Morphological characteristics of antennal sensilla in *Talponia batesi* (Lepidoptera: Tortricidae). *Annals of the Entomological Society of America* 101: 181–188.
- Gräter F, Xu W, Leal W, Grubmüller H. 2006. Pheromone discrimination by the pheromone-binding protein of *Bombyx mori*. *Structure* 14: 1577–1586.
- Hazarika LK, Bhuyan M, Hazarika BN. 2009. Insect pests of tea and their management. *Annual Review of Entomology* 54: 267–284.
- Hazarika L, Puzari K, Wahab S. 2001. Biological control of tea pests, pp. 159–180. In Upadhyay RK, Mukerji KG, Chamola BP [eds.], *Biocontrol Potential and Its Exploitation in Sustainable Agriculture*. Springer, Boston, Massachusetts, USA.
- Honson N, Johnson MA, Oliver JE, Prestwich GD, Plettner E. 2003. Structure-activity studies with pheromone-binding proteins of the gypsy moth, *Lymantria dispar*. *Chemical Senses* 28: 479–489.
- Jin LY, Li ZQ, Zhang YN, Liu NY, Dong SL. 2014. Different roles suggested by sex-biased expression and pheromone binding affinity among three pheromone binding proteins in the pink rice borer, *Sesamia inferens* (Walker) (Lepidoptera: Noctuidae). *Journal of Insect Physiology* 66: 71–79.
- Katsuki T, Sharpless KB. 1980. The first practical method for asymmetric epoxidation. *Journal of the American Chemical Society* 102: 5974–5976.

- Li Z, Yao E, Luo Z, Liu T, Shang Z, Liu Z, Wang S. 1988. Preliminary structural elucidation of sex pheromone components of two geometrids in China: *Semiothisa cinerearia* Bremer & Crey and *Ectropis obliqua* Prout, pp. 29–32. In Proceedings of the China-Japan Seminar on Insect Semiochemicals, 6–8 Oct 1988, Beijing, China.
- Liu NY, He P, Dong SL. 2012. Binding properties of pheromone-binding protein 1 from the common cutworm *Spodoptera litura*. Comparative Biochemistry and Physiology Part B: Biochemistry and Molecular Biology 161: 295–302.
- Liu NY, Yang K, Liu Y, Xu W, Anderson A, Dong SL. 2015a. Two general-odorant binding proteins in *Spodoptera litura* are differentially tuned to sex pheromones and plant odorants. Comparative Biochemistry and Physiology Part A: Molecular & Integrative Physiology 180: 23–31.
- Liu NY, Yang F, Yang K, He P, Niu XH, Xu W, Anderson A, Dong SL. 2015b. Two subclasses of odorant-binding proteins in *Spodoptera exigua* display structural conservation and functional divergence. Insect Molecular Biology 24: 167–182.
- Liu T, Li Z, Luo Z, Jiang Y, Yao E. 1994. Synthesis of some bioactive components on *Ectropis obliqua* sex pheromone. Acta Scientiarum Naturalium Universitatis Nankaiensis 1: 82–85.
- Luo Z, Li Z, Cai X, Bian L, Chen Z. 2017. Evidence of premating isolation between two sibling moths: *Ectropis griseescens* and *Ectropis obliqua* (Lepidoptera: Geometridae). Journal of Economic Entomology 110: 2364–2370.
- Ma L, Li ZQ, Bian L, Cai XM, Luo ZX, Zhang YJ, Chen ZM. 2016a. Identification and comparative study of chemosensory genes related to host selection by legs transcriptome analysis in the tea geometrid *Ectropis obliqua*. PLoS One 11: e0149591. doi: 10.1371/journal.pone.0149591
- Ma T, Xiao Q, Yu YG, Wang C, Zhu CQ, Sun ZH, Chen XY, Wen XJ. 2016b. Analysis of tea geometrid (*Ectropis griseescens*) pheromone gland extracts using GC-EAD and GC×GC/TOFMS. Journal of Agricultural and Food Chemistry 64: 3161–3166.
- Mori K. 2007. Significance of chirality in pheromone science. Bioorganic & Medicinal Chemistry 15: 7505–7523.
- Mutis A, Palma R, Venthur H, Iturriaga-Vásquez P, Faundez-Parraguez M, Mella-Herrera R, Kontodimas D, Lobos C, Quiroz A. 2014. Molecular characterization and in silico analysis of the pheromone-binding protein of the European grapevine moth *Lobesia botrana* (Denis & Schiffermüller) (Lepidoptera, Tortricidae). Neotropical Entomology 43: 266–275.
- Pelosi P, Maida R. 1995. Odorant-binding proteins in insects. Comparative Biochemistry & Physiology Part B: Biochemistry and Molecular Biology 111: 503–514.
- Šali A, Blundell TL. 1993. Comparative protein modelling by satisfaction of spatial restraints. Journal of Molecular Biology 234: 779–815.
- Sun L, Mao TF, Zhang YX, Wu JJ, Bai JH, Zhang YN, Jiang XC, Yin KS, Guo YY, Zhang YJ. 2017. Characterization of candidate odorant-binding proteins and chemosensory proteins in the tea geometrid *Ectropis obliqua* Prout (Lepidoptera: Geometridae). Archives of Insect Biochemistry and Physiology 94: e21383. doi.org/10.1002/arch.21383
- Sun M, Liu Y, Walker WB, Liu C, Lin K, Gu S, Zhang Y, Zhou J, Wang G. 2013a. Identification and characterization of pheromone receptors and interplay between receptors and pheromone binding proteins in the diamondback moth, *Plutella xylostella*. PLoS One 8: e62098. doi: 10.1371/journal.pone.0062098
- Sun M, Liu Y, Wang G. 2013b. Expression patterns and binding properties of three pheromone binding proteins in the diamondback moth, *Plutella xylostella*. Journal of Insect Physiology 59: 46–55.
- Sun XL, Wang GC, Gao Y, Zhang XZ, Xin ZJ, Chen ZM. 2014. Volatiles emitted from tea plants infested by *Ectropis obliqua* larvae are attractive to conspecific moths. Journal of Chemical Ecology 40: 1080–1089.
- Venthur H, Mutis A, Zhou JJ, Quiroz A. 2014. Ligand binding and homology modelling of insect odorant-binding proteins. Physiological Entomology 39: 183–198.
- Yang Y, Zhang L, Guo F, Long Y, Wang Y, Wan X. 2015. Reidentification of sex pheromones of tea geometrid *Ectropis obliqua* Prout (Lepidoptera: Geometridae). Journal of Economic Entomology 109: 167–175.
- Yao EY, Li ZM, Lou ZQ, Shang ZZ, Yin KS, Hong BB. 1991. Report on structural elucidation of sex pheromone components of a tea pest (*Ectropis obliqua* Prout). Progress in Natural Science 6: 452–454.
- Ye GY, Xiao Q, Chen M, Chen XX, Yuan ZJ, Stanley DW, Hu C. 2014. Tea: biological control of insect and mite pests in China. Biological Control 68: 73–91.
- Yu J, Guo F, Yang YQ, Gao HH, Hou RY, Wan XC. 2017. Synthesis of the enantiomers of (3Z, 9Z)-cis-6, 7-epoxy-3, 9-octadecadiene, one of the major components of the sex pheromone of *Ectropis obliqua* Prout. Tetrahedron: Asymmetry 28: 758–761.
- Zhang GH, Yuan ZJ, Zhang CX, Yin KS, Tang MJ, Guo HW, Fu JY, Xiao Q. 2014. Detecting deep divergence in seventeen populations of tea geometrid (*Ectropis obliqua* Prout) in China by COI mtDNA and cross-breeding. PLoS One 9: e99373. doi: 10.1371/journal.pone.0099373