



Land Cover Change and its Environmental Impact in the Upper Reaches of the Yellow River, Northeast Qinghai-Tibetan Plateau

Authors: Yongnian, Zeng, Zhaodong, Feng, and Guangchao, Cao

Source: Mountain Research and Development, 23(4) : 353-361

Published By: International Mountain Society

URL: [https://doi.org/10.1659/0276-4741\(2003\)023\[0353:LCCAIE\]2.0.CO;2](https://doi.org/10.1659/0276-4741(2003)023[0353:LCCAIE]2.0.CO;2)

BioOne Complete (complete.BioOne.org) is a full-text database of 200 subscribed and open-access titles in the biological, ecological, and environmental sciences published by nonprofit societies, associations, museums, institutions, and presses.

Your use of this PDF, the BioOne Complete website, and all posted and associated content indicates your acceptance of BioOne's Terms of Use, available at www.bioone.org/terms-of-use.

Usage of BioOne Complete content is strictly limited to personal, educational, and non - commercial use. Commercial inquiries or rights and permissions requests should be directed to the individual publisher as copyright holder.

BioOne sees sustainable scholarly publishing as an inherently collaborative enterprise connecting authors, nonprofit publishers, academic institutions, research libraries, and research funders in the common goal of maximizing access to critical research.

Zeng Yongnian, Feng Zhaodong, and Cao Guangchao

Land Cover Change and its Environmental Impact in the Upper Reaches of the Yellow River, Northeast Qinghai-Tibetan Plateau

353



Many studies have focused on desertification processes in northern China in an attempt to understand desertification-related ecological problems. Desertification on the Qinghai-Tibetan Plateau has received

little attention, even though desertification resulting from overgrazing is a widespread phenomenon there. This study focuses on the Gonghe Basin in the upper reaches of the Yellow River, located on the northeast Qinghai-Tibetan Plateau, to shed light on desertification processes and associated environmental impacts during a 10-year period from 1987 to 1996. We first used 2 thematic mapper images to classify land cover for 2 summers (1987 and 1996) into 9 land cover types related to desertification. To assess the severity of desertification and to calculate the total soil carbon emission caused by desertification, we first defined severity by comparing land cover differences between 1987 and 1996 and then determined the differences in organic carbon content of the soil in grasslands during different stages of desertification between 1987 and 1996. The results showed that about 240 km² of grasslands was lost to agricultural encroachment and about 190 km² became sand-covered area between 1987 and 1996. During the same period, desertification affected 569.6 km² of grasslands, with desertification assessed as very severe for 41.8%, severe for 15.1%, and moderate for 43.1% of the area. The results also show that carbon emissions from grassland deterioration caused by desertification amounted to 2.06×10^6 tC during the 10-year period, the average annual emission rate being 0.206×10^6 tC.

Keywords: Land cover change; desertification; carbon emission; RS and GIS; Qinghai-Tibetan Plateau.

Peer reviewed: May 2003. **Accepted:** August 2003.

Introduction

Land degradation is a widespread phenomenon, with both ecological and economic impacts that threaten sustainable development throughout the world (Dregne 1990, 1995, 2002; Schlesinger et al 1990; Dregne and Chou 1992; Zhu and Cheng 1994; Daily 1995; Dregne et al 1996; Darkoh 1998; Shen et al 2001; Le Houerou 2002). As a specific expression of land degradation, desertification in arid and semiarid areas directly


affects nearly a quarter of the total land area of the world (ie, about 3.6 billion hectares) and threatens the very existence of about one sixth of the world's population (UNEP 1992). Desertification also degrades vulnerable ecosystems by altering regional hydrological cycles; this has an impact not only on socioeconomic development but also contributes to large-scale climate change (IGU 1998). Desertification has thus become a global concern not only in terms of ecological restoration, but also in terms of understanding the role it plays in global change (IGU 1998).

China is an extreme example, with severe land degradation resulting from a long history of inappropriate land management and increasing population pressure (Zhu and Liu 1981; Zhu and Cheng 1994; Li et al 2001; Shen et al 2001). Because of economic expansion and population increases in arid and semiarid China, desertification-related ecological degradation has been both so intensive and so extensive as to constitute a severe threat to regional sustainable development (Wang et al 1999b). Many studies have focused on desertification processes in northern China to understand desertification-related ecological problems (Zhu and Liu 1981; Zhu and Cheng 1994; Ci 1997; Wang et al 1998, 1999a,b). Desertification in the Qinghai-Tibetan Plateau has received little attention, even though desertification resulting from overgrazing there is widespread (Cheng 1998; Dong et al 1998; Niu 1999; Li et al 2001). The importance of understanding desertification in the Qinghai-Tibetan Plateau cannot be overemphasized, given the significance of the plateau in global climatic systems (Cheng et al 1997; Feng et al 1998; Zheng et al 2002). Equally important are the economic implications of desertification on the plateau: decreases in primary productivity resulting from desertification directly threaten the husbandry-based economy. Hence, there is a pressing need to monitor desertification processes on the plateau to assess environmental impacts and to provide a basis for sustainable economic planning.

The upper reaches of the Yellow River, located on the northeast Qinghai-Tibetan Plateau, consist of extensive grasslands of intrinsic ecological value that have long suffered from grassland degradation and desertification resulting from overgrazing (Cheng 1998; Dong et al 1998; Li et al 2000; Wang and Cheng 2000; Wang et al 2000, Zhang et al 2000). Although both intensive and extensive grassland degradation and desertification have been documented in the upper reaches of the Yellow River, few studies have focused on spatial and temporal variations in land cover and associated desertification.

The aim of this study is to detect changes in land cover and to assess its impacts on the environment in the Gonghe Basin, 1 of the main basins in the upper

FIGURE 1. A–C Population, livestock, and farmland statistics for the Gonghe Basin, 1985 through 1996. Farmland includes dry and irrigated farmland. (Source: SBHQ 1996)



reaches of the Yellow River, where grasslands became severely degraded from the 1950s to the 1980s (Dong et al 1993). It attempts to document land cover changes since 1985 using 2 sets of multispectral thematic mapper (TM) images (1987 and 1996) to detect trends in desertification in the late 1980s and early 1990s, when the climate entered a dry period (Cheng et al 1997; Ye et al 1997; Shang et al 2001) and the Chinese economy entered a new mode (privatization oriented). This study is specifically concerned with the analysis of the processes leading to land cover change and the assessment of related environmental impacts and socioeconomic consequences. Land cover change was the result of physical and anthropogenic factors. To depict land cover change since 1985, 2 sets of multispectral TM images seemed sufficient because the climate mode was generally the same (Cheng et al 1997; Ye et al 1997; Shang et al 2001), and the variables indicating the effects of human activities show nearly linear increasing trends during this period. Although there are some fluctuations in these variables, human and livestock populations and farmland acreage have generally increased since 1985 (Figure 1A–C).

The study area (Figure 2), Gonghe Basin ($35^{\circ}27'–36^{\circ}20'N$, $100^{\circ}–101^{\circ}20'E$), has a total area of about 11,500 km², with elevations ranging from 2400 to 3500 m. A desert-steppe landscape dominates low areas of the basin, whereas steppe landscape is found at high altitudes. The mean summer temperature is about 16°C, and the mean winter temperature is about –12°C. The mean annual precipitation varies from 311 mm in the northern (lower) part to 402 mm in the southern (higher) part of the basin (Dong et al 1993). The dominant sandy soil is low in inherent fertility, field water capacity, and organic matter content.

Methodology

Land cover categories

To detect changes in land cover during a 10-year period (1987–1996) using remote-sensing data, a new set of land cover categories was developed by modifying the National (China) Land Survey and Provincial (Qinghai) Land Survey standards applicable to the study area, taking full account of our field investigations. These categories are presented in hierarchical form. Three ecosystems are identified, first according to major land uses and then further divided into 9 land cover types that can be identified relatively quickly from remote-sensing data and linked to the severity of desertification (Table 1).

Data and processing

Two sets of cloud-free TM data were obtained, one in July 1987 and the other in July 1996, to capture the most

FIGURE 2 Map of the study area. (Map by Zeng Yongnian)




TABLE 1 Hierarchical land cover classification.

| Ecosystem | Land-cover type | Description of land-cover type |
|---------------------------|------------------------------|--|
| Bare ecosystem | Water | Salt lake, freshwater lake, reservoir, river |
| | Sand-covered area | All types of active sand dunes without grass cover |
| | Bare rock | Dry, denuded, rocky mountain terrain |
| Grassy ecosystem | Sparsely covered grassland | Vegetation coverage: 10–30% |
| | Moderately covered grassland | Vegetation coverage: 30–70% |
| | Densely covered grassland | Vegetation coverage: >70% |
| | Marshland | Wetlands in lowlands near foothills |
| Man-made ecosystem | Dry farmland | Farmland with rainfed agriculture |
| | Irrigated land | Irrigated farmland |


informative and representative land cover signatures in a year because July is the warmest and wettest season of the year. Because the climate in 1987 and 1996 was nearly the same, the 2 sets of TM data are representative of changes that occurred during this period. Ancillary data, either to facilitate TM data classification or to assess desertification processes (including environmental and socioeconomic factors), were acquired from the provincial census bureau and meteorological bureau.

The TM images were geometrically corrected, first by identifying ground control points on the original images and on the reference topographical maps and then by applying quadratic polynomial transformation equations. Subsequently, images were resampled by applying a nearest neighbor algorithm, and the root mean square error of <math><0.3</math> pixel (9 m) was obtained to ensure accuracy. The original TM bands 3, 4, and 5

were selected as classification bands, and supervised classification was executed using the maximum likelihood classification algorithm. Because the supervised maximum likelihood classification is based solely on spectral properties, the accuracy of classification results was not sufficient to meet our needs in detecting temporal variations in land cover. A postclassification technique, therefore, was used to improve accuracy. Specifically, thematic information (eg, on water, sand-covered area, and irrigated lands) was first extracted from TM data with the assistance of a TM-based vegetation index and DEM-based geomorphic analyses.

The thematically mapped land cover categories were then overlaid with the land cover categories classified using the maximum likelihood classification (ie, preclassification) to correct previously misclassified land cover categories. This process substantially

FIGURE 3, A AND B Land-cover patterns in Gonghe Basin, 1987 and 1996. (Based on TM data)



improved the results of preclassification. To ensure acceptance based on accuracy, we took sample points from the classified images in 1987 and 1996. Subsequently, direct comparison of land cover types was carried out on the basis of geographical location, with available primary data such as grassland maps in 1986 and 1997, land use maps in 1995, and our own field data. We then used a Kappa coefficient to evaluate accuracy. The Kappa coefficient was 91% for 1987 and

92.3% for 1996, ensuring our confidence in the accuracy of the classification.

Results and discussion

Changes in land cover

Our classification results show that 3 land cover categories changed significantly in the study area during the 10-year period: grassland, dry farmland, and sand-

TABLE 2 Land cover changes from 1987 to 1996.

| Type of land cover | Area in 1987 (km ²) | Area in 1996 (km ²) | Difference (km ²) | Rate of change (%) |
|------------------------------|---------------------------------|---------------------------------|-------------------------------|--------------------|
| Water | 207.4 | 227.9 | 20.5 | 9.9 |
| Bare rock | 19.1 | 19.1 | 0 | 0 |
| Sand-covered area | 1058.5 | 1247.4 | 188.9 | 17.8 |
| Marshland | 12.7 | 15.5 | 2.8 | 22 |
| Dry farmland | 362.1 | 549.7 | 187.6 | 51.8 |
| Irrigated land | 193.4 | 245.9 | 52.5 | 27.1 |
| Sparsely covered grassland | 181.3 | 152.1 | -29.2 | 16.1 |
| Moderately covered grassland | 2673.4 | 2590.2 | -83.2 | 3.1 |
| Densely covered grassland | 6787.7 | 6447.8 | -339.9 | 5.0 |
| Total area | 11,495.6 | 11,495.6 | | |

covered area. Specifically, the total area of dry farmlands increased from 362.1 km² in 1987 to 549.7 km² in 1996, and sand-covered areas increased from 1058.5 km² in 1987 to 1247.4 km² in 1996. The largest net change in terms of acreage (452.3 km²) was of grasslands, which decreased from 9642.4 km² in 1987 to 9190.1 km² in 1996. Among the 452.3 km² of grasslands lost, densely covered grasslands were affected most (339.9 km²), whereas moderately covered grasslands

(83.2 km²) and sparsely covered grasslands (29.2 km²) suffered much less (Figure 3A,B; Table 2).

To determine trends in land cover change and converted acreage from 1 category to another during the 10-year period in relation to desertification, we overlaid a 1987 land cover map with a 1996 land cover map. The results show that the sparsely covered grasslands were reduced by 71.3 km², all of which was transformed into sand-covered areas (Table 3).

TABLE 3 Matrix of land cover conversions from 1987 to 1996 (km²).

| Land-cover type | Water | Bare rock | Sand-covered area | Marshland | Dry farmland | Irrigated land | Sparsely covered grassland | Moderately covered grassland | Densely covered grassland | Total area (1996) |
|------------------------------|-------|-----------|-------------------|-----------|--------------|----------------|----------------------------|------------------------------|---------------------------|-------------------|
| Water | 196.4 | 0 | 7.5 | 0 | 0 | 0 | 0.3 | 10.8 | 12.9 | 227.9 |
| Bare rock | 0 | 19.1 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 19.1 |
| Sand-covered area | 3.4 | 0 | 1005.7 | 0 | 0 | 1.4 | 71.3 | 83.9 | 81.7 | 1247.4 |
| Marshland | 0.6 | 0 | 0 | 8.2 | 0 | 0 | 1 | 0.2 | 5.5 | 15.5 |
| Dry farmland | 0 | 0 | 0 | 0 | 331.7 | 0 | 0 | 1.9 | 216.1 | 549.7 |
| Irrigated land | 0.6 | 0 | 0 | 0 | 0 | 163.6 | 1.6 | 45.4 | 34.7 | 245.9 |
| Sparsely covered grassland | 0 | 0 | 12.3 | 0 | 0 | 0.4 | 53.6 | 81.2 | 4.6 | 152.1 |
| Moderately covered grassland | 5.1 | 0 | 12.3 | 0 | 0.7 | 11.2 | 50.9 | 2276.6 | 233.4 | 2590.2 |
| Densely covered grassland | 1.3 | 0 | 20.7 | 4.5 | 29.7 | 16.8 | 2.6 | 173.4 | 6198.8 | 6447.8 |
| Total area (1987) | 207.4 | 19.1 | 1058.5 | 12.7 | 362.1 | 193.4 | 180.3 | 2673.4 | 6787.7 | 1495.6 |

Moderately covered grasslands were reduced by 210.5 km², 83.9 km² of which was transformed into sand-covered areas, 45.4 km² into irrigated lands, and 81.2 km² into sparsely covered grasslands. Densely covered grasslands were reduced by 565.9 km², 81.7 km² of which was transformed into sand-covered areas, 216.1 km² into dry farmlands, 34.7 km² into irrigated lands, and 233.4 km² into moderately covered grassland. These data demonstrate that 299.7 km² of grasslands were lost and 556.1 km² were degraded to different degrees.

Overgrazing has been widely blamed for grassland deterioration in the study area (SBHQ 1996; OIT 1997). To further confirm the responsibility of overgrazing for grassland deterioration from 1987 to 1996, we looked at the Talatan area of the Gonghe Basin. The total livestock population far exceeded reasonable livestock density in terms of carrying capacity by about 23.5% in the Talatan area (OIT 1997) and by as much as 116% in some hot spots within the area (OIT 1997).

The second land cover/use category of environmental concern is dry farmlands, which increased by 187.6 km² during the period from 1987 to 1996, primarily as a result of cultivation of densely covered grasslands (Tables 2, 3). Cultivation of grassland has destroyed soil structure, increased soil erosion, and triggered desertification and sandstorm activity (Dong et al 1993; Li et al 2001). The increase in sand-covered areas is the ultimate consequence of land degradation. Our analysis shows that sand-covered area increased by 188.9 km² from 1987 to 1996, primarily as a result of overgrazing and cultivation. The increase in sand-covered area during the 10-year period indicates that overexploitation of natural resources in the study area has not been slowed when compared with the trend between the 1950s and the 1980s (Dong et al 1993).

Impacts on the environment

Compared with ecosystems in other climatic regimes, ecosystems in arid and semiarid regions are disproportionately prone to ecological damage from inappropriate forms of land use. According to our field investigations and data analysis, different types of land cover change, from better ecological conditions to worse ones, corresponded to different degrees of desertification severity in the study area.

For example, change from any kind of land cover to sand-covered area is considered very severe desertification. Change of densely covered or moderately covered grassland to sparsely covered grassland is categorized as severe desertification. Change from densely covered grassland to moderately covered grassland is classified as moderate desertification. Change from densely covered grassland (ie, >80%

coverage) to identifiably deteriorated, densely covered grassland (ie, <80% coverage) constitutes mild desertification.

Comparison of land cover between 1987 and 1996 shows that the total area of desertified lands increased by 569.6 km². Very severely desertified land increased at a mean annual rate of 23.8 km² during the 10-year period, severely desertified land at a rate of 8.6 km², and moderately desertified land at a rate of 24.5 km². The desertified acreage and the rate of desertification easily qualify the study area as 1 of the most desertified regions in the Qinghai-Tibetan Plateau, according to overall statistics on desertification in the entire plateau (Li et al 2001).

Terrestrial carbon and its sequestration have become issues of worldwide concern (Post et al 1982, 1990; Wisniewski and Sampson 1993; Fang et al 1996; Cao and Woodward 1998; Wang and Zhon 1999; Lal 2000; Follett 2001; Swift 2001). Land use change and land degradation have been shown to be significant in modulating atmospheric CO₂ levels by emitting or sequestering soil organic carbon (Heimann 1997). Recent studies have shown that in the past 30 years approximately 3.02 Pg C has been emitted from grassland soils on the Qinghai-Tibetan Plateau because of land-use change and grassland degradation (Wang et al 2002).

To assess the trends in soil organic carbon emission resulting from desertification during the 10-year period in the study area, we used the most common and simplest method based on land cover (Wang et al 1999). Organic carbon storage in the soil under a particular land cover in the specified area is a product of the mean soil organic carbon content per unit area and the total area of that particular land cover. The total organic carbon storage is then obtained by adding the organic carbon content of all soils under different forms of land cover in a specific area. Our study used relationships between land cover and associated soil organic carbon content established by field investigations (Shao et al 1988; Shen et al 1992). The soil organic carbon content was directly obtained from the soil organic matter content by multiplication, using the van Bemmelen conversion coefficient (0.58 g Cg⁻¹). The mean soil organic carbon per unit area was estimated by

$$C_i = 0.58 O_i W_i H_i \quad (1)$$

where C is mean soil organic carbon per unit area, i the type of desertified land, 0.58 the van Bemmelen Index, O the mean organic matter content in the soil under a specific land cover, W the mean soil bulk density, and H the mean soil analytical thickness. Because the soils in the plateau are generally thin (Dong et al 1993), 0.7 m of the soil analytical thickness (H) was proposed by Wang et al (2002) and adopted in this study.

TABLE 4 Soil carbon content by category of desertified land.

| Type of desertified land | Soil organic matter content (%) | Mean soil bulk density (g/cm ³) | Soil organic carbon content per unit area (kg/m ²) |
|--|---------------------------------|---|--|
| Identifiably deteriorated densely covered grassland (mildly desertified) | 1.189 | 1.48 | 7.144 |
| Moderately covered grasslands (moderately desertified) | 0.337 | 1.54 | 2.107 |
| Sparsely covered grasslands (severely desertified) | 0.148 | 1.55 | 0.931 |
| Sand-covered area (very severely desertified) | 0.094 | 1.56 | 0.595 |

TABLE 5 Desertification and associated carbon emission.

| Degree of desertification | Land cover conversion | Converted area (km ²) | Change of soil organic carbon content per unit area (10 ² tC) | Carbon emission (10 ² tC) |
|------------------------------------|---|-----------------------------------|--|--------------------------------------|
| Very severe desertification | Sparsely covered grasslands → very severely desertified land | 71.3 | 3.36 | 239.57 |
| | Moderately covered grasslands → very severely desertified land | 83.9 | 15.12 | 1268.57 |
| | Identifiably deteriorated, densely covered grassland → Very severely desertified land | 83.1 | 65.49 | 5442.22 |
| Severe desertification | Moderately covered grasslands → severely desertified land | 81.2 | 11.76 | 954.91 |
| | Identifiably deteriorated, densely covered grassland → severely desertified land | 5.0 | 62.13 | 310.65 |
| Moderate desertification | Identifiably deteriorated, densely covered grassland → moderately desertified land | 245.1 | 50.37 | 12,345.69 |
| Total | | 569.6 | | 20,561.61 |

Soil organic matter content and bulk densities under different degrees of desertification related to land cover change are listed in Table 4 (Shao et al 1988; Shen et al 1992), and soil carbon content per unit area is then calculated using Equation 1. To calculate total soil carbon emission, we first defined the degree of desertification by comparing the land cover differences between 1987 and 1996. We then found the differences in the soil organic carbon content of grass-

land with different degrees of desertification between 1987 and 1996 (Table 5). The estimated total carbon emission caused by desertification was 2.06×10^6 tC during the 10-year period, and the mean annual soil carbon emission flux was 0.206×10^6 tC, suggesting that desertification in the upper reaches of the Yellow River is partly responsible for significant emission of CO₂ into the atmosphere from the Qinghai-Tibetan Plateau.

ACKNOWLEDGMENTS

This research was supported by the National Science Foundation of China (40071066), the Education Ministry Foundation for Cadremen Teachers, the Knowledge Innovation Project of the Chinese Academy of Sciences (CAS) (KZCX1-10-06), and the National Key Basic Research Project (G2000048701).

AUTHORS

Zeng Yongnian

National Laboratory of Western China's Environmental System, Lanzhou University, Lanzhou 730000, People's Republic of China, and Center for Resources and Environment Research of Qinghai-Xizang (Tibetan) Plateau, Qinghai Normal University, Xining 810000, People's Republic of China. Zengyongnian@sina.com

Feng Zhaodong

National Laboratory of Western China's Environmental System, Lanzhou University, Lanzhou 730000, People's Republic of China, and Department of Earth and Environmental Studies, Montclair State University, New Jersey 07043, USA. fengzd@lzu.edu.cn

Cao Guangchao

Center for Resources and Environment Research of Qinghai-Xizang (Tibetan) Plateau, Qinghai Normal University, Xining 810000, People's Republic of China. gc.cao@yeah.net

REFERENCES

- Cao M, Woodward FI.** 1998. Dynamic responses of terrestrial ecosystem carbon cycling to global change. *Nature* 393:249–252.
- Cheng G.** 1998. Some understandings about the eco-environmental protection and buildings in the source region of Yangtze and Yellow Rivers [in Chinese]. *Advance in Earth Sciences* 13 (Supplement):1–5.
- Cheng G, Li P, Zhang X, Guo D, Li S.** 1997. *Influence of Climatic Changes on Snow Cover, Glaciers and Frozen Soils in China* [in Chinese]. Lanzhou: Gansu Cultural Publishing House.
- Ci L.** 1997. *Land Evaluation and Expert System for Combating Desertification: Case Study of the Mu Us Sandland* [in Chinese]. Beijing: China Forestry Publishing House.
- Daily GC.** 1995. Restoring value to the world's degraded lands. *Science* 269:350–354.
- Darkoh MBK.** 1998. The nature, causes and consequences of desertification in the drylands of Africa. *Land Degradation and Development* 9:1–20.
- Dong G, Gao S, Jin J, et al.** 1993. *Study of Desertification and Ways to Combat it in Gonghe Basin, Qinghai* [in Chinese]. Beijing: Science Press.
- Dong G, Yan P, Su Z.** 1998. The urgency of research and control of land desertification in the source area of Yangtze and Yellow Rivers [in Chinese]. *Advance in Earth Sciences* 13 (Supplement):32–36.
- Dregne HE.** 1990. Erosion and soil productivity in Africa. *Journal of Soil and Water Conservation* 45:431–436.
- Dregne HE.** 1995. Erosion and soil productivity in Australia and New Zealand. *Land Degradation and Rehabilitation* 6:71–78.
- Dregne HE.** 2002. Land degradation in the drylands. *Arid Land Research and Management* 16:99–132.
- Dregne HE, Chou N.** 1992. Global desertification and cost. In: Dregne HE, editor. *Degradation and Restoration of Arid Lands*. Lubbock, TX: Texas Tech University, pp 249–282.
- Dregne HE, Xiong Z, Xiong S.** 1996. Soil salinity in China. *Desertification Control Bulletin* 28:28–33.
- Fang J, Liu G, Xu S.** 1996. Carbon pool of terrestrial ecosystem in China. In: Wang G, Wen YM, editors. *Monitoring of Greenhouse Gas Concentration and Emission and Relevant Processes* [in Chinese]. Beijing: China Environment Science Press, pp 95–101.
- Feng S, Tang M, Wang D.** 1998. The new evidence about the Qinghai-Tibetan Plateau is triggering region of climate change in China [in Chinese]. *Chinese Science Bulletin* 43(6):633–636.
- Follett RF.** 2001. Soil management concepts and carbon sequestration. *Soil and Tillage Research* 61:77–92.
- Heimann M.** 1997. A review of the contemporary global carbon cycle and as seen a century ago by Arrhenius and Högbom. *Ambio* 26(1):17–24.
- [IGU] International Geographic Union.** 1998. Land use/cover change. *Land Use Policy* 15(2):165–166.
- Lal R.** 2000. Carbon sequestration in drylands. *Annals of Arid Zone* 39:1–10.
- Le Houerou HN.** 2002. "Man-made" deserts: Desertization processes and threats. *Arid Land Research and Management* 16:1–36.
- Li L, Zhang G, Wang Q, et al.** 2000. Study on evapotranspiration and its impact factors over Yellow River upper stream area [in Chinese]. *Advance in Earth Science* 15(3):256–259.
- Li S, Dong Y, Dong G, et al.** 2001. *Sandy Desertification Problem and Sustainable Development in Qinghai-Xizang (Tibet) Plateau* [in Chinese]. Beijing: China Tibetan Research Press.
- Niu Y.** 1999. The study of environment in the Plateau of Qinghai-Tibet [in Chinese]. *Progress in Geography* 18(2):163–171.
- [OIT] Office of the Item.** 1997. *Investigation Report on Grassland and Animal Husbandry* [in Chinese]. Integrated agriculture exploitation project in Hainan canton, Qinghai province, China. Hainan: OIT
- Post WM, Emanuel WR, Zinke PJ, Stangenberger AG.** 1982. Soil carbon pools and world life zones. *Nature* 298:156–159.
- Post WM, Peng TH, Emanuel WR, King AW, Dale VH, de Angelis DL.** 1990. The global carbon cycle. *American Scientist* 78:310–326.
- [SBHQ] Statistics Bureau of Hainan Canton, Qinghai Province.** 1996. *National Economic Statistics Yearbook During the Period from 1955–1996* [in Chinese]. Hainan: SBHQ.
- Schlesinger WH, Reynolds JF, Cunningham GL, Huenneke LF, Jarrell WM, Virginia RA, Whitford WG.** 1990. Biological feedbacks in global desertification. *Science* 247:1043–1048.
- Shang K, Dong G, Wang S, Yang D.** 2001. Response of climatic change in north China [in Chinese]. *Journal of Desert Research* 21(4):387–392.
- Shao L, Dong G, Lu F.** 1988. Developmental and adverse processes of steppe desertification and law of vegetation succession in Gonghe Basin [in Chinese]. *Journal of Desert Research* 8(1):30–40.
- Shen J, Dong G, Li C, Chen H.** 1992. Desertification and changes of matter content in the soil [in Chinese]. *Journal of Desert Research* 1:40–48.
- Shen Y, Zhang K, Wang X.** 2001. *Desertification* [in Chinese]. Beijing: China Environment Publishing House.
- Swift RS.** 2001. Sequestration of carbon by soil. *Soil Science* 166:858–871.
- [UNEP] United Nations Environment Programme.** 1992. *Status of Desertification and Implementation of the United Nations Plan of Action to*

Combat Desertification. Nairobi, Kenya: UNEP

Wang G, Cheng G. 2000. Eco-environmental changes and causative analysis in the source regions of the Yangtze and Yellow Rivers, China. *Environmentalist* 20(3):221–232.

Wang G, Qian J, Cheng G, Lai Y. 2002. Soil organic carbon pool of grassland soils on the Qinghai-Tibetan plateau and its global implication. *The Science of the Total Environment* 291:207–217.

Wang G, Shen Y, Cheng G. 2000. Eco-environmental changes and causal analysis in the source regions of the Yellow River [in Chinese]. *Journal of Glaciology and Geocryology* 22(3):200–206.

Wang S, Zhou C. 1999. Estimating soil carbon reservoir of terrestrial ecosystem in China [in Chinese]. *Geographical Research* 18(4):349–356.

Wang T, Toshiaki Imagawa, Wu W. 1999a. Methods for monitoring and assessing sandy desertification in North China [in Chinese]. *Journal of Desert Research* 19(1):6–11.

Wang T, Wu W, Wang X. 1998. Remote sensing monitoring and assessing sandy desertification: An example from the sandy desertification region of Northern China [in Chinese]. *Quaternary Sciences* 2:110–130.

Wang T, Zhao H, Xiao H. 1999b. Advances in desertification research of China [in Chinese]. *Journal of Desert Research* 19(4):299–311.

Wisniewski J, Sampson RN. 1993. *Terrestrial Biosphere Carbon Fluxes: Quantification of Sinks and Sources of CO₂*. Water Air Soil Pollution 70. Dordrecht: Kluwer Academic Publishers.

Ye J, Wang S, Chen Z. 1997. Climate variability in China and globalization during the first half of the 1990s. In: Ye DZ, editor. *Research of Climate Change and Climatic Affection of China*. Beijing: China Meteorological Press, pp 186–192.

Zhang G, Li L, Shi X, et al. 2000. Climatic changes over the upper Yellow River and its effects on water resources [in Chinese]. *Advances in Water Science* 11(3):277–283.

Zheng D, Lin Z, Zhang X. 2002. Progress in studies of Tibetan Plateau and global environmental change [in Chinese]. *Earth Science Frontiers* 9(1):95–102.

Zhu Z, Cheng G. 1994. *Sandy desertification in China* [in Chinese]. Beijing: Science Press.

Zhu Z, Liu S. 1981. Desertification and desertification control in northern China [in Chinese]. *Desertification Control Bulletin* 5:13–19.