

## **The Karyotype of the Yellow Dung Fly, *Scathophaga stercoraria*, a Model Organism in Studies of Sexual Selection**

Authors: Sbilordo, Sonja H., Martin, Oliver Y., and Ward, Paul I.

Source: Journal of Insect Science, 10(118) : 1-11

Published By: Entomological Society of America

URL: <https://doi.org/10.1673/031.010.11801>

---

BioOne Complete ([complete.BioOne.org](https://complete.BioOne.org)) is a full-text database of 200 subscribed and open-access titles in the biological, ecological, and environmental sciences published by nonprofit societies, associations, museums, institutions, and presses.

Your use of this PDF, the BioOne Complete website, and all posted and associated content indicates your acceptance of BioOne's Terms of Use, available at [www.bioone.org/terms-of-use](https://www.bioone.org/terms-of-use).

Usage of BioOne Complete content is strictly limited to personal, educational, and non - commercial use. Commercial inquiries or rights and permissions requests should be directed to the individual publisher as copyright holder.

---

BioOne sees sustainable scholarly publishing as an inherently collaborative enterprise connecting authors, nonprofit publishers, academic institutions, research libraries, and research funders in the common goal of maximizing access to critical research.



## The karyotype of the yellow dung fly, *Scathophaga stercoraria*, a model organism in studies of sexual selection

Sonja H. Sbilordo<sup>1,2a\*</sup>, Oliver Y. Martin<sup>2b</sup> and Paul I. Ward<sup>1</sup>

<sup>1</sup>Zoology Museum, University of Zürich, Winterthurerstrasse 190, CH-8057 Zürich, Switzerland

<sup>2</sup>Institute for Integrative Biology IBZ, Experimental Ecology, Universitätsstrasse 16, CH-8092 Zürich, Switzerland

### Abstract

Knowledge of karyotypical characteristics of a species is essential for understanding how sexually selected and sexually antagonistic traits evolve. The yellow dung fly *Scathophaga stercoraria* L. (Diptera: Scathophagidae) is an established model system for studies of sexual selection and sexual conflict, but karyotypical data are lacking to date. Here, the karyotype of *S. stercoraria* was characterized using conventional Giemsa-staining and C-banding techniques. The diploid chromosome set consists of 6 pairs of bi-armed meta- or submetacentric chromosomes. The sex chromosomes are the largest chromosomes and constitute 30% of the total length of the diploid set in females and about 25% in males. Males are the heterogametic sex, and the length of the Y chromosome is about three-quarters of that of the X chromosome. C-banding revealed that both sex chromosomes are largely heterochromatic. In contrast, in the five autosome pairs, heterochromatin is limited to narrow bands in the centromeric regions. This karyotypic information will help provide a more profound understanding of the inheritance of phenotypic variation in reproductive traits and the chromosomal locations of underlying genes.

**Keywords:** antagonistic, dimorphism, Diptera, heterochromatin, *Scathophaga*, sex chromosome, sexual conflict

**Abbreviation:** GLM, general linear model

**Correspondence:** <sup>a</sup> sonja.sbilordo@env.ethz.ch, <sup>b</sup> oliver.martin@env.ethz.ch, \*Corresponding author

**Received:** 23 July 2009, **Accepted:** 8 October 2009

**Copyright :** This is an open access paper. We use the Creative Commons Attribution 3.0 license that permits unrestricted use, provided that the paper is properly attributed.

**ISSN:** 1536-2442 | Vol. 10, Number 118

#### Cite this paper as:

Sbilordo SH, Martin OY, Ward PI. 2010. The karyotype of the yellow dung fly, *Scathophaga stercoraria*, a model organism in studies of sexual selection. *Journal of Insect Science* 10:118 available online: [insectscience.org/10.118](http://insectscience.org/10.118)

## Introduction

---

In dioecious animals with genetic sex determination, the chromosome sets of the two sexes differ in the appearance of the sex chromosomes. Depending on the species, either males or females may be the carriers of the heteromorphic sex chromosome pair (Reinhold 1998). In dipteran insects, the most common sex chromosome system is the XX/XY system. Here females possess two equivalent X chromosomes, while males possess an X and a Y chromosome that is usually degenerated and has hence lost much of the genetic activity (Charlesworth 1996; Rice 1996). The primary signals underlying sex determination may be the ratio of X chromosomes to autosomes, as in *Drosophila melanogaster*, for example. In other species, a specific male-determining factor is responsible for sex determination, e.g. as in *Musca domestica* and several *Chrysomya* species. This factor is often but not always located on the Y chromosome (Ullerich 1976; Hediger et al. 1998a; Schütt and Nöthiger 2000; Ullerich and Schöttke 2006).

Sex chromosome heteromorphism has major implications for the inheritance and expression of X linked genes. In males, the homology between the sex chromosomes is usually restricted to a small region. Therefore, the sex chromosomes can recombine in females, but not in males (Charlesworth 1996; Rice 1996). Males are hemizygous for X linked genes, so the amount of gene products of X linked genes have to be equalized in the two sexes. Several mechanisms have evolved to compensate the expression of X-linked genes. In dipteran insects such as *D. melanogaster* or *Sciara ocellaris*, dosage compensation is achieved by up-regulating

the transcription rates of the genes on the single X chromosome in males (Ruiz et al. 2000). Alternatively, if the sex chromosomes carry only very few genes, dosage compensation may not be necessary, as found, for example, in *M. domestica* (Schütt and Nöthiger 2000).

There is convincing theoretical and empirical evidence that phenotypic variation in sexually selected and sexually antagonistic traits is caused by X-chromosomal genes (Rice 1984; Reinhold 1998). Examples of X-linked expression of sexually selected traits include sex-limited traits such as the size of the ventral sperm receptacle in females (stalk-eyed fly, *Cyrtodiopsis dalmanni*: Johns and Wilkinson 2007) or sperm length in males (yellow dung fly, *Scathophaga stercoraria*: Ward 2000; stalk-eyed fly, *C. dalmanni*: Johns and Wilkinson 2007), as well as sexually dimorphic traits such as eyespan in the stalk-eyed fly *C. dalmanni* (Wolfenbarger and Wilkinson 2001).

Sex-linked expression of X-linked genes may be commensurate with the size of the X chromosome (Wolfenbarger and Wilkinson 2001). Indeed, the relationship between the relative length of the X chromosome in the set and the effect of X linked genes has previously been used to assess the influence of X chromosomal genes on expression of sexually selected traits (discussed in Wolfenbarger and Wilkinson 2001). Crucially, this appears to be based on the assumption that the gene contents of the X chromosome and the autosomes are similar. However, in many insect species, the X chromosome is largely heterochromatic, e.g. in several *Chrysomya* species and in *M. domestica* (Bedo 1991; Hediger et al 1998b; Ullerich and Schöttke 2006), so relative gene density is in fact

probably lower than on euchromatic autosomes, potentially concealing the relative importance of X linked genes. Additionally, dosage compensation by twofold hyperactivation of X-linked genes in males may double the magnitude of X-chromosome effects (Wolfenbarger and Wilkinson 2001). Thus, precise knowledge of karyotypical characteristics of a species, including the distribution of heterochromatin as well as the mechanism of dosage compensation, would be essential for a complete understanding of how phenotypic variation in sexually selected and/or sexually antagonistic traits evolves.

Since the pioneering work of Parker (Parker 1970a, 1970b), the yellow dung fly *Scathophaga stercoraria* L. (Diptera: Scathophagidae) has become a classic model organism for the study of pre- and postcopulatory sexual selection (Simmons 2001; Ward 2007; Bussière et al. 2009) and sexual conflict (Hosken et al. 2001; Martin et al. 2004; reviewed in Parker 2006). A large range of reproductive traits thought to be under sexual selection, such as the reproductive tract of the female (Hosken et al. 1999; Arthur et al. 2008), sperm size (Ward 2000) and sperm utilization patterns (Sbilordo et al. 2009), have been thoroughly investigated in this taxa. However, in contrast to the genome size (Garner 2002), the chromosomal constitution of this species remains unknown to the authors' knowledge. Here, a detailed description of the karyotype of the yellow dung fly is provided with special reference to the sex chromosome system and the morphology of the sex chromosomes, including the distribution of heterochromatin.

## Materials and Methods

Cytological analyses were carried out either

on neural ganglia of larvae or on reproductive tissues of adults. All animals used during this study were derived from laboratory stocks originally established from a population of *S. stercoraria* near Fehraltorf, Switzerland. Animals used were either from the first laboratory generation or derived from parental flies reared for more than one generation in the laboratory. Dung fly larvae and adults were housed in a climate chamber under standard conditions (20° C, 13:11 L:D). The larvae were reared in cow dung and used when they were between five and six days old. After eclosion, the adults were kept in vials (separated by sex, one to three flies per vial) and fed with sugar, water and adult *Drosophila* sp. *ad libitum*. Chromosome preparations of gonadal tissue were made six to twelve days after eclosion of the adults. Prior to killing, adult flies were cooled to 10° C for two days and then heated under a warm light for about 45 minutes to accelerate cell division (personal communication, M. Hediger-Niessen).

## Conventional Giemsa staining


The preparation of chromosome spreads was carried out according to El Agoze et al. (1992) with slight modifications. Briefly, the neural ganglia of the larvae were dissected in 1% sodium citrate, and the gonads of the adults were dissected in Ringer solution. Both tissues were treated hypotonically in 1% sodium citrate for ten to fifteen minutes. The hypotonic solution contained 0.05% of colchicine to increase the amount of metaphasic cells. After hypotonic treatment, the tissues were fixed in freshly prepared 3:1 ethanol-glacial acetic acid for five minutes. This fixative was replaced immediately before slide preparation. The fixed tissues were minced in 60% acetic acid, and two to four drops of the resulting cell suspension were dropped onto cleaned, ice-cold slides. These slides

were then placed on a hot plate at 45° C to dry. During evaporation of the solution, slides were moved gently to distribute the cell suspension evenly over the slide. Slides were then allowed to dry for at least two hours before they were stained with 10% Giemsa in a phosphate buffer solution (pH = 6.8) for four to six hours. After staining, slides were rinsed for one minute with deionized water. Chromosome counts were carried out on 92 well-spread metaphases prepared from the tissues derived from 21 individuals: three larvae, 13 males, and five females. Chromosome lengths were measured on ten good quality mitotic cell spreads from each sex derived from three males and three females, respectively. Measurements were performed on microscope images (100x) conveyed to a computer running the image analysis system KS 300, version 3.0 (Zeiss, www.zeiss.com). The chromosomes were classified morphologically based on the centromeric indices according to Guerra (1986). Digital images of the measured cell spreads were taken for illustration purposes. A multivariate general linear model was performed to compare the length of the two sex chromosomes within each sex. Residuals of the models did not deviate from normality (Kolmogorov-Smirnov tests:  $p > 0.42$ ).

### C-banding

Suitable mitotic chromosome spreads were

marked on slides freshly stained with Giemsa, and the slides subsequently were destained. Slides were passed through an ascending alcohol series (70% and 99% ethanol), washed in deionized water, and air dried. The C-banding technique (Sumner 1972) was then applied. Slides were treated with 0.2N HCl for 30 min at room temperature and rinsed with deionized water. The slides were then placed in a freshly prepared 5% aqueous solution of Ba(OH)<sub>2</sub> at 50° C for five minutes and washed with deionized water. After these treatments, slides were allowed to dry completely at room temperature for 48 h. The slides were then incubated in 2xSSC (0.3 M NaCl containing 0.03 M tri-sodium citrate) for 60 min at 60° C, rinsed with deionized water and stained with 10% Giemsa in phosphate buffer (pH = 6.8) for six hours. After a final washing with bi-distilled water, the slides were air-dried and mounted in Eukitt quick-hardening mounting medium (Fluka, www.sigmaaldrich.com). Digital photographs were taken with a Nikon Eclipse E600 microscope (www.nikon.com) fitted with a ProgRES C5 digital camera (www.progres-camera.com). Measurements were performed on the photograph using the software ImageJ (<http://rsbweb.nif.gov/ij/>). The schematic drawing of C-banded mitotic chromosomes of males was based on the assessment of banding patterns in 20 metaphase spreads.



**Figure 1.** Mitotic cell spreads of a male (A) and a female (B) *Scathophaga stercoraria*. The sex chromosomes marked with asterisks (\*) stain more darkly with Giemsa. High quality figures are available online.

## Results

The modal chromosome set of *S. stercoraria* was diploid and consisted of twelve chromosomes, comprising five pairs of autosomes and one pair of sex chromosomes (Figure 1). All chromosomes of the set were biarmed, metacentric, or submetacentric (Table 1).

The sex chromosomes were the two longest chromosomes of the set constituting 30% of the total length of the diploid set in females and 25% in males, respectively (Figure 2, Table 1). Both of the sex chromosomes were stained more intensely by Giemsa than the autosomes (Figures 1, 2, 3). The more intensive staining of sex chromosomes as compared with autosomes was consistent with previous work on other dipterans, e.g. the housefly *M. domestica* (El Agoze et al.1992). The homologous autosomes were closely associated with each other and arranged in pairs in most of the mitotic cell spreads. This was most

probably a result of somatic pairing during mitosis (see Metz 1916). The sex chromosomes were more loosely associated in the spreads (Figures 1, 3, 4). This observation coincides with the observation of mitotic chromosomes from ovarian tissue and somatic cells of females in *M. domestica* (Metz 1916) and the sex chromosomes of *Melophagus ovinus* males (Cooper 1941).

Males are the heterogametic sex in this species because the sex chromosomes were significantly dimorphic in size (GLM absolute length:  $F_{1, 19} = 27.90$ ,  $p < 0.001$ ; GLM relative length:  $F_{1, 19} = 60.82$ ,  $p < 0.001$ ). In females, sex chromosome length did not differ significantly (GLM absolute length:  $F_{1, 19} = 0.42$ ,  $p = 0.525$ ; GLM relative length:  $F_{1, 19} = 3.815$ ,  $p = 0.067$ ; (Figures 1, 2). C-banding revealed a predominantly heterochromatic character of both sex chromosomes. C-banding applied to highly condensed metaphase spreads resulted in dark staining of both sex



**Table 1.** Morphometric traits of the chromosomes based on 10 mitotic plates for each sex. The chromosomes were slightly more condensed in cell spreads of females than of males.

chromosome pair	1	2	3	4	5	X	Y
<b>male</b>							
chromosome length in $\mu\text{m}$	6.64 $\pm$ 0.22	5.68 $\pm$ 0.25	5.61 $\pm$ 0.38	4.80 $\pm$ 0.29	4.75 $\pm$ 0.43	10.36 $\pm$ 0.43	7.65 $\pm$ 0.28
relative length % of 2n	18.33 $\pm$ 0.39	15.63 $\pm$ 0.34	15.26 $\pm$ 0.45	13.11 $\pm$ 0.42	12.83 $\pm$ 0.63	14.27 $\pm$ 0.39	10.56 $\pm$ 0.28
arm ratio	1.41 $\pm$ 0.21	1.84 $\pm$ 0.11	1.64 $\pm$ 0.15	1.46 $\pm$ 0.08	1.39 $\pm$ 0.07	1.19 $\pm$ 0.04	1.16 $\pm$ 0.05
centromeric index	43.4 $\pm$ 2.3	35.8 $\pm$ 1.41	39.1 $\pm$ 2.19	40.9 $\pm$ 1.17	42.1 $\pm$ 1.18	45.7 $\pm$ 0.88	46.4 $\pm$ 0.97
<b>classification</b>	<b>m</b>	<b>sm</b>	<b>sm</b>	<b>m</b>	<b>m</b>	<b>m</b>	<b>m</b>
<b>female</b>							
chromosome length in $\mu\text{m}$	4.59 $\pm$ 0.30	3.76 $\pm$ 0.27	3.54 $\pm$ 0.31	3.32 $\pm$ 0.21	3.0 $\pm$ 0.19	7.84 $\pm$ 0.59	--
relative length % of 2n	17.68 $\pm$ 0.38	14.43 $\pm$ 0.36	13.49 $\pm$ 0.42	12.78 $\pm$ 0.24	11.63 $\pm$ 0.44	30.0 $\pm$ 0.75	--
arm ratio	1.32 $\pm$ 0.06	1.63 $\pm$ 0.05	1.60 $\pm$ 0.13	1.47 $\pm$ 0.09	1.56 $\pm$ 0.17	1.13 $\pm$ 0.03	--
centromeric index	43.32 $\pm$ 1.14	38.16 $\pm$ 0.71	39.27 $\pm$ 1.66	41.02 $\pm$ 1.52	40.13 $\pm$ 1.97	46.71 $\pm$ 0.43	--
<b>classification</b>	<b>m</b>	<b>sm</b>	<b>sm</b>	<b>m</b>	<b>m</b>	<b>m</b>	--

Data indicate means  $\pm$  SE  
\* according to Guerra 1986

chromosomes over almost their entire length with a poorly resolved banding pattern (Figure 3B). In prometaphase spreads with more extended chromosomes, the sex chromosomes also appeared more strongly stained than the pale C-banding negative parts of the autosomes (Figure 4). In these spreads, the Y chromosome additionally showed a particular staining pattern with large dark staining C-banding positive blocks on either side of the centromeric constriction, as well as on the distal regions of the chromosome arms. A


small interstitial C-banding negative region was visible in the large centromeric heterochromatin block of the long arm. In contrast to the Y chromosome, the X chromosome did not show any prominent dark staining blocks, and the banding pattern was less pronounced (Figures 4, 5). However, in most of the preparations, three small interstitial C-band negative regions were visible on the long arm, and one lighter stained band was visible on the short arm (Figures 4, 5). The observed C-banding pattern with dark blocks and less intense




background staining was similar to those previously observed on extended sex chromosomes of the screwworm *Chrysomya bezziana* (Bedo 1991). Differently stained bands may reflect intrachromosomal differences in the heterochromatin constitution of the sex chromosomes (see Hediger et al. 1998b). The C-banding pattern of autosomes was limited to a single narrow band in the centromeric region in each of the five autosome pairs, regardless of the degree of chromosome condensation (Figure 3B, 4,

5). These C-bands were not consistently visible on every autosome in all spreads. This was probably due to variation in the staining procedure as has been proposed for other insects (Bedo 1991; El Agoze et al. 1992; Ullerich and Schöttke 2006).

Interestingly, one phenotypic female with fully developed ovaries possessed 3 X chromosomes, an aneuploidy observed in all cell spreads stemming from this female (Figure 3). The absolute length of the three sex chromosomes did not differ



**Figure 4.** C-banding of a representative mitotic cell spread of a male *Scathophaga stercoraria*. Both sex chromosomes are more darkly stained than the C-band negative regions of the autosomes, and the Y chromosome additionally exhibits extensive darkly stained blocks. The autosomes are C-banding negative except for narrow bands in the centromeric regions. High quality figures are available online.



**Figure 5.** Diagrammatic representation of the C-banding idiogram of a male *Scathophaga stercoraria*. Strongest labeling is depicted in black, weaker labeling in grey and C-banding negative regions are depicted in white. High quality figures are available online.

significantly (GLM:  $F_{2, 26} = 0.44$ ,  $p = 0.650$ ).

## Discussion

Male heterozygosity for sex chromosomes has been proposed for the yellow dung fly in a previous study (Ward 2000), and this could be confirmed based on the karyological data presented here. *S. stercoraria* has five pairs of autosomes and two sex chromosomes, and there is significant size dimorphism in the sex chromosomes of males. A diploid chromosome complement of 12 chromosomes is concordant with other fly species, e.g. the housefly *M. domestica* (El Agoze et al. 1992), the olive fruit fly *Bactrocera oleae* (Mavragani-Tsipidou 2002), and several blow fly species e.g. *Chrysomya megacephala* and *C. putoria* (Ullerich 1976; Parise-Maltempi and Avancini 2001), *C. chloropyga*, *C. varipes* (Ullerich 1976) and *C. bezziana* (Bedo 1991). While all these species have the same chromosome number, there exists striking diversity in the length of the sex chromosomes. In *B. oleae*, *C. megacephala*, *C. chloropyga*, and *C. varipes*, the sex chromosomes are the two shortest chromosomes of the complement (Ullerich 1976; Parise-Maltempi and Avancini 2001; Mavragani-Tsipidou 2002). In *M. domestica* and *C. putoria* the X chromosomes are approximately autosomal size while the Y chromosome is the smallest chromosome of the set (El Agoze et al. 1992; Parise-Maltempi and Avancini 2001). In marked contrast to all these species, the karyotypes of *C. bezziana* (Bedo 1991) and *S. stercoraria* are dominated by the sex chromosomes, with both being much larger than the autosomes. These few examples indicate that the phylogenetic relationship between species is a very poor predictor for the relative

length of the sex chromosomes within species. Sex chromosomes are thought to evolve very rapidly under antagonistic sexual selection (Van Doorn and Kirkpatrick 2007), and it has been proposed that the influence of sex-linked genes on polygenic sexual dimorphic traits is approximately proportional to the length of the X-chromosome (Lande 1987). However, the X chromosomes are partially or largely heterochromatic in all of these species (Ullerich 1976; Bedo 1991; Hediger et al. 1998b; Parise-Maltempi and Avancini 2001). Therefore, it would be worthwhile to investigate whether observed differences in sex chromosome length between closely related species coincide with divergence in sexually dimorphic traits and/or mating system or whether the length of sex chromosomes solely reflects different proportions of heterochromatin.

One phenotypic female had three sex chromosomes of approximately equal size, which made it reasonable to assume that these three chromosomes were all X chromosomes. Sex chromosome aneuploidies are relatively common in animals, and they most often produce minor abnormalities, although in some cases the carriers are sterile (reviewed in Sumner 2003). Insects with aneuploid sex chromosome complements have been reported for several other Dipteran species e.g. XXY males in *C. chloropyga* and *M. domestica* (Ullerich 1976; Hediger et al. 1998a), XO females in *C. chloropyga* (Ullerich 1976) and an XXX female in *Phormia regina* (Ullerich 1961). The occurrence of aneuploid sex chromosome sets may indicate that sex in the concerned species is not determined by the X:autosome ratio as in *D. melanogaster* (Schütt and Nöthiger 2000), but rather by a male-determining factor (Ullerich 1976).

In conclusion, the evolution of sex chromosomes and their involvement in sexual selection and conflict are highly active fields of research (Arnqvist and Rowe 2005). These rapidly evolving chromosomes are considered to be evolutionary hotspots with a central role in the speciation process (see e.g. Presgraves 2008). A range of phenotypic traits likely to be under sexual selection such as body size (Simmons and Ward 1991) or copulation behavior of males (Mühlhäuser et al. 1996) have already been shown to be partially genetically determined in *S. stercoraria*. Now, with more precise knowledge concerning sex chromosomes, such studies could use this as foundations to provide a more complete picture of sexual selection in this important model organism.

### Acknowledgments

We would like to dedicate this article to the memory of Paul Ward and thank him for all the encouragement he offered. His keen interest in the yellow dung fly model system and postcopulatory sexual selection will be missed. Unfortunately, Paul died too early to contribute beyond the early stages, but we hope that the paper would still meet with his scientific approval. We further thank Monika Hediger-Niessen for providing a wealth of useful information on slide preparation and Georg Ribi, Marco Demont, Martin Schäfer, and an anonymous reviewer for very helpful suggestions.

### References

Arnqvist G, Rowe L. 2005. *Sexual conflict*. Princeton University Press.

Arthur BI, Sbilordo SH, Pemberton AJ, Ward PI. 2008. The anatomy of fertilization in the yellow dung fly *Scathophaga*

*stercoraria*. *Journal of Morphology* 269(5): 630-637.

Bedo DG. 1991. Cytological characterisation of heterochromatin in mitotic and meiotic chromosomes of the old world screwworm fly, *Chrysomya bezziana* (Diptera: Calliphoridae). *Genome* 34(4): 631-637.

Bussière L, Demont M, Pemberton A, Hall M, Ward PI. 2009. The assessment of insemination success in yellow dung flies using competitive PCR. *Molecular Ecology Resources* 10(2): 292-303.

Charlesworth B. 1996. The evolution of chromosomal sex determination and dosage compensation. *Current Biology* 6(2): 149-162.

Cooper KW. 1941. Bivalent structure in the fly, *Melophagus ovinus* L. (Pupipara, Hippoboscidae). *Proceedings of the National Academy of Sciences of the United States of America* 27(2): 109-114.

El Agoze M, Lemeunier F, Periquet G. 1992. Mitotic and salivary gland chromosome analyses in the *Musca domestica* L. (house fly) (Diptera: Muscidae). *Heredity* 69: 57-64.

Garner TWJ. 2002. Genome size and microsatellites: The effect of nuclear size on amplification potential. *Genome* 45(1): 212-215.

Guerra MD. 1986. Reviewing the chromosome nomenclature of Levan et. al. *Revista Brasileira de Genetica* 9(4): 741-743.

Hediger M, Minet AD, Niessen M, Schmidt R, Hilfiker-Kleiner D, Cakir S, Nöthiger R, Dübendorfer A. 1998a. The male-determining activity on the Y chromosome of the housefly (*Musca domestica* L.)

- consists of separable elements. *Genetics* 150(2): 651-661.
- Hediger M, Niessen M, Müller-Navia J, Nöthiger R, Dübendorfer A. 1998b. Distribution of heterochromatin on the mitotic chromosomes of *Musca domestica* L. in relation to the activity of male-determining factors. *Chromosoma* 107(4): 267-271.
- Hosken DJ, Garner TWJ, Ward PI. 2001. Sexual conflict selects for male and female reproductive characters. *Current Biology* 11(7): 489-493.
- Hosken DJ, Meyer EP, Ward PI. 1999. Internal female reproductive anatomy and genital interactions during copula in the yellow dung fly, *Scathophaga stercoraria* (Diptera: Scathophagidae). *Canadian Journal of Zoology* 77(12): 1975-1983.
- Johns PM, Wilkinson GS. 2007. X chromosome influences sperm length in the stalk-eyed fly *Cyrtodiopsis dalmanni*. *Heredity* 99(1): 56-61.
- Lande R. (1987). Genetic correlation between the sexes in the evolution of dimorphism and mating preference. In: Bradbury JW, Andersson MB, editors. *Sexual Selection: Testing the Alternatives*. pp. 83-94. John Wiley.
- Martin OY, Hosken DJ, Ward PI. 2004. Post-copulatory sexual selection and female fitness in *Scathophaga stercoraria*. *Proceedings of the Royal Society of London Series B* 271(1537): 353-359.
- Mavragani-Tsipidou P. 2002. Genetic and cytogenetic analysis of the olive fruit fly *Bactrocera oleae* (Diptera : Tephritidae). *Genetica* 116(1): 45-57.
- Metz CW. 1916. Chromosome studies on the Diptera II. The paired association of chromosomes in the Diptera, and its significance. *Journal of Experimental Zoology* 21(2): 213-279.
- Mühlhäuser C, Blankenhorn WU, Ward PI. 1996. The genetic component of copula duration in the yellow dung fly. *Animal Behaviour* 51: 1401-1407.
- Parise-Maltempi PP, Avancini RMP. 2001. C-banding and FISH in chromosomes of the blow flies *Chrysomya megacephala* and *Chrysomya putoria* (Diptera, Calliphoridae). *Memorias do Instituto Oswaldo Cruz* 96(3): 371-377.
- Parker GA. 2006. Sexual conflict over mating and fertilization: An overview. *Philosophical Transactions of the Royal Society of London Series B* 36 (1466): 235-259.
- Parker GA. 1970a. Sperm competition and its evolutionary consequences in insects. *Biological Reviews of the Cambridge Philosophical Society* 45(4): 525-567.
- Parker GA. 1970b. Sperm competition and its evolutionary effect on copula duration in the fly *Scatophaga stercoraria*. *Journal of Insect Physiology* 16(7): 1301-1328.
- Presgraves DC. 2008. Sex chromosomes and speciation in *Drosophila*. *Trends in Genetics* 24(7): 336-343.
- Reinhold K. 1998. Sex linkage among genes controlling sexually selected traits. *Behavioral Ecology and Sociobiology* 44(1): 1-7.
- Rice WR. 1984. Sex-chromosomes and the evolution of sexual dimorphism. *Evolution* 38(4): 735-742.
- Rice WR. 1996. Evolution of the Y sex chromosome in animals. *Bioscience* 46: 331-343.

- Ruiz MF, Esteban MR, Donoro C, Goday C, Sanchez L. 2000. Evolution of dosage compensation in diptera: The gene *maleless* implements dosage compensation in *Drosophila* (Brachycera suborder) but its homolog in *Sciara* (Nematocera suborder) appears to play no role in dosage compensation. *Genetics* 156(4): 1853-1865.
- Sbilordo SH, Schäfer MA, Ward PI. (2009). Sperm release and use at fertilization by yellow dung fly females (*Scathophaga stercoraria*). *Biological Journal of the Linnean Society* 98(3): 511-518.
- Schütt C, Nöthiger R. 2000. Structure, function and evolution of sex-determining systems in Dipteran insects. *Development* 127(4): 667-677.
- Simmons LW. 2001. *Sperm Competition and Its Evolutionary Consequences in the Insects*. Princeton University Press.
- Simmons LW, Ward PI. 1991. The heritability of sexually dimorphic traits in the yellow dung fly *Scathophaga stercoraria* (L.). *Journal of Evolutionary Biology* 4(4): 593-601.
- Sumner AT. 1972. Simple technique for demonstrating centromeric heterochromatin. *Experimental Cell Research* 75(1): 304-306.
- Sumner AT. 2003. *Chromosomes: Organization and Function*. Blackwell Science Ltd.
- Ullerich F-H. 1976. Chromosomenverhältnisse, konstitutives Heterochromatin und Geschlechtsbestimmung bei einigen Arten der Gattung *Chrysomya* (Calliphoridae, Diptera). *Chromosoma (Berl.)* 58: 113-136.
- Ullerich F-H. 1961. Geschlechtsbestimmung bei der Fliege *Phormia regina*. *Naturwissenschaften* 48: 559-560.
- Ullerich F-H, Schöttke M. 2006. Karyotypes, constitutive heterochromatin, and genomic DNA values in the blowfly genera *Chrysomya*, *Lucilia*, and *Protophormia* (Diptera : Calliphoridae). *Genome* 49 (6): 584-597.
- Van Doorn GS, Kirkpatrick M. 2007. Turnover of sex chromosomes induced by sexual conflict. *Nature* 449(7164): 909-912.
- Ward PI. 2007. Postcopulatory selection in the yellow dung fly *Scathophaga stercoraria* (L.) and the mate-now-choose-later mechanism of cryptic female choice. *Advances in the Study of Behavior* 37: 343-369.
- Ward PI. 2000. Sperm length is heritable and sex-linked in the yellow dung fly (*Scathophaga stercoraria*). *Journal of Zoology* 251: 349-353.
- Wolfenbarger LL, Wilkinson GS. 2001. Sex-linked expression of a sexually selected trait in the stalk-eyed fly, *Cyrtodiopsis dalmanni*. *Evolution* 55(1): 103-110.