

Influence of long-term exposure to simulated acid rain on development, reproduction and acaricide susceptibility of the carmine spider mite, *Tetranychus cinnabarinus*

Authors: Wang, Jin-Jun, Zhang, Jian-Ping, He, Lin, and Zhao, Zhi-Mo

Source: Journal of Insect Science, 6(19) : 1-8

Published By: Entomological Society of America

URL: https://doi.org/10.1673/2006_06_19.1

BioOne Complete (complete.BioOne.org) is a full-text database of 200 subscribed and open-access titles in the biological, ecological, and environmental sciences published by nonprofit societies, associations, museums, institutions, and presses.

Your use of this PDF, the BioOne Complete website, and all posted and associated content indicates your acceptance of BioOne's Terms of Use, available at www.bioone.org/terms-of-use.

Usage of BioOne Complete content is strictly limited to personal, educational, and non - commercial use. Commercial inquiries or rights and permissions requests should be directed to the individual publisher as copyright holder.

BioOne sees sustainable scholarly publishing as an inherently collaborative enterprise connecting authors, nonprofit publishers, academic institutions, research libraries, and research funders in the common goal of maximizing access to critical research.



Influence of long-term exposure to simulated acid rain on development, reproduction and acaricide susceptibility of the carmine spider mite, *Tetranychus cinnabarinus*

Jin-Jun Wang¹, Jian-Ping Zhang^{1,2}, Lin He¹ and Zhi-Mo Zhao¹

¹ Key Laboratory of Entomology and Pest Control Engineering, College of Plant Protection, Southwest University, Chongqing 400716, P. R. China

² Current Address: College of Agriculture, Shihezi University, Shihezi 832003, P. R. China

Abstract

Development, reproduction and acaricide susceptibility of *Tetranychus cinnabarinus* (Boisduvals) (Acari: Tetranychidae) were investigated after long-term (about 40 generations) exposure to various levels of acid rain; pH 2.5, 3.0, 4.0, and 5.6. Deionized water (pH 6.8) served as a control. The mites were reared on eggplant leaves at 28°C, 80%RH and a photoperiod of 14:10 (L:D) in the laboratory. The results showed that the duration of the immature stage was significantly affected by acid rain exposure. The shortest duration (8.90 days) was recorded for populations exposed to pH 5.6 acid rain, while the longest duration (9.37 days) occurred after exposure to pH 2.5 acid rain. Compared with the control population, adult longevity was shortened with an increase in acidity. Similarly, the oviposition duration was also shortened by an increase in acidity. Statistically, female fecundity did not differ significantly between pH 5.6, pH 4.0 and control populations, but did differ significantly between the control population and those exposed to pH 2.5 and pH 3.0 acid rain. This suggested that the mite suffered reproductive defects after long-term exposure to acid rain with higher acidity (pH 2.5 and 3.0). The intrinsic rate of increase among different populations was not significantly affected, but the net reproductive rate of populations exposed to pH 2.5 and 3.0 acid rain was significantly less than pH 4.0, 5.6, and control populations. Bioassay results showed that after long-term exposure to acid rain, susceptibility of the mites to two acaricides, dichlorvos and fenpropathrin, did not change significantly.

Keywords: *Tetranychus cinnabarinus*, acid rain, development, reproduction, susceptibility

Correspondence: jjwang7008@yahoo.com, zhangjp9507@yahoo.com.cn, helinok@163.net, zhaoxl@swau.edu.cn

Received: 10.7.2005 | Accepted: 2.2.2006 | Published: 21.9.2006

Copyright: Creative Commons Attribution 2.5 (<http://creativecommons.org/licenses/by/2.5>)

ISSN: 1536-2442 | Volume 2006, Number 19

Cite this paper as:

Wang J, Zhang J, He L, Zhao Z. 2006. Influence of long-term exposure to simulated acid rain on development, reproduction and acaricide susceptibility of the carmine spider mite, *Tetranychus cinnabarinus*. 8pp. *Journal of Insect Science* 6:19, available online: insectscience.org/6.19

Introduction

The threat of acid rain to the global biosphere has become an environmental problem of worldwide importance. Increased acidity of many lakes and soils, alteration of the bio-communities' equilibria, and decline of forests have been shown to be related to acid rain (Shaefer 1992; Driscoll et al. 2001; Menz and Seip 2004). The impact of acid deposition, usually in the form of simulated acidic rain, on vegetation, including the direct (primary) and indirect (secondary) effects through the influence on soils, has been well documented (Cowling 1982; Lee 1982; Linthurst et al. 1982; Treshow 1984; Shriner 1986; Treshow and Anderson 1989; Dai et al. 1998; Menz and Seip 2004). Acid deposition also influences host-plant susceptibility and suitability to insect herbivores (Bedford 1986; Riemer and Whittaker 1989; Heliövaara and Väisänen 1993; Zhang et al. 2005). While there are numerous studies on the effects of air pollution on insects, the majority of these investigations were conducted in forest systems under the influence of particulate and/or gaseous pollutants. Most references to insect population dynamics state that air pollution is one of the predisposing factors that influences the susceptibility of forest stands to outbreaks (Grodzki et al. 2004). A number of studies have been done concerning effects of air pollution on phytophagous insects in agricultural systems (Zhang et al. 2004).

Regions that have been most affected by acidic deposition include Europe, eastern North America, and Southeast Asia, especially central and southern China (Kuylenstierna et al. 2001). Sulphur emissions have played the dominant role in acidic deposition in these regions. Sulphur emissions in China decreased in the late 1990s, increased from 1999 to 2000, and remained stable up to 2002 (Li and Gao 2002). Chongqing is a city in China that is seriously polluted by acid rain. The average frequency of acid precipitation in Chongqing was about 70–90% and the lowest pH value of acid precipitation was 3.6. Acid rain in Chongqing has caused soil acidification, reduced forested areas and crop yield, and intensified insect pest infestation (Zhao et al. 1999). Based on our investigation, the infestation of the carmine spider mite, *Tetranychus cinnabarinus* (Boisduval), a serious mite pest of vegetable and fibre crops, is positively correlated with the frequency of acid precipitation. Our previous work has focused upon determining the biological and physiological effects of acid rain on the development and reproduction

of *T. cinnabarinus* (Zhao et al. 1999; Wang et al. 2004; Zhang et al. 2004). The results indicated that there was no acute lethality of sprayed simulated acid rain of pH 4.0–3.0 on this mite. However, the development and reproduction of the mite was affected by the direct application of acid rain. Mite population dynamics and host injury increased significantly after acid rain (pH 4.0) was applied to both on the mites and the host plant. Biochemical analysis showed that the peroxidase, superoxide dismutase, catalase and acid phosphatase might be relevant to population changes of mites under acid rain pressure.

Demographic toxicology incorporating life table parameters is applied in meaningful comparative studies. Life table parameters for populations unexposed or exposed to various concentrations of a toxicant or pollutant can be incorporated, and ensuing population responses can be compared. This approach combines ecological and toxicological parameters, resulting in better predictions about the effects of toxicants at the population level. Ultimately, the power of the intrinsic rate of increase (r_m) as an ecologically meaningful bioassay parameter for toxicology studies lies in the fact that this statistic is based on both survivorship (l_x) and fecundity (m_x). Estimates of r_m are environmentally specific and must be viewed in the context of the experimental conditions used (Stark et al. 2003). The relative fitness of a population can be evaluated by the average population size when the population has achieved an equilibrium by natural selection after several generations in a given environment (Carson 1961). To date, the study on the long-term impacts of acid rain on herbivorous insect and mite populations (i.e. impact on insect and mite fitness over several generations) is lacking. The present study was undertaken to evaluate the effects of long-term exposure to acid rain on the development, reproduction and pesticide susceptibility of *T. cinnabarinus*.

Materials and Methods

Mites

The stock culture of carmine spider mite, *T. cinnabarinus* (Boisduval), was originally collected from the cowpea bean (*Vigna sinensis* Endlicher) in Chongqing, China. This culture was maintained on potted kidney bean (*Phaseolus vulgaris* L.) in a insect rearing room at $28 \pm 1^\circ\text{C}$, 75–80%RH, and a photoperiod of 14:10 (L:D). This colony was maintained for more than two years before acid rain treatment. Voucher specimens were deposited

in the insect collection of the Southwest University, Chongqing, P. R. China.

Simulated acid rain treatment

An initial solution was created by adding to deionized water the mean final concentration of 5.35mg/L $(\text{NH}_4)_2\text{SO}_4$, 0.36 mg/L Na_2SO_4 , 0.97 mg/L MgSO_4 , 6.80 mg/L CaSO_4 , 1.52 mg/L KNO_3 , 0.51 mg/L NaNO_3 , 0.77 mg/L MgCl_2 , and 0.29 mg/L NaF as determined from data obtained by the Tongyuan Monitoring Station, Chongqing for the ionic composition of rain (Wang and Qin 1989). An amount of H_2SO_4 was added to bring the initial solution to pH 5.6, 4.0, 3.0 or 2.5. Deionized water, pH6.8, was used as a control. The pH values of all solutions were checked with a pHs-4C meter (Chengdu Instrument Inc., Chengdu, China).

Host plant

Eggplants (*Solanum melongena* L.) grown in pots (7.5 cm height, 4.5 cm diameter) were used for mite rearing. Plants were replaced every 30 days throughout the study. The plants were fertilized with a controlled Hoagland nutritional solution (Hoagland and Arnon 1950) and watered with deionized water as required. The experiment was conducted in the insect rearing room described above. For experimental use the mites were transferred from the stock colony to egg plants. In all experiments, simulated acid rain of different pH values were applied to the potted eggplant with mites from a portable overhead sprinkler system at a height of 75 cm from the ground level. The plants were sprayed every two days with 150 ml of simulated rain per plant each time. The mite colony was transferred to new plants every 30 days or shorter throughout the study. Similar mite population densities were maintained for all acid rain treatments. After one year and three months (about 40 generations), the mites from each treatment were used for developmental, reproductive and acaricide susceptibility studies.

Development and reproduction of mite

This experiment was conducted to test the effects of long-term exposure to acid rain with different acidity on the development and reproduction of the mite. In this experiment, leaves of eggplants sprayed with deionized water were used, and all leaves used were from the same phyllotaxis. Petri dishes (12 x 1.5 cm), each with a piece of water-soaked foam plastic inside, were used as observation units. The water-soaked foam plastic was covered with a layer of Whatman No. 1 filter paper. Five female and two male adult mites from

each treatment were transferred to a detached eggplant leaf placed on the filter paper. The mites were held at 28°C for a 4 hour oviposition period, after which the adults were removed. Petri dishes were placed in growth chambers (Jiangnan Instrument Inc., China) at 28°C, 80%RH, and a photoperiod of 14:10 (L:D). Eggs were checked twice daily and after hatching, 20 larval mites were transferred individually using a camel's-hair brush to a small area isolated by strips of absorbent cotton on the lower surface of a detached leaf. Individual mites were checked twice daily for ecdysis (i.e., for life stage and developmental period) and survivorship. The exuviae were counted to determine molting. Mites were transferred to new leaves every 2 or 3 days throughout the study.

After eclosion, male and female adults were paired. Each pair was transferred daily to a fresh leaf disk and eggs laid were counted under a stereomicroscope. Observations were recorded daily until the last female died.

Bioassay

This experiment was carried out to test the relationship between the susceptibility of the mites to acid rain and acaricides. The susceptibility of the mites after exposure to acid rain treatment to two acaricides, dichlorvos and fenpropathrin, purchased from Chongqing Pesticide-Chemical Company, P. R. China, was assessed following the recommended methods of the FAO (Dennehy et al. 1983). Various concentrations of each insecticide were tested until a satisfactory range was ascertained. Six concentrations were used in the final analysis. Each population was bioassayed by the slide-dip method using six treatments of acaricides, and a control. The acaricides were diluted with water to the required concentrations, and a slide with about 30 adult female spider mites, 3–5 days old, was dipped into the solution for 5s then left to dry. Each treatment was replicated three times. The slides were left in an incubator with temperature and humidity controlled at 25°C and 75% relative humidity. Mortality was assessed after 24 h. Mites that did not move after stimulation from a camel's hair brush were scored as dead.

Data analysis

The life history parameters of mites on eggplant leaves were analyzed among different populations using the general linear model (GLM) procedure of SAS (SAS Institute, 1988) and means were separated by least significant difference (LSD) test. Life table of different strain statistics were

calculated as described by Hulting et al. (1990). The differences in r_m values among populations were also analyzed using Newman-Keul sequential tests (Sokal and Rohlf, 1981) based on jackknife estimates of variance for r_m s (Meyer et al., 1986). For any difference between 2 r_m s from the sequence, in which the r_m s were arrayed in order of magnitude, to be significant at the α level, it must be equal to or greater than:

$$LSR = Q_{\alpha[K.V]} \sqrt{S_{av}^2} \sqrt{\frac{n_i + n_j}{2n_i n_j}}$$

where K is the number of r_m s in the set whose range is tested, and V is the degrees of freedom. The n_i and n_j were sample sizes of the 2 r_m s, and $Q_{\alpha[K.V]}$ is a value from the table of the studentized range. S_{av}^2 is the weighted average variance of r_m and it is calculated as follows:

$$S_{av}^2 = \frac{\sum (n_i - 1) S_i^2}{\sum (n_i - 1)}$$

where a equals the number of r_m s to be tested, the sample size of the i th r_m is n_i . S_i^2 is the jackknife estimate of the variance for the i th r_m .

The bioassay results were corrected for control mortality using Abbott’s (1925) formula. According to the results of bioassay, the log-concentration against probit-mortality lines was developed, and the comparison of mortalities between the populations was made by the concentrations required for 50% mortality (LC₅₀).

Results

Immature development of *T. cinnabarinus*

After long-term exposure to acid rain treatment (about 40 generation), the developmental periods of immature stages of *T. cinnabarinus* varied significantly among populations exposed to different acid rain conditions (Table 1). For egg development, the longest duration was recorded for pH 4.0 acid rain exposure population (4.83 days), while the shortest was for pH 2.5 acid rain exposure populations ($F = 31.472$; $df = 4, 595$; $P < 0.0001$).

In contrast, the length of the larval stage in mites exposed to pH 5.6 and pH 4.0 treatments were both less than for the pH 6.8 control population, while the population exposed to pH 2.5 acid rain was longer than control population ($F = 17.822$; $df = 4, 549$; $P < 0.0001$). The length of the nymphal stage in mites exposed to the pH 2.5 treatment was significantly longer than those of the other four populations ($F = 10.490$; $df = 4, 439$; $P < 0.0001$). The long-term exposure to acid rain also affected the pre-oviposition period significantly ($F = 1.853$; $df = 4, 214$; $P < 0.0001$). The total length of immature stages of mites exposed to the pH 6.8, 5.6, 4.0, 3.0, and 2.5 treatments were 9.09, 8.9, 9.25, 8.92, and 9.37 days, respectively.

Reproduction and life table parameters of *T. cinnabarinus*

Mean female longevity was significantly different among the five populations ($F = 13.729$; $df = 4, 214$; $P < 0.0001$), and was longest (12.70 days) for the control population and shortest (8.83 days) for the population exposed to pH 2.5 acid rain (Table 2). More than 90% of the females began to lay eggs within 24 h after copulation and they continued laying eggs for their entire adult life under all acid rain treatments. However, the numbers of eggs laid per female and oviposition durations were all significantly affected by the different pH acid rain exposure history. In general, the mites exposed to acid rain laid fewer eggs and had shorter oviposition periods than those of control populations. The mean sex ratio for all populations was strongly biased toward females (81.66%). The oviposition duration of pH2.5 acid rain exposure population was shorter than that of other populations ($F = 14.745$; $df = 4, 214$; $P < 0.0001$).

The intrinsic rate of increase (r_m), net reproductive rate (R_o), mean generation time (MT), and population doubling time (t) and finite rate of increase (λ) were calculated for populations of *T. cinnabarinus* exposed to different pH acid rain (Table 3). The r_m values of different populations were not significantly different ($P > 0.05$). There were significant differences in R_o values between

Table 1. Developmental periods (days, M±SE) of immature stages of different *T. cinnabarinus* populations after exposure to acid rain for 40 generations.

Population	Egg (days)	Larva (days)	Nymph (days)	Pre-ovipositional period (days)
pH6.8	4.5167±0.0592b (120)	1.4017±0.0455b (117)	3.1707±0.0792a (82)	1.0303±0.0303ab (33)
pH5.6	4.6333±0.0487b (120)	1.2328±0.0394a (116)	3.0291±0.072a (103)	1.0698±0.0393a (43)
pH4.0	4.8333±0.0619a (120)	1.2400±0.0429a (100)	3.1750±0.0610a (80)	1.0000±0.0000ab (40)
pH3.0	4.2000±0.0716c(120)	1.4951±0.0532b (103)	3.2286±0.0582a (70)	0.9796±0.0204b (49)
pH2.5	4.0250±0.0465d (120)	1.7179±0.0567c(117)	3.6330±0.089b (109)	1.0556±0.0315ab (54)

Means within a column followed by the same letter are not significantly different ($P > 0.05$; GLM) in ANOVA (LSD).

Table 2. Fecundity and longevity (M±SE) of different *T. cinnabarinus* populations after exposure to acid rain for 40 generations

Population	n	Adult longevity	No. eggs laid per female	Sex ratio % (Female)	Oviposition duration (d)
pH6.8	33	12.6970±0.3928a	29.1212±1.6921a	82.27	8.9394±0.3691a
pH5.6	43	11.9302±0.4604ab	25.4419±1.4685ab	81.56	8.3256±0.4045a
pH4.0	40	11.9000±0.4526ab	29.7250±2.1074a	81.26	8.2000±0.5066a
pH3.0	49	10.9592±0.3299b	23.6939±1.4125b	80.76	7.8571±0.3583a
pH2.5	54	8.8333±0.4203c	18.9444±0.8931c	82.45	5.1852±0.3528b

Means within a column followed by the same letter are not significantly different ($P > 0.05$; GLM) in ANOVA (LSD).

mites exposed to pH 2.5 and 3.0 and the control population ($P < 0.05$), however, R_0 values of the other three populations were not significantly different from each other or from the control population ($P > 0.05$). The finite rate of increase (λ) of different populations was 1.16–1.17. The mean generation time of pH 4.0 was longest (14.24 days) and that of pH 2.5 was shortest (12.31 days). Population doubling time of different populations ranged from 4.48 to 4.66 days. The relative fitness (R_f , which is defined as the ratio of r_m values between acid rain exposed and control populations) of pH 5.6, pH 4.0 and pH 3.0 were 1.013, 1.033 and 1.07 respectively, while that of pH 2.5 was 0.993. In general, the relative fitness of different populations did not change significantly.

Dichlorvos and fenpropathrin susceptibility of *T. cinnabarinus*

The susceptibility of *T. cinnabarinus* to dichlorvos and fenpropathrin are shown in Table 4. The slope of the probit lines showed that the populations were relatively homogeneous with regard to susceptibility. This suggested that after 40

generations of exposure to acid rain, the susceptibility of *T. cinnabarinus* to dichlorvos and fenpropathrin did not change significantly.

Discussion

The demographic toxicological approach is not only a measure of the total toxic effect, but also sub-lethal effects that are not perceptible can be evaluated. Results of several studies have indicated that sub-lethal effects can be very subtle and affect populations at concentrations lower than the traditional concentration response curve (Stark et al. 2003). Studying the effects of acidic water on populations of *Daphnia pulex*, Walton et al. 1982 found that 96-hour toxicity tests detected no effects of acidic water, but negative effects were easily discerned using an evaluation of the intrinsic rate of increase. Other studies reported that acid rain enhanced growth and development of insects (Palokangas et al. 1995; Redak et al. 1995; Stinner et al. 1988). Although there was no acute lethality of sprayed simulated acid rain on *T. cinnabarinus*,

Table 3. Comparison of life table parameters of different *T. cinnabarinus* population after exposure to acid rain for 40 generations

Population	n	r_m	R_0	λ	MT	t	R_f
pH6.8	33	0.150±0.005a	7.88±0.446a	1.16	13.74	4.61	1
pH5.6	43	0.152±0.004a	8.52±0.498a	1.16	14.1	4.56	1.013
pH4.0	40	0.155±0.004a	9.04±0.643a	1.17	14.24	4.48	1.033
pH3.0	49	0.154±0.004a	7.60±0.414ab	1.17	13.17	4.5	1.027
pH2.5	54	0.149±0.004a	6.24±0.294b	1.16	12.31	4.66	0.993

n , Number of females in analysis; r_m , jackknife estimate of the intrinsic rate of increase (per capita rate of population growth); R_0 , net reproduction rate; MT , mean generation time (in days); t , doubling time (in days) for population; λ , finite rate of increase. R_f , relative fitness (which is defined as the ratio of r_m values between acid rain exposed and control populations). Means within a column followed by the same letter are not significantly different at $P > 0.05$.

Table 4. Susceptibility of different *T. cinnabarinus* populations to two acaricides after exposure to acid rain for 40 generations

Populations	acaricides	slope	χ^2	LC ₅₀ ±SE (mg/L)	95%CL (mg/L)
pH6.8	dichlorvos	4.5856	1.0063	464.77±12.32a	488.92~440.62
	fenpropathrin	1.1405	1.481	1602.24±178.85a	1952.79~1251.69
pH5.6	dichlorvos	4.2948	1.4084	463.18±13.02a	488.70~450.68
	fenpropathrin	1.3673	9.8075	1777.83±173.50a	2117.89~1437.77
pH4.0	dichlorvos	4.2833	1.6366	470.16±13.28a	496.19~444.13
	fenpropathrin	1.2756	6.3203	1844.81±191.84a	2220.82~1468.80
pH3.0	dichlorvos	4.8076	2.4633	480.97±12.42a	505.31~456.63
	fenpropathrin	1.1095	0.386	1809.86±211.23a	2223.87~1395.85
pH2.5	dichlorvos	5.1519	2.47	481.70±11.62a	504.48~458.92
	fenpropathrin	1.0884	1.6714	1841.06±219.24a	2270.77~1411.35

the development and reproduction of the mite was restrained by direct application of acid rain (pH 4.0-3.0) (Zhao et al. 1999). On the other hand, Wang et al. (2004) reported that development and reproduction of *T. cinnabarinus* were enhanced through the host plant-mediated impact of simulated acid rain (pH 5.6-3.0). The *T. cinnabarinus* population and host plant injury increased significantly after acid rain (pH 4.0) was applied to the mites and the host-plants (Wang and Zhao 2002). Furthermore, Zhang et al. (2004) reported that after long-term exposure to acid rain, physiological aspects of the mite, including some enzymes, were enhanced significantly. Such physiological changes might be related to development, reproduction and acaricide susceptibility of the mites.

The present study clearly demonstrates that immature development and adult reproduction was affected significantly after long exposure to acid rain. The mean generation times at pH 5.6 and pH 4.0 populations were longer, but those at pH 3.0 and pH 2.5 populations were shorter than control. However, doubling times for populations exposed to pH 5.6, pH 4.0 or pH 3.0 acid rain were shorter than controls. In addition, compared with the control population, the relative fitness of mite populations exposed to pH 5.6, 4.0, and 3.0 acid rain were all greater than 1.0. Moreover, Zhang et al. (2005) reported that weak acid rain (pH > 4.0) promoted the activities of three protective enzymes (SOD, POD and CAT) of eggplant leaf and its growth. The growth of *T. cinnabarinus* was also promoted because the content of soluble sugar phosphate and soluble protein changed in eggplant leaves making them more favorable to feeding by mites. However, strong acid rain (pH < 3.0) inhibited the growth of both host plant and the mite. This might be the reason for mite infestations in agricultural areas prone to high rates of acid deposition.

The intrinsic rate of increase (r_m), is often used to determine an insect's ability to thrive on various hosts (Kerns et al. 1989). Comparison of r_m values often provides considerable insight beyond that available from independent analyses of several life history parameters (Petitt et al. 1994). Following Carson (1961), the relative fitness of a population can be evaluated by the average population size when the population has achieved an equilibrium by natural selection after several generations in a given environment. Our previous study showed that the development and reproduction of mites varied

significantly when acid rain was sprayed on mites or host-plants; the jackknife estimate of r_m of pH4.0~3.5 was significantly higher than that of the control population (Zhao et al. 1999; Wang et al. 2004). However, after long-term exposure to acid rain, despite the effects on the development and reproduction of mites, jackknife estimates of r_m and R_f were similar when the simulated acid rain was stopped. This implies that these changes were temporal and able to recover. Acid rain did not induce genetic changes in the mites and only induced life parameter changes that were plant-mediated (Zhang et al. 2005).

Zhang et al. (2004) reported that after long-term exposure to acid rain, the activity of several enzymes closely related to pesticide resistance including peroxidase, superoxide dismutase, catalase, phosphatase and carboxylesterase varied significantly among populations of mites exposed to different acidity treatments. However, bioassay studies reported here indicate that the susceptibility of different *T. cinnabarinus* populations to dichlorvos and fenprothrin did not vary. Thus, although acid rain did influence development and reproduction of *T. cinnabarinus* by direct or indirect effects, acid rain did not affect susceptibility of *T. cinnabarinus* to acaricides.

The present study together with our previous studies give further evidence that *T. cinnabarinus* is remarkably tolerant to a wide range of acidification conditions, and that acid rain, particular pH4.0-5.6, will definitely increase the infestation of the mite. As acid rain significantly affects areas of China, including areas of agricultural and natural biotic importance, further detailed studies investigating the impact of acid rain upon *T. cinnabarinus*-host plant interactions are warranted.

Acknowledgements

This research was funded in part by National Natural Science Foundation (30170617) and Program for New Century Excellent Talents in University, MOE (NCET-04-0854) of P. R. China.

References

- Abbott WJ. 1925. A method of computing the effectiveness of an insecticide. *Journal Economic Entomology* 18: 265-267.

- Bedford BL. 1986. Modification of plant – pest interactions air pollutions. *Proceedings of International Workshop*, Ecosystems Research Center, Cornell University, Ithaca, NY, USA.
- Carson HL. 1961. Heterosis and fitness in experimental populations of *Drosophila melanogaster*. *Evolution* 15: 496-509.
- Cowling EB. 1982. A status report on acid precipitation and its biological consequences as of April 1981. In: D'itri F, editor. *Acid precipitation – effects on ecological systems*, 3-20. Ann Arbor Press, Ann Arbor, MI, USA.
- Dai Z, Liu Y, Wang X, Zhao D. 1998. Changes in pH, CEC, and exchangeable acidity of some forest soils in southern China during the last 32–35 years. *Water, Air, and Soil Pollution* 108: 377-390.
- Dennehy HJ, Granett J, Leigh T. 1983. Relevance of slide-dip and residual bioassay comparisons to detection of resistance in spider mites. *Journal Economic Entomology* 76: 1225-1230.
- Driscoll CT, Lawrence GB, Bulger AJ, Bulter TJ, Cronan CS, Eagar C, Lambert KF, Likens GE, Stoddard JL, Weathers KC. 2001. Acidic deposition in the northeastern United States: sources and inputs, ecosystem effects, and management strategies. *Bioscience* 51: 180-198.
- Grodzki W, McManus M, Knižek M, Meshkova V, Mihalcu V, Novotny J, Turčani M, Slobodyan Y. 2004. Occurrence of spruce beetles in forest stands at different levels of air pollution stress. *Environmental Pollution* 130: 73-83.
- Heliövaara K, Väisänen R. 1993. *Insects and pollution*. Boca Raton: CRC Press Inc.
- Hoagland DR, Arnon DI. 1950. *The water culture method for growing plants without soil*. California Agricultural Experiment Station Circular. 347.
- Hulting FL, Orr DB, Obrycki JJ. 1990. A computer program for calculation and statistical comparison of intrinsic rate of increase and associated life parameters. *Florida Entomologist* 73: 601-612.
- Kerns DL, Puterka GJ, Peters DC. 1989. Intrinsic rate of increase for greenbug (Homoptera: Aphididae) biotypes E, F, G, and H on small grain and sorghum varieties. *Environmental Entomology* 18: 1074-1078.
- Kuylenstierna J, Rodhe H, Cinderby S, Hicks K. 2001. Acidification in developing countries; ecosystem sensitivity and the critical load approach on a global scale. *Ambio* 30: 20-28.
- Lee JJ. 1982. The effects of acid precipitation on crops. In: Acid precipitation – effects on ecological systems. In: D'itri F, editor. *Acid precipitation – effects on ecological systems*, 453–468. Ann Arbor Press, Ann Arbor, MI, USA.
- Li W, Gao J. 2002. Acid deposition and integrated zoning control in China. *Environmental Management* 30: 169-182.
- Linhurst RA, Baker JP, Bartuska AM. 1982. Effects of acidic deposition: A brief review. In: Frederick E, editor. *A specialty conference on atmospheric deposition*. Air Pollution Control Association, Pittsburgh, Pennsylvania, USA.
- Menz FC, Seip HM. 2004. Acid rain in Europe and the United States: an update. *Environmental Science & Policy* 7: 253-265.
- Meyer JS, Ingersoll CG, McDonald LL, Boyce MS. 1986. Estimating uncertainty in population growth rates: Jackknife vs. Bootstrap techniques. *Ecology* 67: 1156-1166.
- Palokangas P, Neuvonen S, Haapala S. 1995. The effects of simulated acid rain on growth and susceptibility to predation of *Phratora polaris* (Col., Chrysomelidae). *Environmental Pollution* 89: 67-71.
- Petitt FL, Loader CA, Schon MK. 1994. Reduction of nitrogen concentration in the hydroponic solution on population growth rate of the aphids (Homoptera: Aphididae) *Aphis gossypii* on cucumber and *Myzus persicae* on pepper. *Environmental Entomology* 23: 930-936.
- Redak RA, Paine TD, Trumble JT. 1995. *Trirhabda geminata* (Coleoptera: Chrysomelidae) resistance to the direct impact of simulated acidic fog on larval growth and mortality. *Environmental Pollution* 90: 61-66.
- Riemer J, Whittaker JB. 1989. Air pollution and insect herbivores: observed interactions and possible mechanisms. In: Bernays EA, editor. *Insect – plant interaction*, 1: 73–106. Boca Raton: CRC Press Inc.
- Shaefer SR. 1992. Responses of microbial pollution in the rhizosphere to deposition of simulated acidic rain into foliage and/or soil. *Environmental Pollution* 76: 267-268.
- Shriner DS. 1986. Terrestrial ecosystems: wet deposition. In: Legge AH, Krupa SV, editors. *Air pollutants and their effects on the terrestrial ecosystem*. *Advances in environmental Sciences and Technology*, 18: 365–388. New York: John Wiley & Sons.
- Sokal RR, Rohlf FJ. 1969. *Biometry – the principles and practice of statistics of statistics in biological research*. San Francisco: Freeman.
- Stark JD, Banks JE. 2003. Population-level effects of pesticides and other toxicants on arthropods. *Annual Review of Entomology* 48: 505-19.
- Stinner DH, Stinner BR, McCartney DA. 1988. Effects of simulated acidic precipitation on plant–insect interactions in agricultural systems: corn and black cutworm larvae. *Journal of Environmental Quality* 17: 371-376.
- Treshow M, Anderson FK. 1989. *Plant stress from air pollution*. New York: John Wiley & Sons.
- Treshow M. 1984. *Air pollution and plant life*. New York: John Wiley & Sons.

- Walton WE, Camptom SM, Allan JD, Daniels JD. 1982. The effect of acid stress on survivorship and reproduction of *Daphnia pulex* (Crustacea: Cladocera). *Canadian Journal of Zoology* 60: 573-579.
- Wang JJ, Zhao ZM, Zhang JP. 2004. The host plant-mediated impact of simulated acid rain on the development and reproduction of *Tetranychus cinnabarinus* (Acari: Tetranychidae). *Journal of Applied Entomology* 128: 397-402.
- Wang JJ, Zhao ZM. 2002. Study on effect of simulated acid rain on carmine spider mite, *Tetranychus cinnabarinus*. In: Li DM, Kang L, Wu JW, Zhang RZ, editors. *Innovation and Development in Entomology*, 693–696. Beijing: Scientific and Technological Press.
- Wang LJ, Qin CL. 1989. Study on the characteristics of the growth of vegetables on the purple soil soaked by simulated acid rain. *Chongqing Environmental Science* 11: 11-15.
- Zhang JP, Wang JJ, Zhao ZM, Dou W, Chen Y. 2004. Effects of simulated acid rain on the physiology of carmine spider mite, *Tetranychus cinnabarinus* (Boisduvals) (Acari: Tetranychidae). *Journal of Applied Entomology* 128: 342-347.
- Zhang JP, Wang JJ, Zhao ZM, Chen Y, Dou W. 2005. Effects of simulated acid rain on physiological and biochemical characters of eggplant, the host plant of *Tetranychus cinnabarinus*. *Chinese Journal of Applied Ecology* 16: 450-454.
- Zhao ZM, Luo HY, Wang JJ. 1999. Effects and Mechanisms of simulated acid rain on plant-mite interactions in agricultural systems. I. The direct effects of simulated acid rain on carmine spider mite, *Tetranychus cinnabarinus*. *Systematic & Applied Acarology* 4: 83-89.