



Eromangateuthis N. Gen., a New Genus for a Late Albian Gladius-Bearing Giant Octobranchian (Cephalopoda: Coleoidea)

Author: Dirk Fuchs

Source: Paleontological Contributions, 2019(21) : 1-3

Published By: The Paleontological Institute at The University of Kansas


URL: <https://doi.org/10.17161/1808.29619>

BioOne Complete (complete.BioOne.org) is a full-text database of 200 subscribed and open-access titles in the biological, ecological, and environmental sciences published by nonprofit societies, associations, museums, institutions, and presses.

Your use of this PDF, the BioOne Complete website, and all posted and associated content indicates your acceptance of BioOne's Terms of Use, available at www.bioone.org/terms-o-use.

Usage of BioOne Complete content is strictly limited to personal, educational, and non - commercial use. Commercial inquiries or rights and permissions requests should be directed to the individual publisher as copyright holder.

BioOne sees sustainable scholarly publishing as an inherently collaborative enterprise connecting authors, nonprofit publishers, academic institutions, research libraries, and research funders in the common goal of maximizing access to critical research.



Paleontological Contributions

October 8, 2019

Number 21

EROMANGATEUTHIS N. GEN., A NEW GENUS FOR A LATE ALBIAN GLADIUS-BEARING GIANT OCTOBRACHIAN (CEPHALOPODA: COLEOIDEA)

Dirk Fuchs

SNSB-Bayerische Staatssammlung für Paläontologie und Geologie, Richard-Wagner-Str. 10, 80333 Munich, Germany; drig.fuchs@gmail.com

ABSTRACT

A comparative analysis shows that the gladius morphology of *Boreopeltis helgolandiae* Engeser & Reitner, 1985 (Aptian), the type species of *Boreopeltis* Engeser & Reitner, 1985, *Boreopeltis sagittata* (Naef, 1921) (Tithonian), and *Boreopeltis smithi* Fuchs & Larson, 2011 (Cenomanian) is fundamentally different from *Boreopeltis soniae* Wade, 1993 (Aptian). I therefore propose *Eromangateuthis* n. gen. in order to accommodate *Eromangateuthis soniae* (Wade, 1993).

Key words: Octobrachia, taxonomy, gladius, lower Cretaceous, Australia.

INTRODUCTION

Mary Wade (1993, fig. 5–8) introduced the new taxon *Boreopeltis soniae* from the middle upper Albian of Queensland (Australia). The species is based on several squid-like gladii, one of which has a length of about 1.2 m, hence, a mantle length twice as long as that of extant *Enteroctopus dofleini* (Wülker, 1910), the Giant Pacific octopus. Wade (1993) placed the new species into the genus *Boreopeltis* Engeser & Reitner, 1985 owing to supposed similarities. I investigated the type material of *Boreopeltis soniae* in 2005 and immediately recognized striking differences between *Boreopeltis soniae* and the type species, lower Aptian *Boreopeltis helgolandiae* Engeser & Reitner, 1985 and all other known species assigned to the genus *Boreopeltis*, i.e., lower Tithonian *Boreopeltis sagittata* (Naef, 1922) and Cenomanian *Boreopeltis smithi* Fuchs & Larson, 2011. Along with the presentation of the first Late Cretaceous member, *Boreopeltis smithi*, I have already mentioned doubts and excluded *Boreopeltis soniae* from the genus *Boreopeltis* (Fuchs & Larson, 2011, p. 242). The erection of a new genus is a logical consequence and is done herein.

MATERIAL & METHODS

The type specimens of *Boreopeltis soniae* are housed in the Queensland Museum in Brisbane. During my stay in Australia (2005), I investigated further specimens in the collection of the Kronosaurus Corner, a public museum in Richmond (northwest

Queensland). Terminology and gladius parameters used below follow the *Treatise Online*, Part M, Coleoidea, chapter 9B: The gladius and gladius vestige in fossil Coleoidea (Fuchs, 2016). The diagnosis below matches the diagnoses and the system presented in the *Treatise Online*, Part M, Coleoidea, chapter 23G: Octobrachia (Fuchs, submitted 2019).

SYSTEMATIC PALEONTOLOGY

Superorder Octobrachia Haeckel, 1866

Suborder Prototeuthina Naef, 1921

Family Plesiotheuthidae Naef, 1921

Eromangateuthis nov. gen.

Type species.—*Boreopeltis soniae* Wade, 1993.

Etymology.—After the so-called Eromanga basin in northeast Australia.

Diagnosis (new).—Large-sized plesiotheuthids; gladius slender to moderately wide (gladius width_{max} to gladius length 0.15–0.25) with pronounced solid median keel, which tapers anteriorly; median field very slender to slender (opening angle 10–15°, median field width_{hypz} to hyperbolar zone length 0.15–0.25); anterior margin distinctly convex, with lateral plate- or channel-like reinforcements; median field area very large (median field area to gladius area 0.85–0.95); lateral fields very slender to slender (lateral fields width_{max} to median field width_{max} 0.45–0.55); hyperbolar zones very short to short, difficult to determine (hyperbolar zone length to




Fig. 1: Comparative gladius morphology of the *Boreopeltis* species complex and *Eromangateuthis soniae*. A, *Boreopeltis helgolandiae* (Aptian; gladius line drawing modified after Engeser & Reitner, 1985); B, *Boreopeltis sagittata* (Tithonian); B1, gladius reconstruction in dorsal view; B2, photograph of specimen BSPG 2011 XLI 332-08; B3, gladius reconstruction in ventrolateral view; C, *Boreopeltis smithi* (Cenomanian); C1, gladius reconstruction in dorsal view; C2, gladius reconstruction in ventrolateral view; D, *Eromangateuthis soniae* n. comb. (Albian); D1, gladius reconstruction in dorsal view; D2, gladius reconstruction in ventrolateral view; D3, photograph of a gladius in the exhibition of the Kronosaurus Korner Museum, Richmond (Queensland, Australia), dorsal view; black dots show position of conus apex; scale bars 10 cm.

median field length 0.15–0.25); conus unusually deep, ventrally oriented (not funnel-like), patella shaped; soft parts unknown.

Included species.—Only the type species.

Stratigraphical and geographical range.—Lower Cretaceous (upper Albian) of Queensland (East Australia).

Differential diagnosis (Fig. 1).—Although Wade (1993) described the gladius morphology of *Boreopeltis soniae* in detail, comparisons with other prototeuthid octobranchians have been neglected. Actu-

ally, *Eromangateuthis soniae* and the *Boreopeltis* species complex have little in common; neither the anterior nor the posterior gladius part show a vague resemblance. A pronounced median keel present in *Eromangateuthis* n. gen. is absent in *Boreopeltis*, where the median line is not solid and only slightly elevated. Conversely, solid lateral reinforcements present on the median field of *Boreopeltis* are absent in *Eromangateuthis* n. gen. The lateral reinforcements in *Boreopeltis* are anteriorly projected; the shape

of the anterior median field margin is therefore fundamentally different from *Eromangateuthis* n. gen., whose anterior margin is evenly rounded. The relative median field width of *Eromangateuthis* n. gen. is close to *Boreopeltis*, but the latter genus is missing the very narrow waist, which is caused by short and rapidly tapering lateral fields and hyperbolar zones. In *Boreopeltis*, the hyperbolar zone is significantly longer (hyperbolar zone length-to-median field length 0.50–0.70). Finally, the conus of *Eromangateuthis* n. gen. is distinct from a funnel-like conus typical for *Boreopeltis* and other plesiotheuthids. The conus of *Eromangateuthis* n. gen. is unusual in being patella-like, similar to muensterelloid teudopseids (Fuchs & others, 2019).

Eromangateuthis soniae (Wade, 1993) n. comb.

Figure 1D.1–1D.3

Type material.—Holotype (QMF25285), seven paratypes (QMF25775, 15209, 21988, 25284, 25286, 25288, 25289).

Type locality.—Marathon near Richmond (Queensland, Australia); southern anabranch of Flinders River (200 m upstream L813).

Type horizon.—Allaru Formation, middle upper Albian (see Wade, 1993).

Studied material.—Apart from the type material, I have studied specimens QMF25990, 25294, 52687, 39836; a specimen in the collection of the Kronosaurus Korner Museum in Richmond (Queensland, Australia), and a fragmentary conus deposited in Natural History Museum of London (NHMUK C.59211).

Description.—*Eromangateuthis soniae* is a plesiotheuthid with a gladius of moderate width (gladius width_{max} to gladius length 0.21). The slender median field (opening angle 10–11°, median field width_{hypz} to hyperbolar zone length 0.19) exhibits a pronounced solid median keel, which tapers anteriorly. The anterior margin of the median field is distinctly convex. The lateral margins are thickened by longitudinal plate- or channel-like reinforcements. The median field is very large (median field area to gladius area

0.91) compared to the lateral fields, which are slender (lateral fields width_{max} to median field width_{max} 0.52) and short (hyperbolar zone length to median field length 0.22). The posterior conus is unusually deep, ventrally oriented (not funnel-like), and therefore patella shaped. Soft parts are unknown. For a detailed description of the type material of *Eromangateuthis soniae*, see Wade (1993).

REFERENCES

- Engeser, Theo, & Joachim Reitner. 1985. Teuthiden aus dem Unterapt ("Töck") von Helgoland (Schleswig-Holstein, Norddeutschland). *Paläontologische Zeitschrift* 59:245–260.
- Fuchs, Dirk. submitted 2019. Part M, Chapter 23G: Systematic Descriptions: Octobranchia. *Treatise Online*.
- Fuchs, Dirk. 2016. Part M, chapter 9B: The gladius and gladius vestige in fossil Coleoidea. *Treatise Online* 83:1–23.
- Fuchs, Dirk, Yasuhiro Iba, Alexander M. Heyng, Masaya Iijima, Christian Klug, N. L. Larson, & Günter. Schweigert. 2019. The Muensterelloidea: Phylogeny and character evolution of Mesozoic stem-octopods. *Special papers in Palaeontology*. DOI: 10.1002/spp2.1254.
- Fuchs, Dirk, & N. L. Larson. 2011. Diversity, morphology, and phylogeny of coleoid cephalopods from the Upper Cretaceous Plattenkalks of Lebanon. Part I: Prototeuthidina. *Journal of Paleontology* 85(2):234–249.
- Haeckel, E. H. P. A. 1866. *Generelle Morphologie der Organismen*. Georg Reimer, Berlin. 462 p.
- Naef, Adolf. 1921. *Fauna und Flora des Golfes von Neapel und der angrenzenden Meeres-Abschnitte*. 35. Monographie: Die Cephalopoden, 1 Teil, 1 Lieferung. R. Friedländer und Sohn. Berlin. 148 p.
- Naef, Adolf. 1922. *Die fossilen Tintenfische: Eine paläozoologische Monographie*. Gustav Fischer, Jena. 322 p.
- Wade, Mary. 1993. New Kelaenida and Vampyromorpha: Cretaceous squid from Queensland. *Memoirs of the Association of Australasian Paleontologists* 15:353–374.
- Wülker, Gerhard. 1910. Über Japanische Cephalopoden. Beiträge zur Kenntnis der Systematik und Anatomie der Dibranchiaten. *Abhandlungen der Mathematisch-Physikalischen Classe der Königlich Bayerischen Akademie der Wissenschaften* 3(supplement 1):1–77.