

DECOTAB: a multipurpose standard substrate to assess effects of litter quality on microbial decomposition and invertebrate consumption

Authors: Andries A. Kampfraath, Ellard R. Hunting, Christian Mulder, Anton M. Breure, Mark O. Gessner, et. al.

Source: Freshwater Science, 31(4) : 1156-1162

Published By: Society for Freshwater Science

URL: <https://doi.org/10.1899/12-075.1>

BioOne Complete (complete.BioOne.org) is a full-text database of 200 subscribed and open-access titles in the biological, ecological, and environmental sciences published by nonprofit societies, associations, museums, institutions, and presses.

Your use of this PDF, the BioOne Complete website, and all posted and associated content indicates your acceptance of BioOne's Terms of Use, available at www.bioone.org/terms-of-use.

Usage of BioOne Complete content is strictly limited to personal, educational, and non-commercial use. Commercial inquiries or rights and permissions requests should be directed to the individual publisher as copyright holder.

BioOne sees sustainable scholarly publishing as an inherently collaborative enterprise connecting authors, nonprofit publishers, academic institutions, research libraries, and research funders in the common goal of maximizing access to critical research.

DECOTAB: a multipurpose standard substrate to assess effects of litter quality on microbial decomposition and invertebrate consumption

Andries A. Kampfraath^{1,6,13}, Ellard R. Hunting^{1,7,13}, Christian Mulder^{2,8},
Anton M. Breure^{2,3,9}, Mark O. Gessner^{4,5,10}, Michiel H. S. Kraak^{1,11}, AND
Wim Admiraal^{1,12}

¹ Aquatic Ecology and Ecotoxicology, Institute for Biodiversity and Ecosystem Dynamics (IBED-AEE),
University of Amsterdam, Sciencepark 904, NL-1098 XH, Amsterdam, The Netherlands

² National Institute for Public Health and the Environment (RIVM), Antonie van Leeuwenhoeklaan 9,
NL-3720 BA, Bilthoven, The Netherlands

³ Department of Environmental Science, Institute for Water and Wetland Research (IWWR),
Radboud University, Heyendaalseweg 135, NL-6525 AJ, Nijmegen, The Netherlands

⁴ Leibniz Institute of Freshwater Ecology and Inland Fisheries (IGB), Alte Fischerhütte 2, 16775 Stechlin, Germany

⁵ Department of Ecology, Berlin Institute of Technology (TU Berlin), Ernst-Reuter-Platz 1, 10587 Berlin, Germany

Abstract. Currently available tools for studying plant litter decomposition and invertebrate consumption in aquatic ecosystems have at least 2 major limitations: 1) the difficulty of manipulating litter chemical composition to provide mechanistic insights into attributes of litter quality controlling decomposition rate, and 2) lack of a standardized litter that hampers comparisons of results among studies. These limitations point to a need for a standard litter surrogate with adjustable chemical composition. We propose using a decomposition and consumption tablet (DECOTAB) consisting of cellulose powder embedded in an agar matrix to evaluate decomposition and consumption rates in aquatic environments. We describe the preparation of DECOTABs and demonstrate some applications in laboratory microcosms and outdoor mesocosms. A leaf shredder, the isopod *Asellus aquaticus*, and a collector-gatherer, the nonbiting midge larva *Chironomus riparius*, readily consumed DECOTABs, leading to massive mass loss of the tablets within 21 d (~90%). The isopod also consumed DECOTABs amended with extracts of riparian plants and soil to create a chemically complex source of organic matter. Our results highlight the potential utility of DECOTABs to assess invertebrate contributions to organic matter decomposition in aquatic systems. In the absence of invertebrates, exposure of basic and complex DECOTABs to microorganisms resulted in significant mass loss within 21 d (10–25%), and addition of an antibiotic and fungicide suppressed microbial decomposition, suggesting that the tablets are useful for studying microbial processes. Complex tablets decomposed faster than the basic tablets, a result illustrating the importance of chemical composition of organic material for microbial decomposers. DECOTABs are a novel, versatile tool for addressing long-standing questions in aquatic ecology and environmental assessment.

Key words: litter decomposition, cellulose degradation, microorganisms, detritivores, benthos, functional ecosystem assessment, aquatic ecosystem health, methodology, standardization.

⁶ E-mail addresses: d.kampfraath@gmail.com

⁷ e.r.hunting@uva.nl

⁸ christian.mulder@rivm.nl

⁹ ton.breure@rivm.nl

¹⁰ gessner@igb-berlin.de

¹¹ m.h.s.kraak@uva.nl

¹² w.admiraal@uva.nl

¹³ Both authors contributed equally

Decomposition of plant litter is a vital ecosystem process driven by both microorganisms and detritivores (Webster and Benfield 1986, Graça 2001, Gessner et al. 2010, Tank et al. 2010). The standard approach to studying decomposition in the field is to measure leaf mass loss from litter bags made of coarse or fine mesh that controls access of invertebrates differing in body size (Boulton and Boon 1991). Although not perfect, partly because of risks of hypoxia in fine mesh bags or litter fragmentation by

turbulent flow in coarse mesh bags (Boulton and Boon 1991), this approach adequately mimics natural conditions in many situations and represents the influences of both litter quality and environmental factors on decomposition dynamics. Litter bags are the method of choice in comparative field studies (comparisons of litter types, locations, etc.) and often yield realistic estimates of local decomposition rates (Webster and Benfield 1986, Boulton and Boon 1991). Laboratory studies in microbial microcosms (Suberkropp 1991, Dang et al. 2009) and feeding trials with detritivores (Cameron and La Point 1978, González and Graça 2003) complement the litterbag approach by providing important insight into the decomposition process. However, rigorous tests of the mechanisms responsible for differences in decomposition are difficult with approaches relying on natural litter that varies in many quality attributes.

Standardized ways to measure decomposition rates that exclude confounding effects of varying litter quality are required to allow comparisons on large geographical and temporal scales, e.g., for studies on wide-ranging impacts of human activities (pollution, habitat modification, etc.) on ecosystem functioning in natural environments (Gessner and Chauvet 2002, Young and Collier 2009, Woodward et al. 2012). Selecting individual plant litter species for large-scale studies (Boyer et al. 2011) is only a partial remedy because litter quality can vary widely even within a given plant species (Lecerf and Chauvet 2008). Filter paper and cotton strips have long been used as substitutes for natural plant litter (Egglisshaw 1964, Hildrew et al. 1984, Tiegs et al. 2007, Imberger et al. 2010). Cotton strips, in particular, are used as a standardized substrate for this purpose (e.g., Boulton and Quinn 2000, Young et al. 2008, Young and Collier 2009) because they are essentially composed of cellulose, a major component of plant litter, and have much greater tensile strength than filter paper.

Despite their potential utility in decomposition studies, cotton strips have limitations (Imberger et al. 2010). First, the previously used standard material (Shirley Soil Burial Test Fabric) is no longer available, so comparisons between present-day studies and earlier studies are complicated, although results with Shirley Soil Burial Fabric and replacement materials, such as 'calico' are correlated (Imberger et al. 2010). However, calicoes are produced from natural cotton and, therefore, have an uncertain and variable chemical composition. Second, the chemical composition of cotton strips is extremely simple (cellulose content > 95%) compared to natural plant litter, such that numerous leaf constituents (N, lignin, tannins, fatty acids, etc.) that are potentially important factors

in decomposer activity and decomposition rate are lacking. Third, the composition of cotton strips is difficult to manipulate experimentally to test for effects of specific chemical plant constituents or of compounds inhibiting microbial or detritivore activity (e.g., antibiotics, fungicides, insecticides; see Rader et al. 1994). These limitations highlight the need to identify a standardized substrate whose composition can be altered according to the needs defined by the question posed.

The objective of our study was to develop and test a standardized proxy material whose chemical composition could be adjusted easily for experimental purposes. Inclusion of specific plant constituents and other compounds at desired concentrations in polycarbohydrate gels (e.g., agar or phytagel) has been useful in studies of allelochemical interactions (Hay et al. 1998), colonization dynamics of sessile invertebrates (Henrikson and Pawlik 1995, Hunting et al. 2010b), and feeding preferences of aquatic herbivores (Pavia and Toth 2000). We assessed whether this matrix could be modified to serve as surrogate plant material whose chemical composition can be controlled and whose texture broadly resembles that of natural litter. However, agar has a low nutritional value for invertebrates and can be degraded by only a limited number of microorganisms (Bärlocher and Porter 1986, Armisen 1991). Therefore, we developed and tested the performance of a decomposition and consumption tablet (DECOTAB) consisting of a high concentration of cellulose powder (75%) embedded in an agar matrix to evaluate the importance of factors affecting microbial decomposition and invertebrate feeding. We describe the preparation of the proposed DECOTABs and evaluate some potential applications in laboratory microcosms and outdoor mesocosms.

Methods

Preparation of DECOTABs

We made basic DECOTABs from a suspension containing 60 g of powdered cellulose (Sigma-Aldrich, St. Louis, Missouri), 20 g of purified agar (Oxoid Ltd., Basingstoke Hampshire, UK), and 60 μmol ascorbic acid/L deionized water (dH_2O) as an antioxidant (Merck GmbH, Darmstadt, Germany) (Niki 1991). We heated the mixture to 100°C to dissolve the agar, allowed it to cool to 50°C with frequent stirring, and poured it into a multiwell polycarbonate mold (15-mm diameter, 5-mm height) to cast tablets with a final volume of 118 mm^3 (Fig. 1A). The tablets initially had a convex surface that quickly flattened during solidification of the agar at 7°C . The tablets could be stored in closed containers




FIG. 1. Polycarbonate mold used to prepare DECOTABs (A), and isopods (*Asellus aquaticus*) rallying around and feeding on a basic cellulose DECOTAB (B).

for up to 3 wk in the refrigerator without noticeable decay or dehydration. We measured the dry mass of freshly prepared tablets (mean \pm 1 SD = 81.3 ± 3.1 mg, $n = 25$) by drying them for 3 d at 70°C and then weighing them with an analytical balance (precision = 0.1 mg; Mettler AT261, Mettler-Toledo, Tiel, The Netherlands).

We prepared DECOTABs including an antibiotic and fungicide as described above except that we added chloramphenicol (60 mg/L dH_2O ; Sigma-Aldrich) and cycloheximide (60 mg/L dH_2O ; Sigma-Aldrich) to the cooled suspension at 50°C . We prepared DECOTABs of more complex composition to enhance their resemblance to natural particulate organic matter (POM). These tablets had the same volume as the basic DECOTABs, but consisted of 46.7 g of cellulose, 20 g of purified agar, and 60 μmol of ascorbic acid/L dH_2O . We extracted plant constituents from bulk standardized garden soil (Baseline, Maxeda DIY, Diemen, The Netherlands), stinging nettle (*Urtica dioica*), and willow leaf extract (*Salix alba*), both common plants in the riparian zone of temperate rivers in Europe. We used 70% acetone as the extraction solvent and air-dried the extract (Hunting et al. 2010a). We added powdered extracts (6.7 g/L for soil; 3.3 g/L for stinging nettle, and 3.3 g/L for willow leaves) to the cooled suspension at 50°C .

Experiment 1: Microbial vs invertebrate-mediated decomposition

We ran an experiment in laboratory microcosms to evaluate the relative contribution of microorganisms and detritivores to decomposition of DECOTABs. We

used 100-mL glass microcosms with sediment containing 50 g quartz sand (0.1–0.5-mm grain size; Dorsilit, Eurogrit, Papendrecht, The Netherlands) and a mixture of standard culture food (7.5 mg) composed of Trouvit[®] (Trouw, Fontaine-les-Vervins, France) and Tetraphyl[®] (Tetrawerke, Melle, Germany) at a ratio 20:1 as organic material. The overlying water consisted of 250 mL Dutch Standard Water (Maas et al. 2002). To obtain a natural microbial inoculum, we added 50 mL of filtered ($<75 \mu\text{m}$) natural surface water from a local shallow lake that typically supports a diverse microbial community (del Giorgio and Gasol 1995). A detailed description of Dutch freshwater bacteria in comparable lakes was published by Zwart et al. (1998, 2002). We aerated the water gently with compressed air throughout the experiment.

We tested 2 invertebrate species: the nonbiting midge larva *Chironomus riparius* (laboratory culture) and the isopod *Asellus aquaticus* (collected from a nearby shallow lake). We used relationships among invertebrate length, fresh mass, and dry mass to standardize invertebrate biomass (Hunting et al. 2012). We added invertebrates based on equal initial dry mass. Thus, we added either 5 *A. aquaticus* (7–9 mm) or 20 *C. riparius* larvae (2nd instar) to each microcosm and placed 1 DECOTAB in each microcosm. Treatments were: 1) basic DECOTAB, no invertebrates; 2) DECOTAB containing an antibiotic and fungicide, no invertebrates; 3) basic DECOTAB and *C. riparius*; and 4) basic DECOTAB and *A. aquaticus*. We did not include a treatment with DECOTAB containing both antibiotics and invertebrates because of the toxicity of both antibiotics to invertebrates (Baliga et al. 1970, Monari et al. 2008). We replicated each treatment 10 times. After 21 d, we removed the DECOTABs with a needle, rinsed and dried (3 d at 70°C) them, and weighed them with an analytical balance (precision = 0.1 mg; Mettler AT261). We subtracted the final dry mass from the estimated initial dry mass to calculate DECOTAB mass loss. We tested for treatment differences with a 1-way nonparametric permutation-based multivariate analysis of variance (PERMANOVA) based on Bray-Curtis distances and 9999 permutations (Anderson 2001), followed by a Bonferroni-corrected PERMANOVA pairwise comparisons in PAST (Hammer et al. 2001).

Experiment 2: Effects of DECOTAB complexity on decomposition and consumption

We ran a 21-d mesocosm experiment in June 2011 to evaluate the performance of DECOTABs under outdoor conditions and to compare basic and complex




FIG. 2. Mean (± 1 SD) mass loss of cellulose DECOTABs in laboratory microcosms inoculated with microorganisms only or additionally stocked with 1 of 2 invertebrate species (*Asellus aquaticus* or *Chironomus riparius*) ($t = 21$ d). Bars with different letters are significantly different (Bonferroni-corrected permutation-based multivariate analysis of variance [PERMANOVA] pairwise comparisons).

DECOTABs containing organic matter extracted from plants and soil. Mesocosms consisted of rectangular (66 cm long \times 34 cm wide \times 30 cm high) 90-L plastic tubs. They contained ~ 40 L of rainwater and 18.5 L of sediment made of standardized garden soil (Baseline) and quartz sand (0.1–0.5 mm; Dorsilit, Eurogrit, Papendrecht, The Netherlands) mixed in a ratio of 5 L soil/25 kg sand. We placed the mesocosms in concrete containers filled with water to buffer temperature fluctuations. We pulled a gauze screen (mesh size = 1 mm) over the concrete containers to reduce colonization by allochthonous fauna. Before starting the experiment, we allowed the mesocosms to sit for 2 d to allow the sediment to settle. We assumed that microbial communities could sufficiently acclimate during these 2 d. Subsequently, each mesocosm received 3 basic and 3 complex DECOTABs and either no invertebrates or *A. aquaticus*. We added the isopods at densities of 402 individuals (ind.)/m² (90 ind./mesocosm), which falls within the density range reported for natural populations (67–586 ind./m²; Adcock 1979). We replicated both treatments 5 times. During the experiment, we gently aerated the overlaying water with a permanently installed air compressor aeration system. After 21 d, we removed the DECOTABs and weighed them as described above. We tested for treatment differences with a




FIG. 3. Mean (± 1 SD) mass loss of basic cellulose DECOTABs and complex DECOTABs containing plant and soil extracts mediated by *Asellus aquaticus* or microorganisms in outdoor mesocosms ($t = 21$ d). Bars with different letters are significantly different (Bonferroni-corrected permutation-based multivariate analysis of variance [PERMANOVA] pairwise comparisons).

2-way factorial PERMANOVA based on Bray–Curtis distances and 9999 permutations (Anderson 2001), followed by Bonferroni-corrected PERMANOVA pairwise comparisons in PAST.

Results

DECOTAB mass loss differed between treatments in the microcosm experiment ($F = 23.61$, $p < 0.0001$; Fig. 2). Both *C. riparius* and *A. aquaticus* fed actively on the basic DECOTABs (Fig. 1B), resulting in 80 to 90% mass loss over the course of the 21-d experiment. In microcosms without invertebrates, mass loss averaged 10 to 25% of the initial mass, and no mass loss occurred when DECOTABs contained an antibiotic and fungicide (Fig. 2).

Mass loss in outdoor mesocosms was higher in the presence of invertebrates than in the microorganism-only treatment ($F = 317.4$, $p = 0.0001$; Fig. 3). Mass loss of complex DECOTABs containing soil and plant extracts was higher than that of basic DECOTABs when only microorganisms were present ($F = 13.0$, $p = 0.002$; pairwise comparison, $p = 0.031$), but this difference was not apparent when isopods were allowed to feed on DECOTABs (pairwise comparison, $p = 0.43$). Mass loss in the presence of invertebrates was

similar in outdoor mesocosms and laboratory microcosms in the presence of *A. aquaticus* (cf. Figs 2, 3).

Discussion

The new standardized plant litter substitute developed and tested in our study was useful for measuring rates of microbial decomposition and consumption by invertebrates in sediments. Both invertebrate test species fed actively on the DECOTABs, resulting in a dramatic mass loss, despite the presence of other sources of organic matter in the sediments of our micro- and mesocosms. Moreover, DECOTABs exposed to microorganisms in the absence of invertebrates also lost mass in the 2 experiments, whereas mass loss was suppressed in microcosms with DECOTABs containing an antibiotic and a fungicide. These results suggest that DECOTABs are a useful substrate for studying both microbial decomposition and consumption of organic matter by invertebrates in aquatic systems. However, the extent to which they reflect decomposition of natural plant litter and could be used to develop functional metrics for assessing impacts of anthropogenic stressors on aquatic ecosystems remains to be tested.

DECOTABs were consumed by a shredder (*A. aquaticus*) and by *C. riparius*, which burrows in sediments and feeds on fine particulate organic matter (Cummins and Klug 1979). Chironomids readily consumed DECOTABs, a result suggesting that DECOTABs could be used to study mechanisms affecting use of organic matter by collector-gatherers (in addition to shredders) and to measure their role in organic-matter dynamics of aquatic ecosystems.

Experimental manipulation of substrate composition facilitates mechanistic insights into decomposition processes. One could test whether and which lipids are triggers for detritivory, growth, and reproduction (Cargill et al. 1985). One could assess whether particular phenolics or mixtures of phenolics affect microbial activity or detritivore performance in laboratory or field conditions. We supplemented cellulose DECOTABs with an antibiotic and fungicide to assess whether and to what extent microorganisms vs detritivores contribute to decomposition.

Investigators have used various approaches to assess the contribution of shredders to litter decomposition (Petersen and Cummins 1974, Cuffney et al. 1990, Hieber et al. 2002, González and Graça 2003), but the available information is still scarce (Boulton and Quinn 2000, Young et al. 2008). We did not compare DECOTABs with natural plant litter, so our results are not directly comparable with estimates

based on measured litter consumption (Hieber and Gessner 2002, Hunting et al. 2012). Nevertheless, our data corroborate the notion that invertebrates can play dominant roles in organic-matter turnover in aquatic ecosystems (Wallace and Webster 1996). Invertebrates contributed up to 55 to 65% of total DECOTAB mass loss, whereas microorganisms contributed only 10 to 25%.

Plant litter is a complex mixture of structurally diverse compounds. To mimic the composition of natural litter in mesocosms, we created DECOTABs containing extracts of riparian plants and soil and offered them to microorganisms only or to microorganisms in combination with *A. aquaticus*. The differential response of microbes to basic cellulose and complex DECOTABs illustrates the importance of resource composition for microbial decomposers. For invertebrates, a similar distinction was not observed, possibly because resemblance in texture masked differences in chemical composition. However, conclusive answers about the relative importance of texture vs chemistry for DECOTAB palatability require tests with tablets differing more widely in chemical composition.

Standardized measures to determine decomposition rates are increasingly important for assessing effects of anthropogenic stressors on ecosystem processes (Gessner and Chauvet 2002). However, methods differ among studies, and some are based on using litter from different plant species that decompose at different rates. Even litter of the same species may vary in chemical composition and texture if collected at different locations (Lecerf and Chauvet 2008) or at different times. This variability hampers comparisons of decomposition dynamics over large spatial and temporal scales (Boyero et al. 2011). Use of a standardized substrate would facilitate comparisons among studies, including studies aimed at assessing the response of ecosystem functioning to anthropogenic stressors. The DECOTAB approach could facilitate such standardized experiments at larger scales and increase power of meta-analyses.

In conclusion, our study demonstrated the potential of the newly developed DECOTABs for a variety of applications in aquatic environments. The DECOTABs described here were based on an aqueous matrix, so they shrink upon dehydration. Therefore, they cannot be used readily in terrestrial systems because dehydration will affect shape and texture. However, the basic concept could be applicable to terrestrial studies if tablets were prepared dry. One of the greatest assets of DECOTABs is that they can be prepared in almost any desired size, shape, or composition to suit the needs of the specific question

to be examined. Thus, they have the potential to become a highly standardized and versatile tool to address long-standing issues in aquatic ecology and environmental assessment.

Acknowledgements

We thank Manuel Graça and 2 anonymous referees for their constructive comments on earlier drafts of this article. In addition, we express our gratitude to Jan van Arkel for providing the picture of the DECOTAB mold.

Literature Cited

- ADCOCK, J. A. 1979. Energetics of a population of the isopod *Asellus aquaticus*: life history and production. *Freshwater Biology* 9:343–355.
- ANDERSON, M. J. 2001. A new method for non-parametric multivariate analysis of variance. *Austral Ecology* 26: 32–46.
- ARMISÉN, R. 1991. Agar and agarose biotechnological applications. *Hydrobiologia* 221:157–166.
- BALIGA, B. S., S. A. COHEN, AND H. N. MUNRO. 1970. Effect of cycloheximide on the reaction of puromycin with poly-some-bound peptidyl-tRNA. *FEBS Letters* 8:249–252.
- BÄRLOCHER, F., AND C. W. PORTER. 1986. Digestive enzymes and feeding strategies of three stream invertebrates. *Journal of the North American Benthological Society* 5:58–66.
- BOULTON, A., AND P. BOON. 1991. A review of methodology used to measure leaf litter decomposition in lotic environments: time to turn over an old leaf? *Australian Journal of Marine and Freshwater Research* 42:1–43.
- BOULTON, A. J., AND J. M. QUINN. 2000. A simple and versatile technique for assessing cellulose decomposition potential in floodplain and riverine sediments. *Archiv für Hydrobiologie* 150:133–151.
- BOYERO, L., R. G. PEARSON, M. O. GESSNER, L. A. BARMUTA, V. FERREIRA, M. A. S. GRAÇA, D. DUDGEON, A. J. BOULTON, M. CALLISTO, E. CHAUVET, J. E. HELSON, A. BRUDER, R. J. ALBARIÑO, C. M. YULE, M. ARUNACHALAM, J. N. DAVIES, R. FIGUEROA, A. S. FLECKER, A. RAMÍREZ, R. G. DEATH, T. IWATA, J. M. MATHOOKO, C. MATHURIAU, J. F. GONÇALVES, JR., M. S. MORETTI, T. JINGGUT, S. LAMOTHE, C. M'ÉRIMBA, L. RATNARAJAH, M. H. SCHINDLER, J. CASTELA, L. M. BURIA, A. CORNEJO, V. D. VILLANUEVA, AND D. C. WEST. 2011. A global experiment suggests climate warming will not accelerate litter decomposition in streams but might reduce carbon sequestration. *Ecology Letters* 14: 289–294.
- CAMERON, G. N., AND T. W. LA POINT. 1978. Effects of tannins on the decomposition of Chinese tallow leaves by terrestrial and aquatic invertebrates. *Oecologia* (Berlin) 32:349–366.
- CARGILL, A. S., K. W. CUMMINS, B. J. HANSON, AND R. R. LOWRY. 1985. The role of lipids as feeding stimulants for shredding aquatic insects. *Freshwater Biology* 15: 455–464.
- CUFFNEY, T. F., J. B. WALLACE, AND G. J. LUGTHART. 1990. Experimental evidence quantifying the role of benthic invertebrates in organic-matter dynamics of headwater streams. *Freshwater Biology* 23:281–299.
- CUMMINS, K. W., AND M. J. KLUG. 1979. Feeding ecology of stream invertebrates. *Annual Review of Ecology, Evolution, and Systematics* 10:147–172.
- DANG, C. K., M. SCHINDLER, E. CHAUVET, AND M. O. GESSNER. 2009. Temperature oscillation coupled with fungal community shifts can modulate warming effects on litter decomposition. *Ecology* 90:122–131.
- DEL GIORGIO, P. A., AND J. M. GASOL. 1995. Biomass distribution in freshwater plankton communities. *American Naturalist* 146:135–152.
- ENGLISHAW, H. J. 1964. The distributional relationship between the bottom fauna and plant detritus in streams. *Journal of Animal Ecology* 33:463–476.
- GESSNER, M. O., AND E. CHAUVET. 2002. A case for using litter breakdown to assess functional stream integrity. *Ecological Applications* 1:498–510.
- GESSNER, M. O., C. M. SWAN, C. K. DANG, B. G. MCKIE, R. D. BARDGETT, D. H. WALL, AND S. HÄTTENSCHWILER. 2010. Diversity meets decomposition. *Trends in Ecology and Evolution* 25:372–380.
- GONZÁLEZ, J. M., AND M. A. S. GRAÇA. 2003. Conversion of leaf litter to secondary production by a shredding caddis-fly. *Freshwater Biology* 48:1578–1592.
- GRAÇA, M. A. S. 2001. The role of invertebrates on leaf litter decomposition in streams: a review. *International Review of Hydrobiology* 86:383–393.
- HAMMER, Ø., D. A. T. HARPER, AND P. D. RYAN. 2001. PAST: Paleontological Statistics Software Package for Education and Data Analysis. *Palaeontologia Electronica* 4(1):9.
- HAY, M. E., J. J. STACHOWICZ, E. CRUZ-RIVERA, S. BULLARD, M. S. DEAL, AND N. LINDQUIST. 1998. Bioassays with marine and freshwater macroorganisms. Pages 39–141 in K. F. HAYNES and J. G. MILLAR (editors). *Methods in chemical ecology*. Volume 2. Bioassay methods. Chapman and Hall, New York.
- HENRIKSON, A. A., AND J. R. PAWLIK. 1995. A new antifouling assay method: results from field experiments using extracts of four marine organisms. *Journal of Experimental Marine Biology and Ecology* 194:157–165.
- HIEBER, M., AND M. O. GESSNER. 2002. Contribution of stream detritivores, fungi, and bacteria to leaf breakdown based on biomass estimates. *Ecology* 83:1026–1038.
- HILDREW, A. G., C. R. TOWNSEND, J. FRANCIS, AND K. FINCH. 1984. Cellulolytic decomposition in streams of contrasting pH and its relationship with invertebrate community structure. *Freshwater Biology* 14:323–328.
- HUNTING, E. R., J. M. DE GOEIJ, M. ASSELMAN, R. W. M. VAN SOEST, AND H. G. VAN DER GEEST. 2010a. Degradation of mangrove-derived organic matter in mangrove associated sponges. *Bulletin of Marine Science* 86:871–877.
- HUNTING, E. R., H. G. VAN DER GEEST, A. J. KRIEG, M. B. L. VAN MIERLO, AND R. W. M. VAN SOEST. 2010b. Mangrove-sponge associations: a possible role for tannins. *Aquatic Ecology* 44:679–684.

- HUNTING, E. R., M. H. WHATLEY, H. G. VAN DER GEEST, C. MULDER, M. H. S. KRAAK, T. BREURE, AND W. ADMIRAAL. 2012. Invertebrate footprints on detritus processing, bacterial community structure and spatiotemporal redox profiles. *Freshwater Science* 31:723–731.
- IMBERGER, S. J., R. M. THOMPSON, AND M. R. GRACE. 2010. Searching for effective indicators of ecosystem function in urban streams: assessing cellulose decomposition potential. *Freshwater Biology* 55:2089–2106.
- LECERF, A., AND E. CHAUVET. 2008. Intraspecific variability in leaf traits strongly affects alder leaf decomposition in a stream. *Basic and Applied Ecology* 9:598–605.
- MAAS, J. L., C. VAN DE GÜCHTE, AND F. C. M. KERKUM. 2002. Methodbeschrijvingen voor de beoordeling van verontreinigde waterbodems volgens de TRIADE benadering. Rijksinstituut voor Integraal Zoetwaterbeheer en Afvalwaterbehandeling nota 93.027. (Available from: www.vliz.be/imisdocs/publications/155685.pdf)
- MONARI, M., J. FOSCHI, P. CORTESI, R. ROSMINI, O. CATTANI, AND G. P. SERRAZANETTI. 2008. Chloramphenicol influence on antioxidant enzymes with preliminary approach on microsomal CYP1A immunopositive-protein in *Chamelea gallina*. *Chemosphere* 73:272–280.
- NIKI, E. 1991. Action of ascorbic acid as a scavenger of active and stable oxygen radicals. *American Journal of Clinical Nutrition* 54:1119–1124.
- PAVIA, H., AND G. B. TOTH. 2000. Inducible chemical resistance to herbivory in the brown seaweed *Ascophyllum nodosum*. *Ecology* 81:3212–3225.
- PETERSEN, R. C., AND K. W. CUMMINS. 1974. Leaf processing in a woodland stream. *Freshwater Biology* 4:343–368.
- RADER, R. B., J. V. MCARTHUR, AND J. M. AHO. 1994. Relative importance of mechanisms determining decomposition in a southeastern blackwater stream. *American Midland Naturalist* 132:19–31.
- SUBERKROPP, K. 1991. Relationships between growth and sporulation of aquatic hyphomycetes on decomposing leaf litter. *Mycological Research* 95:843–850.
- TANK, J. L., E. J. ROSI-MARSHALL, N. A. GRIFFITHS, S. A. ENTREKIN, AND M. L. STEPHEN. 2010. A review of allochthonous organic matter dynamics and metabolism in streams. *Journal of the North American Benthological Society* 29:118–146.
- TIEGS, S. D., S. D. LANGHANS, K. TOCKNER, AND M. O. GESSNER. 2007. Cotton strips as a leaf surrogate to measure decomposition in river floodplain habitats. *Journal of the North American Benthological Society* 26:70–77.
- WALLACE, J. B., AND J. R. WEBSTER. 1996. The role of macroinvertebrates in stream ecosystem function. *Annual Review of Entomology* 41:115–139.
- WEBSTER, J. R., AND E. F. BENFIELD. 1986. Vascular plant breakdown in freshwater ecosystems. *Annual Review of Ecology and Systematics* 17:567–594.
- WOODWARD, G., M. O. GESSNER, P. S. GILLER, V. GULIS, S. HLADYZ, A. LECERF, B. MALMQVIST, B. G. MCKIE, S. D. TIEGS, H. CARISS, M. DOBSON, A. ELOSEGI, V. FERREIRA, M. A. S. GRAÇA, T. FLEITUCH, J. O. LACOURSIERE, M. NISTORESCU, J. POZO, G. RISNOVEANU, M. SCHINDLER, A. VADINEANU, L. B.-M. VOUGHT, AND E. CHAUVET. 2012. Continental-scale effects of nutrient pollution on stream ecosystem functioning. *Science* 336:1438–1440.
- YOUNG, R. G., AND K. J. COLLIER. 2009. Contrasting responses to catchment modification among a range of functional and structural indicators of river ecosystem health. *Freshwater Biology* 54:2155–2170.
- YOUNG, R. G., C. MATTHAEI, AND C. R. TOWNSEND. 2008. Organic matter breakdown and ecosystem metabolism: functional indicators for assessing river ecosystem health. *Journal of the North American Benthological Society* 27:605–625.
- ZWART, G., B. C. CRUMP, M. AGTERVELD, F. HAGEN, AND S. K. HAN. 2002. Typical freshwater bacteria: an analysis of available 16S rRNA gene sequences from plankton of lakes and rivers. *Aquatic Microbial Ecology* 28: 141–155.
- ZWART, G., W. D. HIORNS, B. A. METHÉ, M. P. VAN AGTERVELD, R. HUISMANS, S. C. NOLD, J. P. ZEHR, AND H. J. LAANBROEK. 1998. Nearly identical 16S rRNA sequences recovered from lakes in North America and Europe indicate the existence of clades of globally distributed freshwater bacteria. *Systematic and Applied Microbiology* 21: 546–556.

Received: 16 May 2012

Accepted: 10 August 2012