



## **Interaction between the Autosomal Recessive bar Gene and the Y-Linked Snakeskin Body (Ssb) Pattern Gene in the Guppy, *Poecilia reticulata***

Authors: Phang, Violet P. E., Khoo, Gideon, and Ang, Sock Peng

Source: Zoological Science, 16(6) : 905-908

Published By: Zoological Society of Japan

URL: <https://doi.org/10.2108/zsj.16.905>

---

BioOne Complete ([complete.BioOne.org](https://complete.BioOne.org)) is a full-text database of 200 subscribed and open-access titles in the biological, ecological, and environmental sciences published by nonprofit societies, associations, museums, institutions, and presses.

Your use of this PDF, the BioOne Complete website, and all posted and associated content indicates your acceptance of BioOne's Terms of Use, available at [www.bioone.org/terms-of-use](https://www.bioone.org/terms-of-use).

Usage of BioOne Complete content is strictly limited to personal, educational, and non - commercial use. Commercial inquiries or rights and permissions requests should be directed to the individual publisher as copyright holder.

---

BioOne sees sustainable scholarly publishing as an inherently collaborative enterprise connecting authors, nonprofit publishers, academic institutions, research libraries, and research funders in the common goal of maximizing access to critical research.

# Interaction between the Autosomal Recessive *bar* Gene and the Y-Linked Snakeskin Body (*Ssb*) Pattern Gene in the Guppy, *Poecilia reticulata*

Violet P. E. Phang\*, Gideon Khoo and Sock Peng Ang

Department of Biological Sciences, Faculty of Science, National University of Singapore,  
14 Science Drive 4, Singapore 117543, Republic of Singapore


**ABSTRACT**—Many color varieties of the guppy, *Poecilia reticulata*, are commercially cultured in Singapore for the aquarium industry. In the group of guppy varieties called Snakeskin, males characteristically have snakeskin-like reticulations on the body and caudal fin. The snakeskin pattern on the body of male Snakeskin guppies is due to a Y-linked gene (*Ssb*). Female guppies, being homogametic (XX), do not carry the *Ssb* gene. About 90% of Yellow Snakeskin males have the typical snakeskin pattern on their bodies and tails. The remaining males are different in that the snakeskin body pattern has been modified into four or five vertical bars on the caudal-peduncle region. F<sub>1</sub> and F<sub>2</sub> results of single-pair reciprocal matings of the Yellow Snakeskin variety show that a single gene is responsible for the vertical bar pattern. This gene, *bar*, is autosomal recessive. In the homozygous condition (*barbar*), it interacts with the Y-linked *Ssb* gene to give vertical barring patterns on the caudal-peduncle of Yellow Snakeskin males. This pattern is not expressed when the dominant allele, *bar*<sup>+</sup>, is present.

## INTRODUCTION

The guppy is unique among other teleosts in that almost all the genes responsible for pigmentation and color patterns, with the exception of background body coloration genes, are sex-linked and sex-limited. It is the first organism in which Y-linked inheritance of color genes was demonstrated (Schmidt, 1920; Winge, 1922a, b, 1927). The guppy has 23 pairs of chromosomes, of which 22 are autosomal and one pair the sex chromosomes. Males are heterogametic (XY) while females are homogametic (XX) (Winge, 1922a, b; Winge and Ditlevsen, 1947). Expression of color patterns in domesticated varieties is due to dominant sex-linked genes (Dzwillo, 1959; Nayudu, 1979; Fernando and Phang, 1989; Phang *et al.*, 1989a, b, 1990; Phang and Fernando, 1991; Khoo *et al.*, 1999a, b). The color patterns of these varieties were initially selected from a large gene pool in wild-type populations. Kirpichnikov (1981), in his review, documented 17 Y-linked genes that are passed from father to son through the Y-chromosome, 15 that are X- and Y-linked (found in both males and females but expressed only in males because they are sex-limited and hormone-mediated), and one that is autosomal dominant. In contrast, genes responsible for background body coloration such as blond (*b*), gold (*g*), albino (*a*) and

blue (*bl*) are autosomally inherited and recessive to their wild-type alleles (Kirpichnikov, 1981).

Color patterns on the body and caudal fin of domesticated guppies take the form of single bright colors, snakeskin-like reticulations and variegated mosaic patterns of two or more colors (Fernando and Phang, 1985, 1989; Phang *et al.*, 1989a, b, 1990; Phang and Fernando, 1991; Khoo *et al.*, 1999a, b). Recent surveys of guppy farms in Singapore show



**Fig. 1.** An adult Yellow Snakeskin (YSSbar) male guppy showing the vertical bar pattern (arrow) on its caudal-peduncle region (top) and a normal Yellow Snakeskin (YSS) male without the bar pattern (bottom).

\* Corresponding author: Tel. (65)-874-2694;  
FAX. (65)-779-2486.  
E-mail: dbsphang@nus.edu.sg

the popularity of snakeskin-like and variegated patterns among guppy varieties that are cultured for export (Khoo *et al.*, 1999a). The iridescent snakeskin pattern on the body and tail is the result of two closely linked genes, Snakeskin-body (*Ssb*) and Snakeskin-tail (*Sst*), that are expressed only in males (Phang *et al.*, 1989a, b, 1990; Phang and Fernando, 1991). *Ssb* and *Sst* are thus absent in the homogametic (XX) females as these genes are Y-linked. Observations of Yellow Snakeskin males obtained from stocks in Singapore guppy farms show that about 10% of them differ from normal snakeskin males in having at least four to five prominent iridescent vertical bars on the caudal peduncle region instead of the characteristic r-eticulated snakeskin-like network pattern over the entire body (Fig. 1). Using the Yellow Snakeskin guppy variety as a genetic model, we undertook this study to investigate the genetic basis of the bar pattern.

MATERIALS AND METHODS

Source of the fish

Three- to four-week old fry of the Yellow Snakeskin guppy variety were obtained from Swee Hing & Brothers Aquarium Co. in Singapore. YSS juveniles were separated according to sex and cultured as in Khoo *et al.* (1999a, b).

Description of the fish

The common name, Yellow Snakeskin (YSS), is given by commercial guppy breeders to males ( ) and females ( ) of this guppy variety. The color phenotypes of adult Yellow Snakeskin males are Yellow Snakeskin without bar pattern (YSS ) and Yellow Snakeskin with bar (YSSbar ) as shown in Fig. 1. YSS and YSSbar males have pale yellowish background body coloration and a yellow caudal fin that is overlaid with delicate snakeskin-like reticulations. The two phenotypes differ in that YSS males have the characteristic reticulated snakeskin pattern over the whole body while the snakeskin pattern of YSSbar males is modified to four or five prominent

vertical bars on the caudal-peduncle region (Fig. 1). All Yellow Snakeskin females (YSS ) show only pale yellow background coloration on the body and tail.

Reciprocal matings

Single-pair reciprocal matings of males (YSS and YSSbar ) and females (YSS ) of the Yellow Snakeskin variety were set up to determine the inheritance of the bar pattern. The following notations were used: YSS × YSS (Mating 1) and YSSbar × YSS (Mating 2) (Table 1). Single-pair full-sib F<sub>1</sub> males and females were mated to obtain the F<sub>2</sub> generation. Each mating pair (two-month old mature virgin fish) was kept in a 3.5-liter breeding tank. Broods were usually produced 4-6 weeks after mating. F<sub>1</sub> and F<sub>2</sub> offspring were segregated according to phenotypes and sex after about eight weeks of age. Maintenance and grow-out of newly born fry and juveniles were as described by Khoo *et al.* (1999a, b).

Statistical analyses

Observed phenotypic distributions were tested for goodness-of-fit with predicted proportions using the chi-square ( $\chi^2$ ) test (Sokal and Rohlf, 1981; Strickberger, 1990). Since observed and expected numbers in the phenotypic classes and sample sizes were small ( $n < 200$ ), Yates' (1934) correction for continuity was included in the calculation of  $\chi^2$  to improve the approximation to the  $\chi^2$  distribution, as shown by the  $\chi^2_{adj}$  values.

RESULTS AND DISCUSSION

Segregation of *bar* in F<sub>1</sub> and F<sub>2</sub> offspring of YSS × YSS

Single-pair matings of Yellow Snakeskin (YSS) males and females gave two different groups of F<sub>2</sub> progenies (Table 1, Fig. 2). The first group (Mating 1a) of eight single-pair matings produced F<sub>1</sub> and F<sub>2</sub> offspring where all the males had the reticulated snakeskin-like body pattern on a pale yellowish body and yellow caudal fin that is characteristic of the Yellow Snakeskin phenotype (Figs. 1, 2). All the females were also

**Table 1.** Segregation data of the F<sub>1</sub> and F<sub>2</sub> generations of single-pair reciprocal matings of: normal Yellow Snakeskin males (YSS ) and Yellow Snakeskin females (YSS ) (Mating 1a), YSS and YSS (with *bar* pattern gene that was unexpressed due to the absence of the Y-linked *Ssb* gene) (Mating 1b), Yellow Snakeskin males with bar pattern (YSSbar ) and YSS (Mating 2a), and YSSbar and YSS (with unexpressed *bar* gene) (Mating 2b). Observed segregation numbers for F<sub>2</sub> male offspring with and without the bar pattern were subjected to chi-square ( $\chi^2$ ) goodness-of-fit analyses. Female progenies were excluded from  $\chi^2$  tests (\*) because those that carried the *bar* gene could not be distinguished from those with the *bar*<sup>+</sup> allele.

Mating (Mating no.)	Generation	No. of matings (No. of broods)	Observed no. for each phenotypic class (Expected no.)			Expect- ed ratio	Chi-square Goodness-of-fit Test ( <i>df</i> =1)	
			YSS	YSSbar	YSS		$\chi^2$	$\chi^2_{adj}$
YSS    ×YSS (Mating 1a)	F <sub>1</sub>	8 (10)	65 (61.50)	0	58 (61.50)	1:1	0.398	0.293
	F <sub>2</sub>	8 (10)	88 (90.50)	0	93 (90.50)	1:1	0.138	0.088
YSS    ×YSS (Mating 1b)	F <sub>1</sub>	3 (5)	17 (17.50)	0	18 (17.50)	1:1	0.029	0.000
	F <sub>2</sub>	3 (5)	20 (19.50)	6 (6.50)	32 (29.00)	3:1*	0.051	0.000
			20 + 6 = 26 (29.00)			1:1	0.621	0.431
YSSbar    × YSS (Mating 2a)	F <sub>1</sub>	7 (7)	85 (81.50)	0	78 (81.50)	1:1	0.301	0.221
	F <sub>2</sub>	6 (6)	24 (23.25)	7 (7.75)	30 (30.50)	3:1*	0.097	0.011
			24 + 7 = 31 (30.50)			1:1	0.016	0.000
YSSbar    × YSS (Mating 2b)	F <sub>1</sub>	6 (6)	0	16 (14.50)	13 (14.50)	1:1	0.310	0.138
	F <sub>2</sub>	5 (5)	0	9 (10.50)	12 (10.50)	1:1	0.429	0.190

\* : expected ratio (3:1) of normal YSS to YSSbar . All other hypothetical ratios (1:1) were for male to female progenies.

Mating 1				Mating 2			
YSS ♂♂		YSS ♀♀		YSSbar ♂♂		YSS ♀♀	
a	$bar^+bar^+ X Y_{Ssb}$	×	$bar^+bar^+ X X$	P	$barbar X Y_{Ssb}$	×	$bar^+bar^+ X X$
b			$barbar X X$				$barbar X X$
a	$bar^+bar^+ X Y_{Ssb}$ YSS	×	$bar^+bar^+ X X$ YSS	F <sub>1</sub>	$barbar^+ X Y_{Ssb}$ YSS	×	$barbar^+ X X$ YSS
b	$bar^+bar X Y_{Ssb}$ YSS	×	$bar^+bar X X$ YSS		$barbar X Y_{Ssb}$ YSSbar	×	$barbar X X$ YSS
a	$bar^+bar^+ X Y_{Ssb}$ YSS		$bar^+bar^+ X X$ YSS	F <sub>2</sub>	$bar^+bar^+ X Y_{Ssb}$ YSS $barbar^+ X Y_{Ssb}$ YSS $bar^+bar X Y_{Ssb}$ YSS $barbar X Y_{Ssb}$ YSSbar		$bar^+bar^+ X X$ YSS $barbar^+ X X$ YSS $bar^+bar X X$ YSS $barbar X X$ YSS
b	$bar^+bar^+ X Y_{Ssb}$ YSS $bar^+bar X Y_{Ssb}$ YSS $barbar^+ X Y_{Ssb}$ YSS $barbar X Y_{Ssb}$ YSSbar		$bar^+bar^+ X X$ YSS $bar^+bar X X$ YSS $barbar^+ X X$ YSS $barbar X X$ YSS		$barbar X Y_{Ssb}$ YSSbar		$barbar X X$ YSS
F <sub>2</sub> Phenotypic ratio				F <sub>2</sub> Phenotypic ratio			
YSS ♂♂ × YSS ♀♀				YSSbar ♂♂ × YSS ♀♀			
Mating 1a: 1 YSS ♂♂ : 1 YSS ♀♀				Mating 2a: 3 YSS ♂♂ : 1 YSSbar ♂♂			
Mating 1b: 3 YSS ♂♂ : 1 YSSbar ♂♂				Mating 2b: 1 YSSbar ♂♂ : 1 YSS ♀♀			

erozygous for *bar*, i.e., *bar*<sup>+</sup>*bar* (Table 1). Thus, parental females of Mating 1b were homozygous for this gene (genotype: *barbarXX*) (Fig. 2). Only male guppies are able to express the bar pattern because they possess the Y-linked *Ssb* gene. The dominant allele of this locus, *bar*<sup>+</sup>, does not modify the snakeskin body pattern of YSS males.

### Segregation of *bar* in F<sub>1</sub> and F<sub>2</sub> offspring of YSSbar × YSS

As described for Mating 1, F<sub>1</sub> and F<sub>2</sub> progenies of Mating 2 could also be separated into Matings 2a and 2b according to their phenotypes (Table 1, Fig. 2). Seven single-pair matings between YSSbar males and YSS females of Mating 2a gave a total of 85 F<sub>1</sub> males and 78 females that were all YSS in the expected 1:1 male to female ratio (Table 1, Fig. 2). In the F<sub>2</sub> generation, however, there were 24 YSS males and seven YSSbar males. These observed numbers of F<sub>2</sub> males concurred with the 3 YSS : 1 YSSbar male ratio expected in the F<sub>2</sub> generation of a monohybrid cross (Table 1, Fig. 2). This shows that the parental YSSbar males in Mating 2a most likely had a genotype of *barbarXY*<sub>Ssb</sub> while the females were *bar*<sup>+</sup>*bar*<sup>+</sup>XX. For Mating 2b, the F<sub>1</sub> and F<sub>2</sub> offspring comprised only YSSbar males and YSS females that conformed to the male to female ratio of 1:1 (Table 1, Fig. 2). These results prove that YSSbar males and YSS females of the parental generation of Mating 2b were homozygous for the autosomal recessive *bar* gene (Fig. 2).

### Inheritance of *bar* and its interaction with the *Ssb* gene

To date, the number of sex-linked color pattern genes described for the guppy far outnumber the autosomal ones (Winge, 1927, 1934; Winge and Ditlevsen, 1947; Dzwillo, 1959; Nayudu, 1979; Kirpichnikov, 1981; Fernando and Phang, 1989; Phang *et al.*, 1989a, b, 1990; Phang and Fernando, 1991; Khoo *et al.*, 1999a, b). Only four genes, blond (*b*), gold (*g*), albino (*a*) and blue (*bl*), that are responsible for background body coloration have been shown to be autosomally inherited in the guppy (Kirpichnikov, 1981). These genes are recessive to their wild-type alleles. Our observations of all parental and full-sib matings (Mating 1: YSS × YSS and Mating 2: YSSbar × YSS) show evidence of Mendelian inheritance of the vertical bar pattern in guppies with reticulated snakeskin-like patterns (Table 1).

This study demonstrates, for the first time, that a single autosomal body pattern gene, *bar*, is responsible for the recessive vertical bar trait in male guppies of Snakeskin varieties. Segregation data for Matings 1 and 2 shows that the occurrence of *bar* in the homozygous condition modifies the Y-linked Snakeskin-body (*Ssb*) gene of Yellow Snakeskin males (Table 1, Fig. 2), resulting in a pattern of four to five vertical bars on the caudal-peduncle region (Fig. 1). Females that are homozygous for the *bar* gene do not express this barring pattern due to absence of *Ssb* (Phang *et al.*, 1989a, b, 1990; Phang and Fernando, 1991). The *Ssb* gene is not modified by the dominant allele of the bar pattern locus, *bar*<sup>+</sup>.

## ACKNOWLEDGEMENTS

This project was funded by a research grant from the National University of Singapore (RP800024) to V.P.E. Phang (P.I.). The authors thank Mr. K.J. Goh for photographing the guppies.

## REFERENCES

- Dzwillo M (1959) Genetische Untersuchungen an domestizierten Stämmen von *Lebistes reticulatus* Peters. Mitt Hamburg Zool Mus Inst 57: 143–186
- Fernando AA, Phang VPE (1985) Culture of the guppy, *Poecilia reticulata*, in Singapore. Aquaculture 51: 49–63
- Fernando AA, Phang VPE (1989) X-linked inheritance of red and blue tail colorations of domesticated varieties of guppy, *Poecilia reticulata*, and its implications to the farmer. Singapore J Pri Ind 17: 10–18
- Khoo G, Lim TM, Chan WK, Phang VPE (1999a) Genetic basis of the variegated tail pattern in the guppy, *Poecilia reticulata*. Zool Sci 16: 431–437
- Khoo G, Lim TM, Chan WK, Phang VPE (1999b) Sex-linkage of the black caudal-peduncle and red tail genes in the Tuxedo strain of the guppy, *Poecilia reticulata*. Zool Sci 16: 629–638
- Kirpichnikov VS (1981) The genetics of aquarium fish species. In: "Genetic Bases of Fish Selection." Translated by GG Gause. Springer-Verlag, Berlin-Heidelberg, Germany, pp 77–103
- Nayudu P (1979) Genetic studies of melanistic color patterns, and atypical sex determination in the guppy, *Poecilia reticulata*. Copeia 1979: 225–231
- Phang VPE, Ng LN, Fernando AA (1989a) Inheritance of the snakeskin color pattern in the guppy, *Poecilia reticulata*. J Hered 80: 393–399
- Phang VPE, Ng LN, Fernando AA (1989b) Genetics of the color of the yellow snakeskin variety of the guppy, *Poecilia reticulata*. Singapore J Pri Ind 17: 19–28
- Phang VPE, Fernando AA, EWK Chia (1990) Inheritance of the color patterns of the blue snakeskin and red snakeskin varieties of the guppy, *Poecilia reticulata*. Zool Sci 7: 419–425
- Phang VPE, Fernando AA (1991) Linkage analysis of the X-linked green tail and blue tail color genes in the guppy, *Poecilia reticulata*. Zool Sci 8: 975–981
- Schmidt J (1920) Racial investigations. IV. The genetic behavior of a secondary sexual character. CR Trav Lab Carlsberg 14: 1–8
- Sokal RR, Rohlf FJ (1981) Biometry. The Principles and Practice of Statistics in Biological Research. 2nd ed, WH Freeman and Co, New York, USA, 859 p
- Strickberger MW (1990) Genetics. 3rd ed, Macmillan Publishing Co, New York, USA, 842 p
- Winge Ö (1922a) A peculiar mode of inheritance and its cytological explanation. J Genet 12: 137–144
- Winge Ö (1922b) One-sided masculine and sex-linked inheritance in *Lebistes reticulatus*. J Genet 12: 145–162
- Winge Ö (1927) The location of eighteen genes in *Lebistes reticulatus*. J Genet 18: 1–43
- Winge Ö, Ditlevsen E (1947) Color inheritance and sex determination in *Lebistes*. Heredity 1: 65–83
- Yates F (1934) Contingency tables involving small numbers and the  $\chi^2$  test. J R Stat Soc (suppl) 1: 217–235

(Received April 30, 1999 / Accepted June 26, 1999)