

Estrogen-Independent Ovary Formation in the Medaka Fish, *Oryzias latipes*

Authors: Kawahara, Tetsuya, and Yamashita, Ichiro

Source: Zoological Science, 17(1) : 65-68

Published By: Zoological Society of Japan

URL: <https://doi.org/10.2108/zsj.17.65>

BioOne Complete (complete.BioOne.org) is a full-text database of 200 subscribed and open-access titles in the biological, ecological, and environmental sciences published by nonprofit societies, associations, museums, institutions, and presses.

Your use of this PDF, the BioOne Complete website, and all posted and associated content indicates your acceptance of BioOne's Terms of Use, available at www.bioone.org/terms-of-use.

Usage of BioOne Complete content is strictly limited to personal, educational, and non - commercial use. Commercial inquiries or rights and permissions requests should be directed to the individual publisher as copyright holder.

BioOne sees sustainable scholarly publishing as an inherently collaborative enterprise connecting authors, nonprofit publishers, academic institutions, research libraries, and research funders in the common goal of maximizing access to critical research.

Estrogen-Independent Ovary Formation in the Medaka Fish, *Oryzias latipes*

Tetsuya Kawahara and Ichiro Yamashita*

Center for Gene Science, Hiroshima University, Kagamiyama 1-4-2,
Higashihiroshima 739-8527, Japan

ABSTRACT—To investigate whether a female sex steroid, estrogen, acts as a natural inducer of female gonadal sex determination (or ovary formation) in the medaka fish, *Oryzias latipes*, the effects of an aromatase inhibitor and anti-estrogens on sexual differentiation of gonads were examined. We found that both drugs did not show any discernible effects on the genetically determined sex differentiation in both sexes. However, the aromatase inhibitor impaired the paradoxical effects of androgen (a male sex steroid), and the anti-estrogens inhibited the male-to-female sex reversal caused by estrogen. Treatments of the fertilized eggs with androgen disturbed the gonadal sex developments in both sexes, suggesting that sex steroid synthesis is detrimental to the gonadal sex developments in the medaka embryos. These results are consistent with the previous observation that sex steroids are not synthesized before the onset of gonadal sex differentiation, and suggest that ovary formation in the genetic females of the medaka fish is not dependent on estrogen.

INTRODUCTION

Gonadal sex determination is a process of choice between testicular and ovarian developments from the same primordial gonad. Sex determination in non-mammalian vertebrates may be fundamentally different from that of mammals, and is thought to be under the control of genetics and various environmental factors. In mammals, sex steroids (androgen and estrogen) are considered gonadal products synthesized as a result of gonadal sex differentiation, and they are believed to act in establishing the secondary sexual phenotype. However, this is clearly not the case in lower vertebrates, in which sex steroids can affect or reverse gonadal sex development if applied early enough during development. In chickens (Scheib, 1983) and turtles (Dorizzi *et al.*, 1991), estrogens are synthesized by morphologically undifferentiated female gonads. Treatment with an aromatase inhibitor (which blocks the synthesis of estrogen from androgen) causes females to develop testes (Elbrecht and Smith, 1992; Richard-Mercier *et al.*, 1995). Treatment with an anti-estrogen (which binds to estrogen receptors and compete with estrogens) does not disrupt but disturb ovarian development showing partial masculinization (some testicular appearance of female gonads) (Scheib, 1983; Dorizzi *et al.*, 1991). The evidence is in favor of a role for estrogen in ovarian differentia-

tion. However, it is not demonstrated unambiguously whether estrogen is identical with a natural sex inducer in female and whether there exists another, estrogen-independent pathway for ovarian development, parallel to the estrogen-dependent one. In contrast, histological and ultrastructural studies in some fishes including medaka, *Oryzias latipes*, reveal that steroid hormone biosynthesis and steroid-producing cells appear after the completion of gonadal sex differentiation (Iwasaki, 1973; Takahashi and Iwasaki, 1973; Kagawa and Takano, 1979; Schreiber *et al.*, 1982; van den Hurk *et al.*, 1982; Kanamori *et al.*, 1985). This suggests that gonadal sex determination does not depend on sex steroids in these fishes. However, pharmacological studies of these fishes using aromatase inhibitors and anti-estrogens have not been done, thus, the view remains an open question. We show here that both drugs have no discernible effects on genetically determined gonadal development of medaka fish, indicating that female sex determination of this fish is independent of estrogen.

MATERIALS AND METHODS

We used the d-rR strain of medaka with orange-red (in body color) male (X^Y^R) and white female (X^X) (Yamamoto, 1969). The genotype of sex can be judged by body color with more than 99% reliability (Yamamoto, 1969). Morphological sex differentiation of germ cells occurs at the time of hatching (Sato and Egami, 1972; Hamaguchi, 1982). The fertilized eggs (1- or 2-day old after fertilization) were incubated in Yamamoto's salt solution at 25°C under artificial photoperiod of 14L:10D. The fry were then transferred to plastic aquaria and reared to adult by powdered TetraMin under the same condition for 3 months. Adult fish were dissected for sexing gonads under a

* Corresponding author: Tel. +81-824-24-6271;
FAX. +81-824-22-7184.
E-mail: iyama@ipc.hiroshima-u.ac.jp
iyama@hiroshima-u.ac.jp
(after April in 2000)

dissecting microscope.

RESULTS AND DISCUSSION

Treatment with aromatase inhibitor and androgen

Fertilized eggs were incubated in the presence of sublethal concentrations of an aromatase inhibitor, and after hatching fry were reared to adult by normal diet, dissected for sexing gonads. We could not detect any defects in both ovarian and testicular developments (Table 1): each sex developed normal size of ovary or testis which contained normal number of oocytes or sperms, respectively. The result is consistent with the notion that sex steroids are not synthesized in embryos and not involved in sex determination.

To examine whether exogenous androgen may disturb gonadal differentiation, eggs were incubated in the presence of methylandrostenediol (MA) (Table 1). Some genetic females sex-reversed to males with testes. This contrasts to the absence of sex-reversal by androgen treatment of chickens and turtles in which female embryos contain aromatase activity (Elbrecht and Smith, 1992; Richard-Mercier *et al.*, 1995). Higher dosages of MA frequently produced fishes without visible gonads under a dissecting microscope, indicating the toxic action of androgen (or its metabolites) (Yamamoto, 1969). MA also sex-reversed from genetic males to females with ovaries. This was unexpected because oral administration of androgen does not cause the male-to-female sex reversal in medaka (Yamamoto, 1969). This paradoxical effect of androgen is probably the result of conversion of androgen to estrogen by aromatase present in male embryos because co-administration of aromatase inhibitor impaired the androgen effect. Another androgen, methyltestosterone, showed essentially the same results as MA did (data not shown). Taken together, androgen synthesis may be detrimental to genetically determined gonadal development in medaka embryos: in female embryos, even a low level of androgen could not be

aromatized to estrogen and induced male gonadal development; and in male embryos, androgen could be aromatized to estrogen which induced female gonadal development.

Treatment with anti-estrogens

We next examined for the effects of anti-estrogens on female gonadal differentiation (Table 2). It is well known that newly hatched fry of genetic males can be sex-reversed by oral administration of estrogen (Yamamoto, 1969), where gonadogenesis of the sex-reversed fry is morphologically the same as that of genetic female (Onitake, 1972). The sex reversal caused by oral administration of 17 β -estradiol was completely inhibited by co-administration of anti-estrogen, tamoxifen, or partially by another anti-estrogen, ICI182780. We further examined for the effects of anti-estrogens on newly hatched fry destined to be female by incubating eggs in the presence of 17 β -estradiol. Both anti-estrogens again impaired the male-to-female sex reversal. These results indicate that an estrogen receptor is involved in the estrogen-induced sex-reversal. However, each anti-estrogen could not affect normal ovarian development of genetic females at all. We also found that treatments of fertilized eggs with sublethal concentrations of anti-estrogens could not affect the genetically determined gonadal developments (data not shown). These results further support the conclusion that female sex determination in medaka is not dependent on estrogen. Recently, Miyata and colleagues (1999) reported the lack of detectable effects of an aromatase inhibitor on the gonadal sex differentiation in *Xenopus laevis*. They also found that expression of the aromatase gene was absent at the beginning of the estradiol-sensitive period and proposed that estradiol synthesis is not naturally involved in the gonadal sex differentiation.

If our conclusion is correct, one would expect that partial masculinization by anti-estrogens of female gonads in chickens and turtles could be explained by the presence of two parallel pathways for ovarian development: one is estrogen-dependent and another estrogen-independent. We assume that activation of estrogen receptor plays a key role in both pathways: one activator is estrogen and another activation mechanism may be similar to those recently proposed as ligand-independent activation of estrogen receptor (Power *et al.*, 1991; Zwijzen *et al.*, 1998). In medaka fish and *Xenopus* frogs, the estrogen-independent activation of estrogen receptor may be a primary pathway to the female gonadal development. In other lower vertebrates such as salmon (Piferrer *et al.*, 1994), amphibia (Yu *et al.*, 1993), and reptiles (Lance and Bogart, 1992), treatment with aromatase inhibitors causes the female-to-male sex reversal or disrupts normal ovarian developments, suggesting the importance of estrogen in the ovary formation in these animals. However, it should be noted formally that the findings with aromatase inhibitors only demonstrate the synthesis of sex steroid hormones during the sex determination period and the ability of androgens to cause female-to-male sex reversal in these animals but not the importance of estrogen as a natural sex inducer in the gonadal differentiation. It remains to be seen whether exogenous

Table 1. Aromatase inhibitor has no effects on gonadal differentiation. Eggs were incubated in Yamamoto's salt solution containing aromatase inhibitor (AI; Fadrozole; Novartis Pharma Inc.) and/or methylandrostenediol (MA) at the indicated concentrations ($\mu\text{g/ml}$), and after hatching fry were reared to adult by normal diet. Each value represents total number of fish carrying the indicated gonad from two or three independent experiments with similar results. O, ovary; T, testis; OT, ovotestis; and NF, gonads not found in abdominal cavity.

AI	MA	Orange-red (X^iY^R)				White (X^iX^i)			
		T	O	OT	NF	O	T	OT	NF
10	—	101	0	0	0	105	0	0	0
20	—	57	0	0	0	65	0	0	1
50	—	33	0	0	0	35	1	0	0
—	0.5	0	4	0	15	5	0	0	11
—	0.1	32	70	7	11	65	4	3	11
—	0.01	17	6	3	3	40	10	0	3
10	0.1	32	2	5	6	36	1	3	3
20	0.1	59	25	6	9	53	1	0	4
50	0.1	6	1	1	0	8	0	3	2

Table 2. Anti-estrogen does not disturb female gonadal development. Newly hatched fry were reared to adult by Tetramin containing 17 β -estradiol (E₂), tamoxifen (TAM), or ICI182780 (ICI; Zeneca Pharmaceuticals) at the indicated concentrations (μ g/g of diet). Eggs were incubated in the presence of E₂ at the indicated concentration (μ g/ml), and after hatching fry were reared to adult by Tetramin or the same diet containing TAM or ICI.

Egg E ₂	Oral after hatching			Orange-red (X ^Y ^R)				White (X ^X ^Y)			
	E ₂	TAM	ICI	T	O	OT	NF	O	T	OT	NF
–	5	–	–	47	0	1	0	32	0	0	0
–	10	–	–	7	9	23	0	19	0	0	0
–	20	–	–	0	106	0	0	105	0	0	1
–	50	–	–	0	21	1	0	22	0	0	0
–	10	100	–	29	6	6	0	17	0	0	0
–	10	1000	–	26	7	5	0	30	0	0	0
–	20	1000	–	42	19	11	0	45	0	0	0
–	20	2000	–	61	0	0	0	60	0	0	0
–	20	5000	–	49	0	0	0	51	0	0	0
–	20	–	2000	23	6	0	2	22	0	0	0
–	–	10	–	11	0	0	0	29	0	0	0
–	–	100	–	16	0	0	0	26	0	0	0
–	–	1000	–	22	0	0	0	43	0	0	0
–	–	2500	–	10	0	0	0	37	0	0	0
–	–	5000	–	24	0	0	0	55	0	0	0
–	–	–	1000	15	0	0	0	15	0	0	0
–	–	–	2000	29	0	0	0	37	0	0	1
0.2	–	–	–	2	89	0	1	75	0	0	0
0.2	–	2000	–	20	40	0	2	74	0	0	0
0.2	–	–	2000	8	19	0	2	34	0	0	0

estrogens can cause the male-to-female sex reversal at the same concentrations found in undifferentiated female embryos or whether anti-estrogens can disrupt female gonadal developments.

ACKNOWLEDGMENTS

We thank H. Okada for help at the early stage of this work. We also thank Novartis Pharma Inc. and Zeneca Pharmaceuticals for supply of drugs.

REFERENCES

- Dorizzi M, Mignot T-M, Guichard A, Desvages G, Pieau C (1991) Involvement of oestrogens in sexual differentiation of gonads as a function of temperature in turtles. *Differentiation* 47: 9–17
- Elbrecht A, Smith RG (1992) Aromatase enzyme activity and sex determination in chickens. *Science* 255: 467–470
- Hamaguchi S (1982) A light- and electron-microscopic study on the migration of primordial germ cells in the teleost, *Oryzias latipes*. *Cell Tissue Res* 227: 139–151
- Iwasaki Y (1973) Histochemical detection of Δ^5 -3 β -hydroxysteroid dehydrogenase in the ovary of medaka, *Oryzias latipes*, during annual reproductive cycle. *Bull Fac Fish Hokkaido Univ* 23: 177–184
- Kagawa H, Takano K (1979) Ultrastructure and histochemistry of granulosa cells of pre- and post-ovulatory follicles in the ovary of the medaka, *Oryzias latipes*. *Bull Fac Fish Hokkaido Univ* 30: 191–204
- Kanamori A, Nagahama Y, Egami N (1985) Development of the tissue architecture in the gonads of the medaka *Oryzias latipes*. *Zool Sci* 2: 695–706
- Lance VA, Bogart MH (1992) Disruption of ovarian development in alligator embryos treated with an aromatase inhibitor. *Gen Comp Endocrinol* 86: 59–71
- Miyata S, Koike S, Kubo T (1999) Hormonal reversal and the genetic control of sex differentiation in *Xenopus*. *Zool Sci* 16: 335–340
- Onitake K (1972) Morphological studies of normal sex-differentiation and induced sex-reversal process of gonads in the medaka, *Oryzias latipes*. *Annot Zool Japon* 45: 159–169
- Piferrer F, Zanuy S, Carrillo M, Solar II, Devlin RH, Donaldson EM (1994) Brief treatment with an aromatase inhibitor during sex differentiation causes chromosomally female salmon to develop as normal, functional males. *J Exp Zool* 270: 255–262
- Power RF, Mani SK, Codina J, Conneely OM, O'Malley BW (1991) Dopaminergic and ligand-independent activation of steroid hormone receptors. *Science* 254: 1636–1639
- Richard-Mercier N, Dorizzi M, Desvages G, Girondot M, Pieau C (1995) Endocrine sex reversal of gonads by the aromatase inhibitor Letrozole (CGS 20267) in *Emys orbicularis*, a turtle with temperature-dependent sex determination. *Gen Comp Endocrinol* 100: 314–326
- Satoh N, Egami N (1972) Sex differentiation of germ cells in the teleost, *Oryzias latipes*, during normal embryonic development. *J Embryol Exp Morph* 28: 385–395
- Scheib D (1983) Effects and role of estrogens in avian gonadal differentiation. *Differentiation* 23 (Suppl): S87–S92
- Schreibman MP, Berkowitz EJ, van den Hurk R (1982) Histology and histochemistry of the testis and ovary of the platyfish, *Xiphophorus maculatus*, from birth to sexual maturity. *Cell Tissue Res* 224: 81–87
- Takahashi H, Iwasaki Y (1973) The occurrence of histochemical activity of 3 β -hydroxysteroid dehydrogenase in the developing testes of *Poecilia reticulata*. *Dev Growth Differ* 15: 241–253
- van den Hurk R, Lambert JGD, Peute J (1982) Steroidogenesis in the gonads of rainbow trout fry (*Salmo gairdneri*) before and after the onset of gonadal sex differentiation. *Reprod Nutr Dev* 22: 413–425
- Yamamoto T (1969) Sex differentiation In "Fish Physiology 3" Ed by WS Hoar, DJ Randall Academic Press, New York, pp 117–175

Yu NW, Hsu CY, Ku HH, Chang LT, Liu HW (1993) Gonadal differentiation and secretions of estradiol and testosterone of ovaries of *Rana catesbiana* tadpoles treated with 4-hydroxyandrostenedione. *J Exp Zool* 265: 252–257

Zwijsen RML, Buckle RS, Hijmans EM, Loomans CJM, Bernards R (1998) Ligand-independent recruitment of steroid receptor coactivators to estrogen receptor by cyclin D1. *Genes Dev* 12: 3488–3498

(Received June 16, 1999 / Accepted July 14, 1999)