



## **Biosynthesis and Biological Actions of Neurosteroids in Brain Neurons**

Author: Tsutsui, Kazuyoshi

Source: Zoological Science, 18(8) : 1043-1053

Published By: Zoological Society of Japan

URL: <https://doi.org/10.2108/zsj.18.1043>

---

BioOne Complete ([complete.BioOne.org](http://complete.BioOne.org)) is a full-text database of 200 subscribed and open-access titles in the biological, ecological, and environmental sciences published by nonprofit societies, associations, museums, institutions, and presses.

Your use of this PDF, the BioOne Complete website, and all posted and associated content indicates your acceptance of BioOne's Terms of Use, available at [www.bioone.org/terms-of-use](http://www.bioone.org/terms-of-use).

Usage of BioOne Complete content is strictly limited to personal, educational, and non - commercial use. Commercial inquiries or rights and permissions requests should be directed to the individual publisher as copyright holder.

---

BioOne sees sustainable scholarly publishing as an inherently collaborative enterprise connecting authors, nonprofit publishers, academic institutions, research libraries, and research funders in the common goal of maximizing access to critical research.

**[REVIEW]**

# Biosynthesis and Biological Actions of Neurosteroids in Brain Neurons

Kazuyoshi Tsutsui\*

*Laboratory of Brain Science, Faculty of Integrated Arts and Sciences, Hiroshima University, Higashi-Hiroshima 739-8521, and Core Research for Evolutional Science and Technology (CREST), Japan Science and Technology Corporation, Tokyo, Japan*

## CONTENTS

1. INTRODUCTION TO NEUROSTEROID RESEARCH
  - (1) *Classical concept: Brain is a target site of peripheral steroids*
  - (2) *New evidence: Brain is also a steroidogenic site*
  - (3) *A model for the study of biosynthesis and biological actions of neurosteroids*
2. NEUROSTEROIDS IN VERTEBRATE BRAINS
  - (1) *Biosynthesis of neurosteroids in mammalian brains*
  - (2) *Biosynthesis of neurosteroids in nonmammalian brains*
  - (3) *Biosynthetic pathway of neurosteroids*
3. PHYSIOLOGICAL CHANGES IN NEUROSTEROIDS IN THE VERTEBRATE BRAIN
4. NEUROSTEROIDOGENESIS IN BRAIN NEURONS
  - (1) *Identification of neurosteroidogenic cells*
  - (2) *Purkinje neuron is a major site of neurosteroidogenesis*
  - (3) *Other neurons*
5. BIOLOGICAL ACTIONS OF NEUROSTEROIDS PRODUCED IN PURKINJE NEURONS
  - (1) *Purkinje neuron serves as an excellent model for the study of neurosteroid actions*
  - (2) *Nongenomic actions as a novel neuromodulator of neurotransmission*
  - (3) *Genomic actions on neuronal growth and synaptogenesis*
6. OTHER EVIDENCE FOR NEUROSTEROID ACTIONS
7. CONCLUSIONS AND FUTURE DIRECTIONS

**Key words:** neurosteroids, biosynthesis, genomic action, nongenomic action, neurons

## 1. INTRODUCTION TO NEUROSTEROID RESEARCH

### **(1) Classical concept: Brain is a target site of peripheral steroids**

Steroid hormones supplied by the peripheral steroidogenic glands regulate several important brain functions during development which persist into adulthood in vertebrates. Peripheral steroid hormones cross the blood-brain barriers, due to their chemically lipid solubility, and act on brain tissues through intracellular receptor-mediated mechanisms that regulate the transcription of specific genes (Fuxe *et al.*, 1981; McEwen, 1991). By diverse actions on the brain, peripheral steroids, in particular sex steroids, have profound effects on behavior of vertebrate animals. Extensive studies have been conducted to understand the mechanisms for steroid actions

on several kinds of behaviors including courtship, copulatory, aggressive and parental behaviors. Some approach has been to measure sex steroid levels in blood, and to correlate these hormone levels with the display of discrete behaviors. These kinds of studies on a variety of wild and captive, intact and castrated, reproductive and non-reproductive animals have established a relationship between the presence and activation of the gonads, increased blood levels of androgens, estrogens or progestins in adult males and females, and expression of adult typical reproductive behaviors.

Gonadal androgens, for instance, act on the brain to influence several male reproductive behaviors in vertebrates. Castration of adult male birds leads to decreases or losses of aggressive, courtship, and copulatory behaviors and replacement therapy with androgens restores these behaviors (Adkins and Adler, 1972; Arnold, 1975; Pröve, 1978, Tsutsui and Ishii, 1981; Ishii and Tsutsui, 1982; Balthazart, 1983; Wingfield and Marler, 1988; Wingfield and Farner, 1993). Many of the brain regions that control a variety of reproductive behaviors con-

\* Corresponding author: Tel. +81 824-24-6571;  
FAX. +81 824-24-0759.  
E-mail: tsutsui@hiroshima-u.ac.jp

tain a high proportion of cells that concentrate androgenic hormones in male birds (Arnold *et al.*, 1976; Korsia and Bottjer, 1989; Watson and Adkins-Regan, 1989). Therefore, the brain is considered to be a target site of peripheral steroids.

### **(2) New evidence: Brain is also a steroidogenic site**

As mentioned above, a great deal was known about the brain as a target site of steroid hormones more than 10 years before. On the other hand, new findings from several laboratories over the past decade have established unequivocally that the nervous system itself forms steroids *de novo* from cholesterol. The pioneering discovery of Baulieu and his colleagues using mammals (for a review, see Baulieu, 1997) and our studies with nonmammals (for a review, see Tsutsui *et al.*, 1999) have opened the door of a new research field. The new concept that steroids could be synthesized *de novo* in the brain derived from observations made by Baulieu and colleagues. They found that several steroids such as pregnenolone, dehydroepiandrosterone, and their sulfate and lipoidal esters highly accumulated within the brain of several mammalian species (Corpéchet *et al.*, 1981, 1983; Robel and Baulieu, 1985; Lanthier and Patwardhan, 1986; Robel *et al.*, 1986, 1987; Jo *et al.*, 1989; Mathur *et al.*, 1993). The brain content of these steroids remained constant even after the removal of peripheral steroids by procedures such as adrenalectomy, castration and hypophysectomy. These results suggested that the brain can synthesize steroids *de novo* from cholesterol (Corpéchet *et al.*, 1981, 1983; Robel and Baulieu, 1985; Robel *et al.*, 1986, 1987; Jo *et al.*, 1989). In contrast to mammalian studies, little has been known regarding *de novo* steroidogenesis in the brain of nonmammalian vertebrates. We therefore looked for steroids formed from cholesterol in the brain of birds (Tsutsui and Yamazaki, 1995; Usui *et al.*, 1995; Tsutsui *et al.*, 1997a, 1997b; Ukena *et al.*, 1999b, 2001; Matsunaga *et al.*, 2001; Tsutsui and Schlinger, 2001), amphibians (Takase *et al.*, 1999) and fish (Sakamoto *et al.*, 2001a). Independently, other groups, such as Vaudry's laboratory (Mensah-Nyagan *et al.*, 1994) and Schlinger's laboratory (Vanson *et al.*, 1996), also contributed to this area. The formation of several steroids from cholesterol is now known to occur in both mammalian and nonmammalian vertebrates. Such steroids synthesized in vertebrate brains are called neurosteroids.

### **(3) A model for the study of biosynthesis and biological actions of neurosteroids**

When we understand the physiological role of neurosteroids in brain functions, it is essential to identify the cells involved in neurosteroidogenesis. In recent years knowledge has been accumulated in both mammals and nonmammals that glial cells play an important role in neurosteroid formation and metabolism in the brain. Both oligodendrocytes and astrocytes are considered to be the primary site for pregnenolone synthesis, an initial step of neurosteroidogenesis. However, whether neurons located in the brain produce neurosteroids has remained unclear. With these findings as a

background, we have demonstrated the presence and activity of neurosteroidogenic enzymes in brain neurons. Interestingly, the Purkinje cell, a typical cerebellar neuron, possesses neurosteroidogenic enzymes and produces neurosteroids *de novo* in a variety of vertebrates including mammalian species (Usui *et al.*, 1995; Tsutsui *et al.*, 1997a, 1997b; Ukena *et al.*, 1998; Takase *et al.*, 1999; Ukena *et al.*, 1999a). This is the first discovery of neuronal *de novo* neurosteroidogenesis in the brain and serves as an excellent model for the study of neurosteroid actions in the brain.

This review summarizes the advances made in our understanding of biosynthesis and biological actions of neurosteroids in neurons. For detailed information of neurosteroids in glial cells the reader is referred to excellent reviews (Baulieu, 1997; Compagnone and Mellon, 2000).

## **2. NEUROSTEROIDS IN VERTEBRATE BRAINS**

### **(1) Biosynthesis of neurosteroids in mammalian brains**

Pregnenolone, a 3 $\beta$ -hydroxy- $\Delta^5$ -steroid, is a main precursor of steroid hormones secreted by peripheral steroidogenic glands, such as gonads, adrenals, and placentae. The formation of pregnenolone is initiated by the cleavage of the cholesterol side-chain by cytochrome P450<sub>scc</sub>, a rate-limiting mitochondrial enzyme originally found in peripheral steroidogenic glandular cells. Therefore, it is essential to demonstrate the formation of pregnenolone in the brain. A number of studies with mammals have reported that the brain contains abundant quantities of 3 $\beta$ -hydroxy- $\Delta^5$ -steroids, i.e., pregnenolone, dehydroepiandrosterone, and their fatty acid or sulfate esters. Furthermore, the content of these steroids in the brain is virtually constant even after the removal of peripheral steroids (Corpéchet *et al.*, 1981, 1983; Robel and Baulieu, 1985; Lanthier and Patwardhan, 1986; Robel *et al.*, 1986, 1987; Jo *et al.*, 1989; Mathur *et al.*, 1993; Ukena *et al.*, 1998). It has also been demonstrated that certain structures in the mammalian brain possess the P450<sub>scc</sub> enzyme (Hu *et al.*, 1987; Le Goascogne *et al.*, 1987; Jung-Testas *et al.*, 1989; Iwahashi *et al.*, 1990; Papadopoulos *et al.*, 1992; Mellon and Deschepper, 1993; Compagnone *et al.*, 1995a; Ukena *et al.*, 1998). Recent studies have shown that both P450<sub>scc</sub> protein and its messenger RNA (mRNA) are expressed in the rat brain (Jung-Testas *et al.*, 1989; Baulieu and Robel, 1990; Baulieu, 1991; Mellon and Deschepper, 1993; Compagnone *et al.*, 1995a; Kohchi *et al.*, 1998; Ukena *et al.*, 1998). Thus, the formation of pregnenolone and its ester from cholesterol is well established in the mammalian brain. In contrast, the formation of dehydroepiandrosterone, one of the first neurosteroids identified in the adult rat brain, remains to be clarified, even though glial cells isolated from neonatal rat brains may be able to convert pregnenolone to dehydroepiandrosterone (Zwain and Yen, 1999).

Progesterone, a 3-*oxo*- $\Delta^4$ -steroid, and its metabolites, such as 3 $\alpha$ -dihydroprogesterone and 3 $\alpha$ ,5 $\alpha$ -tetrahydroprogesterone, are also produced and accumulate in the mammalian brain as neurosteroids. The biosynthesis of

progesterone is performed by  $3\beta$ -hydroxysteroid dehydrogenase/ $\Delta^5$ - $\Delta^4$ -isomerase ( $3\beta$ -HSD) which catalyzes the dehydrogenation and isomerization of the  $\Delta^5$ - $3\beta$ -hydroxysteroids (pregnenolone and dehydroepiandrosterone) into  $\Delta^4$ -ketosteroids (progesterone and androstenedione, respectively) and is highly expressed in the peripheral steroidogenic glands (Mason, 1993). The expression of both  $3\beta$ -HSD protein and its mRNA has been reported in mammalian brains (Dupont *et al.*, 1994; Guennoun *et al.*, 1995a; Sanne and Krueger, 1995; Kohchi *et al.*, 1998; Ukena *et al.*, 1999a). In addition,  $3\beta$ -HSD activity has been demonstrated biochemically in mammalian brain tissues and cultured cells (Weidenfeld *et al.*, 1980; Akwa *et al.*, 1993; Kabbadj *et al.*, 1993; Ukena *et al.*, 1999a).

## (2) Biosynthesis of neurosteroids in nonmammalian brains

The concept of *de novo* steroidogenesis from cholesterol in the brain of nonmammalian vertebrates derived from our observations made in the 1990s. As an initial step in the demonstration of pregnenolone biosynthesis in the avian brain, Tsutsui and Yamazaki (1995) measured the concentrations of pregnenolone and its sulfate ester in the quail brain using a specific radioimmunoassay. The pregnenolone concentration in adult birds was much higher in the brain than in plasma. The accumulation of pregnenolone in the quail brain may be largely independent of peripheral steroidogenic glands, because a high level of pregnenolone persisted in the hypophysectomized birds. Pregnenolone sulfate ester was also detectable in the avian brain. Subsequently, the formation of pregnenolone from cholesterol was found in intact mitochondria derived from the quail brain (Tsutsui and Yamazaki, 1995). To investigate the presence of cytochrome P450scc in the quail brain, Tsutsui and Yamazaki (1995) carried out Western immunoblot analysis with an antibody against purified bovine P450scc after SDS-gel electrophoresis of brain homogenates. In the brain, the antibody against P450scc predominantly recognized a protein band of electrophoretic mobility in the proximity of bovine P450scc. A similar result was obtained in the brain of another bird, the ring dove (Clark *et al.*, 1999; Tsutsui *et al.*, 1999; Lea *et al.*, 2001). Taken together, these biochemical and immunochemical studies indicate that avian brains possess cytochrome P450scc and produce pregnenolone from cholesterol (for reviews, see Tsutsui *et al.*, 1997a, 1997b, 1999).

Subsequently we have extended our understanding of pregnenolone biosynthesis in the brains of lower vertebrates. Takase *et al.* (1999) demonstrated that the amphibian brain possesses P450scc and produces pregnenolone and its sulfate ester. The concentrations of pregnenolone and its sulfate ester in the brain of *Xenopus laevis* were greater than those in the gonads and plasma (Takase *et al.*, 1999). An immunoreactive protein band of electrophoretic mobility in the proximity of bovine P450scc was detected in the *Xenopus* brain by Western blot analysis (Takase *et al.*, 1999). As a lower vertebrate species also possesses P450scc in the brain, the presence of P450scc is considered as a conserved property

of vertebrate brains. This is also true for the presence of  $3\beta$ -HSD, because  $3\beta$ -HSD activity has also been found in the brain of both avian (Vanson *et al.*, 1996; Pignataro *et al.*, 1998; Ukena *et al.*, 1999b) and amphibian species (Mensah-Nyagan *et al.*, 1994). Recently, Ukena *et al.* (1999b) demonstrated the expression of  $3\beta$ -HSD mRNA in the avian brain. Ukena and Tsutsui (2001) further demonstrated that the embryonic avian brain actively metabolizes progesterone to  $5\beta$ -dihydroprogesterone. In addition to these nonmammalian vertebrates, the expression of  $3\beta$ -HSD and progesterone formation were also obtained in the brain of zebrafish (Sakamoto *et al.*, 2001a). Thus, it is now established that *de novo* steroidogenesis from cholesterol occurs in vertebrate brains (Tsutsui *et al.*, 1999).

## (3) Biosynthetic pathway of neurosteroids

In the peripheral steroidogenic glands, the production of steroid hormones requires the coordinate action of steroidogenic enzymes that start with cholesterol as the initial substrate, and catalyze a series of reactions that ultimately produce several kinds of steroids. If a variety of neurosteroids are synthesized in the brain, then each of these enzymes must be present in the vertebrate brain. To clarify the biosynthetic and metabolic pathways of neurosteroids in the brain, extensive studies with several vertebrates, especially mammals, have been carried out by many laboratories. As indicated above, the presence of cytochrome P450scc and  $3\beta$ -HSD has been well established in the vertebrate brain, whereas limited information has been available for the enzyme  $17\alpha$ -hydroxylase/ $c17,20$ -lyase (cytochrome P450<sub>17 $\alpha$ ,lyase</sub>), which converts pregnenolone to dehydroepiandrosterone, one of the most abundant neurosteroids in the brain. Therefore, Kohchi *et al.* (1998) investigated expression of the mRNAs encoding for three key steroidogenic enzymes, i.e., cytochrome P450scc,  $3\beta$ -HSD and cytochrome P450<sub>17 $\alpha$ ,lyase</sub>, using rats at different postnatal ages in order to characterize the biosynthetic pathway of abundant neurosteroids, such as  $3\beta$ -hydroxy- $\Delta^5$ -steroids and  $3$ -oxo- $\Delta^4$ -steroids, in the brain from cholesterol. The expression of P450scc mRNA occurred throughout the brain at a similar level, while  $3\beta$ -HSD mRNA expression was higher in the cerebellum and cerebrum than in other brain regions (Kohchi *et al.*, 1998). On the other hand, the P450<sub>17 $\alpha$ ,lyase</sub> mRNA was highly expressed in the mesencephalon (Kohchi *et al.*, 1998). Higher expression of the cerebellar and cerebral  $3\beta$ -HSD mRNAs was observed only during neonatal life, but the expression of both P450scc mRNA and P450<sub>17 $\alpha$ ,lyase</sub> mRNA was relatively constant during neonatal life and in adulthood (Kohchi *et al.*, 1998). These results indicate that in the postnatal rat the expression of  $3\beta$ -HSD or P450<sub>17 $\alpha$ ,lyase</sub> mRNA may be age- or region-dependent, unlike P450scc mRNA expression. Although other investigators (Guennoun *et al.*, 1995; Sanne and Krueger, 1995) also demonstrated  $3\beta$ -HSD mRNA expression in the rat brain, a pattern of age-related changes in brain  $3\beta$ -HSD mRNA expression has not previously been reported in any mammalian species. Recently, an age-dependent expression of  $3\beta$ -HSD in the cerebellum was confirmed

by both biochemical and HPLC analysis (Ukena *et al.*, 1999a). According to Ukena *et al.* (1999a), 3 $\beta$ -HSD activity in the cerebellum also increases during neonatal life. A great expression of cerebellar and cerebral 3 $\beta$ -HSD mRNAs during neonatal life suggests some functional role for the products, such as progesterone and its metabolites, of 3 $\beta$ -HSD activity.

As for P450<sub>17 $\alpha$ ,lyase</sub>, Strömstedt and Waterman (1995) also found, using RT-PCR analysis followed by Southern blots, a higher expression of the P450<sub>17 $\alpha$ ,lyase</sub> mRNA in the brain stem of postnatal rats and mice. In addition, Compagnone *et al.* (1995a) reported that rat embryonic cells expressing P450<sub>17 $\alpha$ ,lyase</sub> are located in the mesencephalic region as well as the medulla and spinal cord. Our recent studies also indicated the expression of P450<sub>17 $\alpha$ ,lyase</sub> mRNA in the quail brain (Matsunaga *et al.*, 2001). This enzyme was also highly expressed in the quail mesencephalon (Matsunaga *et al.*, 2001). Since the level of 3 $\beta$ -HSD mRNA expression is low in the mesencephalon, dehydroepiandrosterone but not progesterone, 17 $\alpha$ -hydroxy-progesterone and androstenedione may be produced as a principal neurosteroid in this brain region. Further study is needed to obtain the detailed understanding of neurosteroid production in specific brain regions and different developmental stages.

### 3. PHYSIOLOGICAL CHANGES IN NEUROSTEROIDS IN THE VERTEBRATE BRAIN

*De novo* steroidogenesis from cholesterol appears in the brain of several vertebrates, as described above. Physiological changes in neurosteroid concentrations in the brain must be taken into account when understanding the function of neurosteroids in the vertebrate brain. If neurosteroids are involved in important brain functions, we expect that they would change under different physiological conditions. To test this hypothesis, wild animal species may serve as excellent models. In contrast to the laboratory and domestic animals, the reproductive activity of most species of wild animals inhabiting the temperate and subtropical zones demonstrates a seasonal variation with a short breeding period. Such a variation is the consequence of interaction between external environmental and internal factors. Puberty in young individuals generally coincides with the onset of the breeding phase.

We examined seasonal changes in the concentrations of pregnenolone and its sulfate ester in the brain of *Rana nigromaculata*, a seasonally breeding amphibian (Takase *et al.*, 1999). Pregnenolone sulfate concentrations in the *Rana* brain were high during the active seasons, i.e., breeding phase (female) and post-breeding phase (male), and low during the quiescent season, i.e., hibernating phase (both sexes); whereas brain pregnenolone concentrations were virtually constant throughout the year (Takase *et al.*, 1999). Such a seasonal change in pregnenolone sulfate observed in the brain may be independent of peripheral steroidogenic glands, because the change in the concentration of plasma pregnenolone sulfate was significantly different from that in the brain (Takase *et al.*, 1999). We further found a seasonal

change in progesterone in the brain of newt *Cynops pyrrhogaster*. The progesterone concentration in this wild urodele brain was maximal in the breeding season in both sexes (Inai *et al.*, unpublished). A seasonal change in progesterone in the urodele brain was also independent of the plasma steroid level (Inai *et al.*, unpublished).

Recently, we have collaborated with Lea and his colleagues to analyze seasonal changes in neurosteroid concentrations using a seasonally breeding bird, the ring dove *Streptopelia risoria*. This bird also showed a seasonal change in progesterone in the brain (Clark *et al.*, 1999; Tsutsui *et al.*, 1999). Progesterone concentrations in the male dove diencephalon may increase during the brooding season, as a consequence of an increase in the 3 $\beta$ -HSD activity (Lea *et al.*, unpublished). It is considered that in the ring dove the transition from courtship to parental and associated aggressive behaviors is induced by progesterone. The expression of progesterone receptors was higher in the preoptic area in male doves during the parenting period, whereas plasma progesterone concentrations were low throughout the breeding cycle (Askew *et al.*, 1997). Accordingly, the increase of progesterone produced in the diencephalon may mediate the transition to and maintenance of parental behavior of the male birds.


### 4. NEUROSTEROIDOGENESIS IN BRAIN NEURONS

#### (1) Identification of neurosteroidogenic cells

Identification of neurosteroidogenic cells in the brain is essential to analyze the action of neurosteroids. In the first immunohistochemical description of cytochrome P450<sub>scc</sub> by Le Goascogne *et al.* (1987), an intense immunoreaction was detected in the white matter zone throughout the rat brain. The biochemical study in the rat further demonstrated that oligodendrocyte mitochondria convert cholesterol to pregnenolone (Hu *et al.*, 1987). The oligodendrocyte is a particular type of glial cell and produces the myelin of white matter. Thus, the expression and activity of P450<sub>scc</sub> in the glial cell have been established immunohistochemically and biochemically. In mammals, glial cells are considered to play a major role in neurosteroid formation and metabolism in the brain and both oligodendrocytes and astrocytes are the primary site for pregnenolone synthesis (Hu *et al.*, 1987; Jung-Testas *et al.*, 1989; Baulieu and Robel, 1990; Akwa *et al.*, 1991; Baulieu, 1991; Papadopoulos *et al.*, 1992). This is also true for the presence of P450<sub>scc</sub> in glial cells located in the telencephalic and diencephalic regions of the quail (Usui *et al.*, 1995; Tsutsui *et al.*, 1997a) and the ring dove (Lea *et al.*, 2001).

In contrast to glial cells, the concept of *de novo* neurosteroidogenesis in neurons in the brain has been uncertain. We have found the cerebellar neuron to be an active neurosteroidogenic cell, which possesses both cytochrome P450<sub>scc</sub> and 3 $\beta$ -HSD and produces pregnenolone, pregnenolone sulfate and progesterone, in several vertebrate species (Fig. 1) (Usui *et al.*, 1995; Tsutsui *et al.*, 1997a, 1997b; Ukena *et al.*, 1998; Takase *et al.*, 1999; Ukena *et al.*, 1999a).

## Neurosteroidogenesis in Purkinje Neuron



**Fig. 1. Biosynthetic pathway for neurosteroids in the cerebellar Purkinje neuron.**

StAR, P450scc and 3β-HSD are localized in this neuron. The expression of P450scc remains during neonatal development and in adulthood, indicating the constant production of pregnenolone and its sulfate. This neuron also produces highly progesterone and/or its metabolite(s), due to an increase of 3β-HSD activity, only during neonatal life. StAR: steroidogenic acute regulatory protein, P450scc: cytochrome P450 side-chain cleavage enzyme, 3β-HSD: 3β-hydroxysteroid dehydrogenase/Δ<sup>5</sup>-Δ<sup>4</sup>-isomerase

Thus, our studies provided the first evidence for the location of P450scc and 3β-HSD in the brain neuron and gave the opportunity to understand neuronal neurosteroidogenesis in the brain.

### (2) Purkinje neuron is a major site of neurosteroidogenesis

In our immunohistochemical studies of the quail brain using an antibody against P450scc, the striking observation was the distribution of immunoreactive cells in the cerebellar cortex, although other immunopositive cells were detected in telencephalic and diencephalic regions (Usui *et al.*, 1995; Tsutsui *et al.*, 1997a, 1997b). The distribution of immunoreactive cell bodies and fibers in the cerebellar cortex was coincident with the location of somata and dendrites of Purkinje cells (Usui *et al.*, 1995; Tsutsui *et al.*, 1997a, 1997b). Western immunoblot analysis confirmed the presence of P450scc in Purkinje cells (Usui *et al.*, 1995). These findings obtained in

the avian brain have provided the first evidence for the location of cytochrome P450scc in neurons in the brain, because the Purkinje cell is a typical cerebellar neuron.

Whether neurons located in the brain of other vertebrate species possess cytochrome P450scc and produce pregnenolone and its sulfate ester still remained unclear. Therefore, we investigated the presence of P450scc in the cerebellar Purkinje neuron using a mammalian species (Ukena *et al.*, 1998). Immunoreaction with P450scc was confined to the somata and dendrites of Purkinje neurons in the rat cerebellum (Ukena *et al.*, 1998). An antibody against inositol triphosphate (IP<sub>3</sub>) receptor, a marker of the Purkinje neuron, recognized P450scc-immunoreactive cerebellar cells that showed no immunoreaction with glial fibrillary acidic protein (GFAP), a specific marker of glial cells (Ukena *et al.*, 1998). In addition, the expressions of both P450scc protein and P450scc mRNA were detected in the rat cerebellum (Ukena *et al.*, 1998). Interestingly, P450scc appeared in the rat Purkinje neuron immediately after its differentiation and the expression of this enzyme persisted during neonatal development into adulthood (Ukena *et al.*, 1998). In addition to higher vertebrates, our recent studies with amphibians further identified the presence of P450scc in the cerebellar Purkinje neuron of *Xenopus laevis* and *Rana nigromaculata* (Takase *et al.*, 1999). Taken together, these findings obtained in both higher and lower vertebrates (Usui *et al.*, 1995; Tsutsui *et al.*, 1997a, 1997b; Ukena *et al.*, 1998; Takase *et al.*, 1999) indicate that Purkinje neurons possess P450scc and produce pregnenolone and its sulfate ester (Fig. 1).

Subsequently, we have extended our understanding of 3β-HSD expression in this neuron. RT-PCR and biochemical analyses showed the expression of 3β-HSD and its enzymatic activity in the cerebellum of neonatal and adult rats (Ukena *et al.*, 1999a). Employing *in situ* hybridization of 3β-HSD mRNA, the site of 3β-HSD expression was localized in Purkinje neurons and external granule cells (Ukena *et al.*, 1999a). Thus, both P450scc and 3β-HSD are expressed in Purkinje neurons (Fig. 1). The expression of 3β-HSD, however, increased during the neonatal period, unlike P450scc (Ukena *et al.*, 1999a). In contrast to a constant production of pregnenolone, Purkinje neurons produced not only progesterone but also its metabolite(s) during neonatal life (Fig. 1) (Tsutsui and Ukena, 2000; Tsutsui *et al.*, 2001). Such an age-dependent expression of 3β-HSD was confirmed by biochemical studies together with HPLC analysis, indicating an increase of progesterone formation during neonatal life (Ukena *et al.*, 1999a). Notwithstanding such a difference in the age-dependent expression, our studies have demonstrated that the Purkinje neuron is an important neurosteroidogenic cell in the vertebrate brain (Fig. 1) (for reviews, see Tsutsui and Ukena, 1999; Tsutsui *et al.*, 1999, 2000, 2001).

On the other hand, the steroidogenic acute regulatory protein (StAR) has recently been found in Purkinje neurons (Furukawa *et al.*, 1998). StAR is involved in the transport of cholesterol to the inner mitochondrial membrane, in which P450scc is localized, and thus plays a key role in steroid bio-

synthesis in the peripheral steroidogenic glands (Clark *et al.*, 1994; Stocco and Clark, 1996). StAR may also contribute to the regulation of neurosteroidogenesis in the Purkinje neuron (Fig. 1).

### (3) Other neurons

Recently, the localization of neurosteroidogenic enzymes in other brain neurons has been characterized. For instance, the expressions of P450scc, P450<sub>17 $\alpha$</sub> lyase, and P450arom were detected in the rat hippocampal neurons (Kawato *et al.*, 1999). In addition to brain neurons, P450scc was also found in neurons of the retinal ganglion, sensory neurons in the dorsal root ganglia and motor neurons in the spinal cord of the rat (Guarneri *et al.*, 1994; Compagnone *et al.*, 1995b).

## 5. BIOLOGICAL ACTIONS OF NEUROSTEROIDS PRODUCED IN PURKINJE NEURONS

### (1) Purkinje neuron serves as an excellent model for the study of neurosteroid actions


To understand neurosteroid actions in the brain, we need data on the specific synthesis in particular sites of the brain at particular times. Such informations are crucial to allow one to develop hypotheses predicting the potential roles of particular neurosteroids in the developing or adult brain. Therefore, studies for this exciting area of research should be focused on the mode of action of neurosteroids produced locally in the identified neurosteroidogenic cells underlying important brain functions. We have identified the Purkinje neuron as a major site of neurosteroidogenesis in the brain. This neuron expresses several kinds of neurosteroidogenic enzymes in a variety of vertebrates (for reviews, see Tsutsui and Ukena, 1999; Tsutsui *et al.*, 1999, 2000, 2001). In addition, the Purkinje neuron is known to play an important role in the process of memory and learning. Thus, this neuron may serve as an excellent cellular model for the study of neurosteroid actions.

### (2) Nongenomic actions as a novel neuromodulator of neurotransmission

Until recently, we believed that all steroid hormones regulate biological functions by genomic mechanisms. The genomic action of steroid hormones presumes that steroid hormones cross the plasma membrane and bind to and activate specific intracellular steroid receptors. The activated steroid receptors modulate gene transcription and protein synthesis. However, new findings have been obtained that some neurosteroids, such as pregnenolone, pregnenolone sulfate, progesterone and progesterone metabolite(s), may mediate their actions through ion-gated channel receptors rather than by genomic mechanisms. Our recent studies have focused on neurosteroid actions in the Purkinje neuron and indicated that pregnenolone sulfate contributes to important events in the cerebellum by nongenomic mechanisms.

To understand the mode of action of neurosteroids, produced in Purkinje neurons, we examined the effects of pregnenolone and its sulfate ester on synaptic currents in Purkinje

## Nongenomic Action of Pregnenolone Sulfate



**Fig. 2. Nongenomic action of pregnenolone sulfate produced in Purkinje neurons.**

Pregnenolone sulfate modulates GABAergic transmission by means of the nongenomic action on GABAergic neurons.

neurons using the rat (Tsutsui *et al.*, 1997a; Tsutsui and Ukena, 1999; Tsutsui and Ukena, 2000; Tsutsui *et al.*, 2000). Inhibitory postsynaptic currents (IPSCs) in Purkinje neurons were recorded in a cerebellar slice by the patch-clamp method. Pregnenolone sulfate increased, in a dose-related way, the frequency of IPSCs within 1 min of perfusion, indicating that this effect is unlikely to be induced *via* gene transcription. In contrast, pregnenolone had no effect on the frequency of IPSCs. The IPSCs recorded in the Purkinje neurons were completely blocked by bicuculline, a  $\gamma$ -aminobutyric acid A (GABA<sub>A</sub>) receptor antagonist, suggesting that they are mediated by GABA<sub>A</sub> receptors. Thus, pregnenolone sulfate, produced in Purkinje neurons, may modulate GABAergic transmission by nongenomic actions on GABAergic neurons rather than by genomic mechanisms (Fig. 2).

### (3) Genomic actions on neuronal growth and synaptogenesis

Purkinje neurons produce not only pregnenolone but also progesterone during the neonatal period, as the expression of 3 $\beta$ -HSD and its enzymatic activity increased in neonatal rats (Ukena *et al.*, 1999a). Recently, we also found some metabolite(s) of progesterone, such as 3 $\alpha$ ,5 $\alpha$ -tetrahydroprogesterone, in the neonatal cerebellum (Tsutsui and Ukena, 2000; Tsutsui *et al.*, 2001). It is well known that in the rat cerebellum dramatic morphological changes occur during neonatal life. According to Altman (1972a, 1972b), rat Purkinje neurons completely differentiate at 3 days of age and locate in a narrow zone between the molecular and granular layers. The external granular layer mainly develops at around 10 days

of age, followed by a migration of external granule cells into the granular layer through the Purkinje neurons, and the external granular layer disappears. The formation of the cerebellar cortex is almost complete after around 21 days of age. Thus, postnatal development in the cerebellum is dramatic during neonatal life, showing a higher expression of  $3\beta$ -HSD in the Purkinje neuron. Accordingly, progesterone and/or its metabolite(s) may be involved in the formation of the cerebellar neuronal circuit that occurs during neonatal life through promoting neuronal growth and neuronal synaptic contact by genomic actions.

To test this hypothesis, we examined the effect of progesterone, produced as a neurosteroid in the Purkinje neuron only during neonatal life, on neuronal growth in the cerebellum. Interestingly, *in vitro* studies using cultured cerebellar slices of newborn rats showed that progesterone promotes dendritic growth of the Purkinje neuron (Sakamoto *et al.*,


2001b). A similar result was obtained by *in vivo* studies. Electron microscopic analysis further revealed that progesterone induces an increase of the density of synapses on the Purkinje neuron (Sakamoto *et al.*, 2001b). Furthermore, intranuclear receptors for progesterone were expressed in the Purkinje neuron (Sakamoto *et al.*, 2001b). In contrast to progesterone, we could not detect any significant effects of  $3\alpha,5\alpha$ -tetrahydroprogesterone, a progesterone metabolite, on Purkinje development. These results indicate that progesterone promotes the dendritic growth and synaptogenesis of Purkinje neurons by genomic mechanisms (Fig. 3) (Sakamoto *et al.*, 2001b; Tsutsui *et al.*, 2001). Such an action of progesterone may contribute to the formation of the cerebellar neuronal circuit during neonatal life.

## 6. OTHER EVIDENCE FOR NEUROSTEROID ACTIONS

There is evidence indicating that in mammals neurosteroids mediate their actions through ion-gated channel receptors, such as GABA<sub>A</sub> and *N*-methyl-D-aspartate (NMDA) rather than through intracellular steroid receptors which promote the classical genomic actions (Majewska *et al.*, 1986; Majewska and Schwartz, 1987; Lambert *et al.*, 1990; Lan *et al.*, 1990; Morrow *et al.*, 1990; Puia *et al.*, 1990; Shingai *et al.*, 1991; Wu *et al.*, 1991; Majewska, 1992). For example, pregnenolone and its sulfate ester are thought to act as an agonist and antagonist of GABAergic neurotransmission (Majewska and Schwartz, 1987; Majewska *et al.*, 1988; Mienville and Vicini, 1989). In addition, pregnenolone sulfate potentiates the opening probability of the NMDA subtype of glutamate receptors in cultured neurons (Wu *et al.*, 1991; Bowlby *et al.*, 1993; Irwin *et al.*, 1992, 1994; Fahey *et al.*, 1995). Progesterone and its metabolite(s) also act through ion-gated channel receptors, such as GABA<sub>A</sub> and glycine, to modulate interneuronal communication and excitability as well as through nuclear steroid receptors (Majewska *et al.*, 1986; Lan *et al.*, 1990; Morrow *et al.*, 1990; Puia *et al.*, 1990; Shingai *et al.*, 1991; Majewska, 1992; Paul and Purdy, 1992; Valera *et al.*, 1992; Rupprecht *et al.*, 1993; Patchev *et al.*, 1994). Recently, a physiological function of pregnenolone sulfate with respect to memory has been suggested in rats (Vallée *et al.*, 1997). According to Vallée *et al.* (1997), hippocampal content of pregnenolone sulfate took part in preserving and/or enhancing cognitive abilities in aged rats, possibly *via* an interaction with central cholinergic systems. Pregnenolone sulfate may contribute to memory through mechanisms that potentiate the Ca<sup>2+</sup> conductivity of NMDA receptors (Wu *et al.*, 1991; Bowlby *et al.*, 1993; Irwin *et al.*, 1992, 1994; Fahey *et al.*, 1995).

It has also been found that progesterone promotes myelination in the peripheral nervous system (Koenig *et al.*, 1995). This result is in agreement with our hypothesis that progesterone may be involved in the formation of the cerebellar neuronal circuit in neonatal life. In addition,  $3\alpha,5\alpha$ -tetrahydroprogesterone may regulate nerve growth in rat cultured neurons (Brinton, 1994). On the other hand, progesterone and  $3\alpha,5\alpha$ -tetrahydroprogesterone may enhance

### Genomic Action of Progesterone



**Fig. 3. Genomic action of progesterone produced in Purkinje neurons.**

Progesterone acts on the Purkinje neuron through intracellular receptor-mediated mechanisms that promote dendritic growth and synaptogenesis in this neuron by genomic mechanisms. Such an action of progesterone may contribute to the formation of the cerebellar neuronal circuit during neonatal life.

sexual motivation, receptivity and proceptivity through membrane receptor-mediated mechanisms in female rats and mice (Frye and Gardiner, 1996; Frye *et al.*, 1998; Frye and Vongher, 1999).

Dehydroepiandrosterone and its sulfate ester are also abundant neurosteroids in the brain (Corpéchet *et al.*, 1981, 1983; Jo *et al.*, 1989). Recently, Compagnone and Mellon (1998) reported a stimulatory action of these neurosteroids on neuronal growth using primary cultures of mouse embryonic neocortical neurons. Dehydroepiandrosterone selectively increased the length of axons and the incidence of varicosities and basket-like process formations *in vitro*, whereas dehydroepiandrosterone sulfate selectively promoted branching and dendritic growth *in vitro* (Compagnone and Mellon, 1998). We also reported a stimulatory action of progesterone on dendritic growth and synaptogenesis in Purkinje neurons during cerebellar cortical formation (Sakamoto *et al.*, 2001b). Therefore, these neurosteroids may play an important role in cortical organization in both the cerebellum and cerebrum during development.

## 7. CONCLUSIONS AND FUTURE DIRECTIONS

*De novo* steroidogenesis from cholesterol is now established in the vertebrate brain. Steroids synthesized in the brain as well as other nervous system are called neurosteroids. Neurosteroidogenic enzymes are expressed in both neurons and glial cells. The Purkinje cell, a cerebellar neuron, is considered to play a major role in neurosteroid formation and metabolism in the brain. This neuron possesses the neurosteroidogenic enzymes cytochrome P450<sub>scc</sub> and 3 $\beta$ -HSD and produces pregnenolone, pregnenolone sulfate and progesterone from cholesterol. Cytochrome P450<sub>scc</sub> appears in the Purkinje neuron immediately after its differentiation. The expression of P450<sub>scc</sub> remains during neonatal development and in adulthood, indicating the constant production of pregnenolone and its sulfate. This neuron also produces significant amounts of progesterone, as a product of an increase of 3 $\beta$ -HSD activity, only in a limited neonatal period. Pregnenolone sulfate modulates GABAergic transmission by nongenomic actions on GABAergic neurons rather than genomic mechanisms. On the other hand, progesterone is involved in the promotion of dendritic growth and synaptogenesis of the Purkinje neuron by genomic mechanisms. This action of progesterone may contribute to the formation of the cerebellar neuronal circuit during neonatal life. These serve as an excellent model for the study of physiological roles of neurosteroids in the cerebellum, because Purkinje neurons play an important role in the process of memory and learning. Therefore, future attention should be focused on behavioral studies using neurosteroidogenic enzyme-knockout animals and electrophysiological studies on the occurrence of long-term potentiation (LTP) and/or long-term depression (LTD). In other brain regions, future study is also required to clarify the mode of action of neurosteroids produced locally in the identified cells underlying important brain functions.

## ACKNOWLEDGMENTS

This work was supported in part by Grants-in-Aid for Scientific Research from the Ministry of Education, Science, and Culture, Japan (11170237, 11354010, 12440233, 12894021 and 13210101 to K.T.) and by the International Joint Research Project of the British Council. We are grateful to Drs. K. Ukena, H. Sakamoto, Y. Honda, Y. Inai, M. Matsunaga, Y. Furukawa, M. Takase, C. Kohchi, T. Yamazaki and S. Kominami (Hiroshima University, Higashi-Hiroshima, Japan) and Dr. R. W. Lea (University of Central Lancashire, Preston, U.K.) for their collaboration.

## REFERENCES

- Adkins E, Adler N (1972) Hormonal control of behavior in the Japanese quail. *J Comp Physiol Psychol* 81: 27–36
- Akwa Y, Young J, Kabbadj K, Sancho MJ, Zucman D, Vourc'h C, Jung-Testas I, Hu ZY, Le Goascogne C, Jo DH, Corpéchet C, Simon P, Baulieu EE, Robel P (1991) Neurosteroids: Biosynthesis, metabolism and function of pregnenolone and dehydroepiandrosterone in the brain. *J Steroid Biochem Mol Biol* 40: 71–81
- Akwa Y, Sananès N, Gouézou M, Robel P, Baulieu EE, Le Goascogne C (1993) Astrocytes and neurosteroids: metabolism of pregnenolone and dehydroepiandrosterone. Regulation by cell density. *J Cell Biol* 121: 135–143
- Altman J (1972a) Postnatal development of the cerebellar cortex in the rat. I. The external germinal layer and the transitional molecular layer. *J Comp Neurol* 145: 353–397
- Altman J (1972b) Postnatal development of the cerebellar cortex in the rat. II. Phases in the maturation of Purkinje cells and of the molecular layer. *J Comp Neurol* 145: 399–463
- Arnold A (1975) The effects of castration and androgen replacement on song, courtship, and aggression in zebra finches (*Poephila guttata*). *J Exp Zool* 191: 309–326
- Arnold A, Nottebohm F, Pfaff DW (1976) Hormone-concentrating cells in vocal control and other areas of the brain of the zebra finch. *J Comp Neurol* 165: 487–512
- Askew JA, Georgiou GC, Sharp PJ, Lea RW (1997) Localization of progesterone receptor in brain and pituitary of the ring dove: influence of breeding cycle and estrogen. *Horm Behav* 32: 105–113
- Balthazart J (1983) Hormonal correlates of behavior. In: Eds by DS Farner, KC Parker, *Avian Biology*. Academic Press, New York, pp 221–365
- Baulieu EE (1991) Neurosteroids: a new function in the brain. *Biol Cell* 71: 3–10
- Baulieu EE (1997) Neurosteroids: of the nervous system, by the nervous system, for the nervous system (review). *Rec Progr Hormone Res* 52:1–32
- Baulieu EE, Robel P (1990) Neurosteroids: A new brain function? *J Steroid Biochem Mol Biol* 37: 395–403
- Bowlby MR (1993) Pregnenolone sulfate potentiation of *N*-methyl-D-aspartate receptor channels in hippocampal neurons. *Mol Pharmacol* 43: 813–819
- Brinton RD (1994) The neurosteroid 3 $\alpha$ -hydroxy-5 $\alpha$ -pregnan-20-one induces cytoarchitectural regression in cultured fetal hippocampal neurons. *J Neurosci* 14: 2763–2774
- Clark BJ, Wells J, King SR, Stocco DM (1994) The purification, cloning, and expression of a novel luteinizing hormone-induced mitochondrial protein in MA-10 mouse Leydig tumor cells. Characterization of the steroidogenic acute regulatory protein (StAR). *J Biol Chem* 269: 28314–28322
- Clark JA, Tsutsui K, Ukena K, Lea RW (1999) Changes in central progesterone during the reproductive cycle of the ring dove (*Streptopelia risoria*). 15th National Meeting of the British Neuro-

- science Association, Harrogate, UK, p 15 (abstract)
- Compagnone NA, Bulfone A, Rubenstein JLR, Mellon SH (1995a) Steroidogenic enzyme P450c17 is expressed in the embryonic central nervous system. *Endocrinology* 136: 5212–5223
- Compagnone NA, Bulfone A, Rubenstein JLR, Mellon SH (1995b) Expression of the steroidogenic enzyme P450scc in the central and peripheral nervous systems during rodent embryogenesis. *Endocrinology* 136: 2689–2696
- Compagnone NA, Mellon SH (1998) Dehydroepiandrosterone: a potential signalling molecule for neocortical organization during development. *Proc Natl Acad Sci USA* 95: 4678–4683
- Compagnone NA, Mellon SH (2000) Neurosteroids: biosynthesis and function of these novel neuromodulators (review). *Frontiers Neuroendocrinol* 21: 1–56
- Corpéchet C, Robel P, Axelson M, Sjövall J, Baulieu EE (1981) Characterization and measurement of dehydroepiandrosterone sulfate in rat brain. *Proc Natl Acad Sci USA* 78: 4704–4707
- Corpéchet C, Synguelakis M, Talha S, Axelson M, Sjövall J, Vihko R, Baulieu EE, Robel P, (1983) Pregnenolone and its sulfate ester in rat brain. *Brain Res* 270: 119–125
- Dupont E, Simard J, Luu-The V, Labrie F, Pelletier G (1994) Localization of 3 $\beta$ -hydroxysteroid dehydrogenase in rat brain as studied by *in situ* hybridization. *Mol Cell Neurosci* 5: 119–123
- Fahey JM, Lindquist DG, Pritchard GA, Miller LG (1995) Pregnenolone sulfate potentiation of NMDA-mediated increases in intracellular calcium in cultured chick cortical neurons. *Brain Res* 669: 183–188
- Frye CA, Bayon LE, Pursnani NK, Purdy RH (1998) The neurosteroids, progesterone and 3 $\alpha$ , 5 $\alpha$ -THP, enhance sexual motivation, receptivity, and proceptivity in female rats. *Brain Res* 808: 72–83
- Frye CA, Gardiner SG (1996) Progestins can have a membrane-mediated action in rat midbrain for facilitation of sexual receptivity. *Horm Behav* 30: 682–691
- Frye CA, Vongher JM (1999) Progesterone has rapid and membrane effects in the facilitation of female mouse sexual behavior. *Brain Res* 815: 259–269
- Furukawa A, Miyatake A, Ohnishi T, Ichikawa Y (1998) Steroidogenic acute regulatory protein (StAR) transcripts constitutively expressed in the adult rat central nervous system: Colocalization of StAR, cytochrome P-450scc (CYP 11A1), and 3 $\beta$ -hydroxysteroid dehydrogenase in the rat brain. *J Neurochem* 71: 2231–2238
- Fuxe K, Gustafsson JA, Wetterberg L (1981) *Steroid Hormone Regulation of the Brain*. Pergamon Press, Oxford
- Guarneri P, Guarneri R, Cascia C, Pavaasant P, Piccoli F, Papadopoulos V (1994) Neurosteroidogenesis in rat retinas. *J Neurochem* 63: 86–96
- Guennoun R, Fiddes RJ, Gouézou M, Lombès M, Baulieu EE (1995) A key enzyme in the biosynthesis of neurosteroids, 3 $\beta$ -hydroxysteroid dehydrogenase/ $\Delta^5$ - $\Delta^4$ -isomerase (3 $\beta$ -HSD), is expressed in rat brain. *Mol Brain Res* 30: 287–300
- Hu ZY, Bourreau E, Jung-Testas I, Robel P, Baulieu EE (1987) Neurosteroids: oligodendrocyte mitochondria convert cholesterol to pregnenolone. *Proc Natl Acad Sci USA* 84: 8215–8219
- Irwin RP, Maragakis NJ, Rogawski MA, Purdy RH, Farb DH, Paul SM (1992) Pregnenolone sulfate augments NMDA receptor mediated increases in intracellular Ca<sup>2+</sup> in cultured rat hippocampal neurons. *Neurosci Lett* 141: 30–34
- Irwin RP, Lin SZ, Long RT, Paul SM (1994) *N*-methyl-D-aspartate induces a rapid, reversible, and calcium-dependent intracellular acidosis in cultured fetal rat hippocampal neurons. *J Neurosci* 14: 1352–1357
- Ishii S, Tsutsui K (1982) Hormonal control of aggressive behavior in male Japanese quail. In: Eds by CG Scanes, MA Ottinger, AD Kenney, J Balthazart, J Cronshaw, JC Jones, *Aspects of Avian Endocrinology: Practical and Theoretical Implications*. Texas Tech Press, Texas, pp 125–131
- Iwahashi K, Ozaki HS, Tsubaki M, Ohnishi J, Takeuchi Y, Ichikawa Y (1990) Studies of the immunohistochemical and biochemical localization of the cytochrome P-450scc-linked monooxygenase system in the adult rat brain. *Biochim Biophys Acta* 1035: 182–189
- Jo DH, Abdallah MA, Young J, Baulieu EE, Robel P (1989) Pregnenolone, dehydroepiandrosterone, and their sulfate and fatty acid esters in the rat brain. *Steroids* 54: 287–297
- Jung-Testas I, Hu ZY, Baulieu EE, Robel P (1989) Neurosteroids: biosynthesis of pregnenolone and progesterone in primary cultures of rat glial cells. *Endocrinology* 125: 2083–2091
- Kabbadj K, El-Etr M, Baulieu EE, Robel P (1993) Pregnenolone metabolism in rodent embryonic neurons and astrocytes. *Glia* 7: 170–175
- Kawato S, Kimoto T, Ohta Y, Tsurugizawa T, Makino J, Hojo Y, Takahashi T (1999) Localization and activities of neurosteroidogenic systems in the hippocampal neurons. In: Eds by M Okamoto, Y Ishimura, H Nawata, *Molecular Steroidogenesis*. Universal Academy Press, Tokyo, Japan, pp 385–388
- Koenig HL, Schumacher M, Ferzaz B, Do Thi AN, Ressouches A, Guennoun R, Jung-Testas I, Robel P, Akwa Y, Baulieu EE (1995) Progesterone synthesis and myelin formation by Schwann cells. *Science* 268: 1500–1503
- Kohchi C, Ukena K, Tsutsui K (1998) Age- and region-specific expressions of the messenger RNAs encoding for steroidogenic enzymes P450scc, P450c17 and 3 $\beta$ -HSD in the postnatal rat brain. *Brain Res* 801: 233–238
- Korsia S, Bottjer SW (1989) Developmental changes in the cellular composition of a brain nucleus involved with song learning in zebra finches. *Neuron* 3: 451–460
- Lambert JJ, Peters JA, Sturgess NC, Hales TG (1990) Steroid modulation of the GABA<sub>A</sub> receptor complex: electrophysiological studies. *Ciba Found Symp* 153: 56–71
- Lan NC, Chen JS, Belelli D, Pritchett DB, Seeburg PH, Gee KW (1990) A steroid recognition site is functionally coupled to an expressed GABA<sub>A</sub>-benzodiazepine receptor. *Eur J Pharmacol* 188: 403–406
- Lanthier A, Patwardhan VV (1986) Sex steroids and 5-en-3 $\beta$ -hydroxysteroids in specific regions of the human brain and cranial nerves. *J Steroid Biochem* 25: 445–449
- Le Goascogne C, Robel P, Gouézou M, Sananès N, Baulieu EE, Waterman M (1987) Neurosteroids: cytochrome P-450scc in rat brain. *Science* 237: 1212–1215
- Lea RW, Clark JA, Tsutsui K (2001) Changes in central steroid receptor expression, steroid synthesis and dopaminergic activity related to the reproductive cycle of the ring dove (review). *Microsc Res Tech (MRT)*, in press
- Majewska MD (1992) Neurosteroids: endogenous bimodal modulators of the GABA<sub>A</sub> receptor: mechanism of action and physiological significance. *Prog Neurobiol* 38: 379–395
- Majewska MD, Schwartz RD (1987) Pregnenolone-sulfate: an endogenous antagonist of the  $\gamma$ -aminobutyric acid receptor complex in the brain? *Brain Res* 404: 355–360
- Majewska MD, Harrison NL, Schwartz RD, Barker JL, Paul SM (1986) Steroid hormone metabolites are barbiturate-like modulators of the GABA receptor. *Science* 232: 1004–1007
- Majewska MD, Mienville JM, Vicini S (1988) Neurosteroid pregnenolone sulfate antagonizes electrophysiological responses to GABA in neurons. *Neurosci Lett* 90: 279–284
- Mason JI (1993) The 3 $\beta$ -hydroxysteroid dehydrogenase gene family of enzymes. *Trends Endocrinol Metab* 4: 199–203
- Mathur C, Prasad VVK, Raju VS, Welch M, Lieberman S (1993) Steroids and their conjugates in the mammalian brain. *Proc Natl Acad Sci USA* 90: 85–88
- Matsunaga M, Ukena K, Tsutsui K (2001) Expression and localization of cytochrome P450 17 $\alpha$ -hydroxylase/c17, 20-lyase in the

- avian brain. *Brain Res* 899: 112–122
- McEwen BS (1991) Steroid hormones are multifunctional messengers in the brain. *Trends Endocrinol Metab* 2: 62–67
- Mellon SH, Deschepper CF (1993) Neurosteroid biosynthesis: genes for adrenal steroidogenic enzymes are expressed in the brain. *Brain Res* 629: 283–292
- Mensah-Nyagan AG, Feuilloley M, Dupont E, Do-Rego JL, Leboulenger F, Pelletier G, Vaudry H (1994) Immunocytochemical localization and biological activity of  $3\beta$ -hydroxysteroid dehydrogenase in the central nervous system of the frog. *J Neurosci* 14: 7306–7318
- Mienville JM, Vicini S (1989) Pregnenolone sulfate antagonizes GABA<sub>A</sub> receptor-mediated currents via a reduction of channel opening frequency. *Brain Res* 489: 190–194
- Morrow AL, Pace JR, Purdy RH, Paul SM (1990) Characterization of steroid interactions with  $\gamma$ -aminobutyric acid receptor-gated chloride ion channels: evidence for multiple steroid recognition sites. *Mol Pharmacol* 37: 263–270
- Papadopoulos V, Guarneri P, Krueger KE, Guidotti A, Costa E (1992) Pregnenolone biosynthesis in C6-2B glioma cell mitochondria: Regulation by a mitochondrial diazepam binding inhibitor receptor. *Proc Natl Acad Sci USA* 89: 5113–5117
- Patchev VK, Shoab M, Holsboer F, Almeida OFX (1994) The neurosteroid tetrahydroprogesterone counteracts corticotropin-releasing hormone-induced anxiety and alters the release and gene expression of corticotropin-releasing hormone in the rat hypothalamus. *Neuroscience* 62: 265–271
- Paul SM, Purdy RH (1992) Neuroactive steroids. *FASEB J* 6: 2311–2322
- Pignataro L, Lerner AAC, Baranao JL, de Plazas SF (1998) Biosynthesis of progesterone derived neurosteroids by developing avian CNS: *in vitro* effects on the GABA<sub>A</sub> receptor complex. *Int J Dev Neurosci* 16: 433–441
- Pröve VE (1978) Quantitative untersuchungen zu wechselbeziehungen zwischen balzaktivität und testosteronitern bei männlichen Zebrafinken (*Taeniopygia guttata*). *J Tierpsychology* 48: 47–67
- Puia G, Santi MR, Vicini S, Pritchett DB, Purdy RH, Paul SM, Seeburg PH, Costa E (1990) Neurosteroids act on recombinant human GABA<sub>A</sub> receptors. *Neuron* 4: 759–765
- Robel P, Baulieu EE (1985) Neuro-steroids,  $3\beta$ -hydroxy- $\Delta^5$ -derivatives in the rodent brain. *Neurochem Int* 7: 953–958
- Robel P, Corpéchet C, Clarke C, Groyer A, Synguelakis M, Vourc'h C, Baulieu EE (1986) Neuro-steroids:  $3\beta$ -hydroxy- $\Delta^5$ -derivatives in the rat brain. In: Eds by G Fink, AJ Harmar, KW McKerns, *Neuroendocrine Molecular Biology*. Plenum Press, New York, pp 367–377
- Robel P, Bourreau E, Corpéchet C, Dang DC, Halberg F, Clarke C, Haug M, Schlegel ML, Synguelakis M, Vourc'h C, Baulieu EE (1987) Neuro-steroids:  $3\beta$ -hydroxy- $\Delta^5$ -derivatives in rat and monkey brain. *J Steroid Biochem* 27: 649–655
- Rupprecht R, Reul JM, Trapp T, van Steensel B, Wetzel C, Damm K, Ziegglänsberger W, Holsboer F (1993) Progesterone receptor-mediated effects of neuroactive steroids. *Neuron* 11: 523–530
- Sakamoto H, Ukena K, Tsutsui K (2001a) Activity and localization of  $3\beta$ -hydroxysteroid dehydrogenase/ $\Delta^5$ - $\Delta^4$ -isomerase in the zebrafish central nervous system. *J Comp Neurol*, in press
- Sakamoto H, Ukena K, Tsutsui K (2001b) Effects of progesterone synthesized *de novo* in the developing Purkinje cell on its dendritic growth and synaptogenesis. *J Neurosci*, 21: 6221–6232
- Sanne JL, Krueger KE (1995) Expression of cytochrome P450 side-chain cleavage enzyme and  $3\beta$ -hydroxysteroid dehydrogenase in the rat central nervous system: a study by polymerase chain reaction and *in situ* hybridization. *J Neurochem* 65: 528–536
- Shingai R, Sutherland ML, Barnard EA (1991) Effects of subunit types of the cloned GABA<sub>A</sub> receptor on the response to a neurosteroid. *Eur J Pharmacol* 206: 77–80
- Stocco DM, Clark BJ (1996) Regulation of the acute production of steroids in steroidogenic cells. *Endocr Rev* 17: 221–244
- Strömstedt M, Waterman MR (1995) Messenger RNAs encoding steroidogenic enzymes are expressed in rodent brain. *Mol Brain Res* 34: 75–88
- Takase M, Ukena K, Yamazaki T, Kominami S, Tsutsui K (1999) Pregnenolone, pregnenolone sulfate and cytochrome P450 side-chain cleavage enzyme in the amphibian brain and their seasonal changes. *Endocrinology* 140: 1936–1944
- Tsutsui K, Ishii S (1981) Effects of sex steroids on aggressive behavior of adult male Japanese quail. *Gen Comp Endocrinol* 44: 480–486
- Tsutsui K, Yamazaki T (1995) Avian neurosteroids. I. Pregnenolone biosynthesis in the quail brain. *Brain Res* 678: 1–9
- Tsutsui K, Yamazaki T, Usui M, Furukawa Y, Ukena K, Kohchi C, Kominami S (1997a) P450scc activity in the brain. In: Eds by S Harvey, RJ Etches, *Perspectives in Avian Endocrinology*. Journal of Endocrinol Ltd, Bristol, pp 427–436
- Tsutsui K, Usui M, Yamazaki T, Ukena K, Kominami S (1997b) Neurosteroids in the avian brain. In: Ed by SK Maitra, *Frontiers in Environmental and Metabolic Endocrinology*. Burdwan Press, Burdwan, pp 151–159
- Tsutsui K, Ukena K (1999) Neurosteroids in the cerebellar Purkinje neuron and their actions (review). *Int J Mol Med* 4: 49–56
- Tsutsui K, Ukena K, Takase M, Kohchi C, Lea RW (1999) Neurosteroid biosynthesis in vertebrate brains (review). *Comp Biochem Physiol C* 124: 121–129
- Tsutsui K, Ukena K (2000) Neuronal neurosteroidogenesis in the cerebellum. In: Eds by M Okamoto, Y Ishimura, H Nawata, *Molecular Steroidogenesis*. Universal Academy Press, Tokyo, pp 397–400
- Tsutsui K, Ukena K, Usui M, Sakamoto H, Takase M (2000) Novel brain function: biosynthesis and actions of neurosteroids in neurons (review). *Neurosci Res* 36: 261–273
- Tsutsui K, Schlinger BA (2001) Steroidogenesis in the avian brain. In: Eds by A Dawson, CM Chaturvedi, *Avian Endocrinology*. Narosa Publishing House, New Delhi, pp 59–77
- Tsutsui K, Ukena K, Sakamoto H (2001) Novel cerebellar function: Neurosteroids in the Purkinje neurons and their genomic and nongenomic actions. In: Eds by R J Handa, S Hayashi, *Neuroplasticity, Development, and Steroid Hormone Action*. CRC Press, Boca Raton, USA, pp 101–116
- Ukena K, Usui M, Kohchi C, Tsutsui K (1998) Cytochrome P450 side-chain cleavage enzyme in the cerebellar Purkinje neuron and its neonatal change in rats. *Endocrinology* 139: 137–147
- Ukena K, Kohchi C, Tsutsui K (1999a) Expression and activity of  $3\beta$ -hydroxysteroid dehydrogenase/ $\Delta^5$ - $\Delta^4$ -isomerase in the rat Purkinje neuron during neonatal life. *Endocrinology* 140: 805–813
- Ukena K, Honda Y, Inai Y, Kohchi C, Lea RW, Tsutsui K (1999b) Expression and activity of  $3\beta$ -hydroxysteroid dehydrogenase/ $\Delta^5$ - $\Delta^4$ -isomerase in different regions of the avian brain. *Brain Res* 818: 536–542
- Ukena K, Honda Y, Lea RW, Tsutsui K (2001) Developmental changes in progesterone biosynthesis and metabolism in the quail brain. *Brain Res* 898: 190–194
- Usui M, Yamazaki T, Kominami S, Tsutsui K (1995) Avian neurosteroids. II. Localization of a cytochrome P450scc-like substance in the quail brain. *Brain Res* 678: 10–20
- Valera S, Ballivet M, Bertrand D (1992) Progesterone modulates a neuronal nicotinic acetylcholine receptor. *Proc Natl Acad Sci USA* 89: 9949–9953
- Vallée M, Mayo W, Darnaudéry M, Corpéchet C, Young J, Koehl M, Moal ML, Baulieu EE, Robel P, Simon H (1997) Neurosteroids: deficient cognitive performance in aged rats depends on low pregnenolone sulfate levels in the hippocampus. *Proc Natl Acad Sci USA* 94: 14865–14870

- Vanson A, Arnold AP, Schlinger BA (1996)  $3\beta$ -Hydroxysteroid dehydrogenase/isomerase and aromatase activity in primary cultures of developing zebra finch telencephalon: dehydroepiandrosterone as substrate for synthesis of androstenedione and estrogens. *Gen Comp Endocrinol* 102: 342–350
- Watson JT, Adkins-Regan E (1989) Neuroanatomical localization of sex steroid-concentrating cells in the Japanese quail (*Coturnix japonica*): autoradiography with [ $^3$ H] testosterone, [ $^3$ H] estradiol, and [ $^3$ H] dihydrotestosterone. *Neuroendocrinology* 49: 51–64
- Weidenfeld J, Siegel RA, Chowdhury I (1980) In vitro conversion of pregnenolone to progesterone by discrete brain areas of the male rat. *J Steroid Biochem* 13: 961–963
- Wingfield JC, Marler P (1988) Endocrine basis of communication in reproduction and aggression. In: Eds by E Knobil, J Neill, *The Physiology of Reproduction*. Raven Press, New York, pp 1647–1677
- Wingfield JC, Farner DS (1993) Endocrinology of reproduction in wild species. In: Eds by DS Farner, JR King, KC Parkes, *Avian Biology*. Academic Press, New York, pp 163–327
- Wu F-S, Gibbs TT, Farb DH (1991) Pregnenolone sulfate: a positive allosteric modulator at the *N*-methyl-D-aspartate receptor. *Mol Pharmacol* 40: 333–336
- Zwain IH, Yen SSC (1999) Dehydroepiandrosterone: biosynthesis and metabolism in the brain. *Endocrinology* 140: 880–887

(Received May 18, 2001 / Invited Review)