

# Sex Inversion of Sexually Immature Honeycomb Grouper (Epinephelus merra) by Aromatase Inhibitor

Authors: Bhandari, Ramji Kumar, Komuro, Hiroki, Higa, Mikihiko, and Nakamura, Masaru

Source: Zoological Science, 21(3): 305-310

Published By: Zoological Society of Japan

URL: https://doi.org/10.2108/zsj.21.305

BioOne Complete (complete.BioOne.org) is a full-text database of 200 subscribed and open-access titles in the biological, ecological, and environmental sciences published by nonprofit societies, associations, museums, institutions, and presses.

Your use of this PDF, the BioOne Complete website, and all posted and associated content indicates your acceptance of BioOne's Terms of Use, available at <u>www.bioone.org/terms-of-use</u>.

Usage of BioOne Complete content is strictly limited to personal, educational, and non - commercial use. Commercial inquiries or rights and permissions requests should be directed to the individual publisher as copyright holder.

BioOne sees sustainable scholarly publishing as an inherently collaborative enterprise connecting authors, nonprofit publishers, academic institutions, research libraries, and research funders in the common goal of maximizing access to critical research.

# Sex Inversion of Sexually Immature Honeycomb Grouper (*Epinephelus merra*) by Aromatase Inhibitor

Ramji Kumar Bhandari, Hiroki Komuro, Mikihiko Higa and Masaru Nakamura\*

Sesoko Station, Tropical Biosphere Research Center, University of the Ryukyus, Sesoko, Motobu, Okinawa 905-0227, Japan

**ABSTRACT**—Previous studies have shown that estrogen plays an important role in sex change of protogynous honeycomb grouper, and that the treatments with aromatase inhibitor (AI) cause estrogen depletion and complete sex inversion of pre-spawning females into functional males. In the present study, we examined whether AI causes sex inversion of sexually immature females. Female honeycomb groupers were implanted with various doses of Fadrozole (0, 100, 500 and 1000 µg/fish) in the non-breeding season, and resultant changes in the gonadal structures and the plasma levels of sex steroid hormones (estradiol-17β, E2; testosterone, T; 11-ketotestosterone, 11-KT) were examined three months after implantation. Vehicle-implanted groups did not change sex, while 100 and 500 µg Al-implanted groups had turned into transitionals with intersex gonad. In contrast, the highest dose receiving group exhibited both transitional and male phases. Transitional phase gonad had attretic occytes and spermatogenic germ cells at the late stages of spermatogenesis, while male phase testis contained spermatozoa accumulated in the seminiferous tubules. All males released sperm upon slight pressure on the abdomen. In the Al-implanted fish, plasma levels of E2 decreased in a dose-dependent manner, while the levels of 11-KT were high in the highest dose receiving group. Present results suggest that estrogen plays an important role in sex change of protogynous honeycomb grouper, and that treatments with AI potentially inhibits endogenous E2 production in vivo, causing oocyte degeneration and subsequently the sex inversion from female to male. The Fadrozole could be an important tool for manipulating the sex of hermaphrodite fishes.

Key words: sex change, hermaphrodite fish, honeycomb grouper, aromatase inhibitor, estrogen

# INTRODUCTION

Teleost fishes exhibit a wide variety of reproductive tactics, including adult sex change. Gonochoristic fishes, after gonadal differentiation at the early stages of life cycle, do not reverse sex, while hermaphrodite fishes change their sex in response to social stimuli, age, growth rate and seasons. The honeycomb grouper (Epinephelus merra) is a protogynous hermaphrodite fish. Due to its relatively smaller size among groupers, this species serves as a good model for grouper research. The sex change occurs at an adult stage, particularly in the non-breeding season. In the field populations of honeycomb groupers, the onset of sex change was associated with the oocyte degeneration and low levels of serum estradiol-17 $\beta$ , which was subsequently accompanied by gradual increase in the 11-KT levels and spermatogonia proliferation toward the center of ovigerous lamellae (Bhandari et al., 2003). In the protogynous wrasse

\* Corresponding author: Tel. +81-980-47-6072; FAX. +81-980-47-6072. E-mail: masaru@lab.u-ryukyu.ac.jp (*Thalassoma duperrey*), the onset of sex change was associated with oocyte degeneration and significant drop in E2 levels (Nakamura *et al.*, 1989). These results suggest that the endogenous level of estrogens, when drop below the threshold levels *in vivo*, could be stimulatory to sex change in the field populations of protogynous hermaphrodite fishes.

In fish, estrogen selectively acts as an ovarian inducer (Nakamura *et al.*, in press). The estrogen biosynthesis is mediated by steroidogenic enzyme cytochrome P450 aromatase (arom), which is believed to play an important role in ovarian differentiation of various fish species (Nakamura *et al.*, 1998, 2000; Guigen *et al.*, 1999; Kitano *et al.*, 2000; D'Cotta *et al.*, 2001). The aromatase inhibitor (AI) treatments around the period of gonadal sex differentiation have caused estrogen depletion and various degrees of masculinization, including complete and functional sex reversal in some species (Piferrer *et al.*, 1997; Kitano *et al.*, 2000; Kwon *et al.*, 2000, 2002; Afonso *et al.*, 2001). In hermaphrodite fishes, AI treatments of adult females have resulted in either partial (Kroon and Liley, 2000) or complete sex reversal (Higa *et al.*, in press; Bhandari *et al.*, in press). However,

very little is known about effects of AI on sex inversion of sexually immature hermaphrodite fishes. In the present study, we collected female honeycomb groupers in the nonbreeding season and implanted them with various doses of non-steroidal AI (Fadrozole) to elucidate the role of estrogen in sex change of protogynous hermaphrodite fishes.

# MATERIALS AND METHODS

Wild honeycomb groupers of average total length of 18 to 20 cm were collected from fishermen of Nakijin village, northern Okinawa. We have found that female honeycomb groupers have total length below 20 cm (Bhandari et al., 2003). The spawning season starts during the full moon of May and lasts in July (Soyano K, personal communication). Generally, October through January are considered as non-breeding period. In the present study, sexually immature females were selected to implant with AI (a gift from Norvartis Pharma, Tokyo, Japan) and as controls. Altogether, 42 fish were implanted, of which 10 fish were sacrificed for initial control, and remaining 32 fish were divided into four groups with eight fish in each treatment. On January 6, 2003, control fish received implantation of vehicle (cocoa butter only) and fish in the treatment groups received 100, 500 and 1000 µg AI (dissolved in the cocoa butter), respectively into the body cavity. Later, they were reared in the polyethylene cages (80 cm×80 cm×80 cm) placed over 5 tons cemented tanks with flow-through sea water. All fish were sampled in April 6, 2003. After anaesthetization with phenoxyethanol (0.05%), the total length and wet body weight were measured, blood was collected from caudal vein into a 1 ml heparinized syringe, and fish were sacrificed for the collection of pituitary and gonads. Gonads were weighed to determine gonadosomatic index (GSI) and a small piece of gonad from each individual was preserved in the Bouin's solution for the histological analysis. The collected blood was centrifuged at 3000 rpm for 15 minutes and obtained plasma was stored at -30°C until analysis. The plasma levels of E2, testosterone (T) and 11-ketotestosterone (11-KT) were determined by enzyme-linked immunosorbent assay (ELISA) according to the routine procedure. Later, preserved gonadal tissues were embedded in the paraffin, sectioned at 7  $\mu$ m, and stained with Delafield's hematoxylin and eosin. The prepared slides were examined under light microscope for histological changes.

### RESULTS

#### GSI and Histological changes in gonads

At the end of experiment, the GSI of control fish was significantly higher than that in the remaining groups (Fig. 1A). The gonadal restructuring in the AI-implanted fish were categorized into four stages: female, early transition, late transition and male, according to Bhandari et al (2003). At the start of experiment, the ovary of initial control females had oocytes at the perinucleolar stages, except some at the pre-vitellogenic stages (Fig. 2A). At the end of experiment, the ovary of control females had vitellogenic oocytes at the volk globule stage (Fig. 2B). A fish from control group had ovotestis at the early transition stage. About 75% of 100  $\mu$ g Al-implanted fish had gonads at early transition stage, where perinucleolar stage oocytes were undergoing atresia, and spermatogonia were sparsely proliferating at the periphery of the ovigerous lamella, indicating sex change (Fig. 2C). In the intersexual gonads of 500 µg Al-implanted fish,



Treatment

**Fig. 1.** Changes in gonadosomatic index (GSI), plasma levels of estradiol-17 $\beta$  (E2), testosterone (T) and 11-ketotestosterone (11-KT) in the honeycomb grouper implanted with various doses of Fadrozole. Numbers in the bars indicate total number of fish sampled. \* indicates statistically significant difference at *P*<0.05 from the connected means.

spermatogenic germ cells were extensively proliferating, and previtellogenic oocytes were undergoing atresia, indicating the sex change was progressing (Fig. 2D). Out of eight 1000  $\mu$ g Al-implanted fish, the gonads of two fish were at the early transition stage, three were at the late transition stage, and three had already transformed into testis. Those testes had various stages of spermatogenic germ cells, including an accumulation of sperm in the seminiferous tubules (Fig. 2E). The degree of sex change in each treatment is shown in the Fig. 3. The gonads of majority of 100



**Fig. 2.** Light micrograph of gonadal structure of honeycomb grouper before and after implantation with Fadrozole: (A) initial control (B) vehicle, (C) 100 μg, (D) 500 μg, and (E) 1000 μg Fadrozole/fish. Abbreviations: AO, atretic oocyte; PO, primary oocyte; VO, vitellogenic oocyte; SG, spermatogonia; SC, spermatocyte; ST, spermatid; and SZ, spermatozoa. Scale bar=100 μm.

and 500  $\mu$ g Al-implanted fish were at early or late transition stages, whereas those in 1000  $\mu$ g Al-implanted fish were at the late transition stage or fully transformed into testis.

# Changes in the plasma levels of sex steroid hormones

The plasma levels of sex steroid hormones are shown in the Fig. 1. Compared to initial controls, the plasma levels of E2 significantly increased in the control fish (Fig. 1), whereas in the Al-implanted fish, the levels decreased in a dose-dependent manner. The plasma levels of T did not change, while the levels of 11-KT significantly increased in the 1000 µg Al-implanted fish.

# DISCUSSION

In the present study, we investigated whether AI causes sex inversion of sexually immature female honeycomb grouper. The low doses of AI-receiving fish had shown the progress in sex change, whereas the majority of highest dose receiving fish had either late transitional gonad or transformed into testis, suggesting AI has dose-dependent effects on sex inversion of honeycomb grouper.



**Fig. 3.** Sex ratio of honeycomb grouper implanted with various doses of Al. Abbreviations: F, Female phase; ET, early transition phase; LT, late transition phase; M, Male phase (according to Bhandari *et al.*, 2003). Numbers in bars indicate the total number of fish. The testes of LT and Male phase fish contained spermatogenic germ cells at various stages of development, including spermatozoa.

The steroidogenic enzyme aromatase that converts androgens into estrogens plays an important role in ovarian differentiation (Nakamura et al., 1998; Strüssmann and Nakamura, 2002; Guigen et al., 1999, D'Cotta et al., 2001). The AI inhibition of aromatase activity is believed to cause oocyte degeneration, and induce sex reversal in various fish species. The efficacy of AI has been tested in a variety of gonochoristic species, where treatments around the period of gonadal sex differentiation have caused complete sex reversal from females to functional males (Piferrer et al., 1997; Nakamura et al., 1999; Kitano et al., 2000, Afonso et al., 2001), while its effects on sex reversal of adult gonochorists are largely unknown. In contrast, in the hermaphrodite fishes, AI treatments have shown variable results in terms of its effectiveness. For the honeycomb grouper, Al-implantation caused complete sex inversion of pre-spawning females to functional males in two and half months (Bhandari et al., in press), while the oral administration caused complete sex inversion of wrasse in six weeks (Higa et al., unpublished), whereas the treatments for the same period caused formation of transitional gonads in the blackeye goby (Kroon and Liley, 2000). Together with previous results, present results support the hypothesis that AI treatments cause sex inversion of protogynous hermaphrodite fishes; however, the effectiveness probably depends on species, the extent of gonadal development, and the dose of AI.

In the wild honeycomb grouper, low levels of serum E2 and the degeneration of perinucleolar stage oocytes were associated with the onset of sex change (Bhandari *et al.*, 2003), while in the wrasse, *Thalassoma duperrey*, a rapid drop in E2 levels, and degeneration of pre-vitellogenic oocytes were observed at the onset of sex change (Nakamura *et al.*, 1989). In the present study, the plasma E2 levels were decreased dose- dependently in the AI-implanted fish. Treatments with AI caused a significant drop in the E2 production by ovarian fragments of sexually mature female honeycomb grouper *in vitro*, while AI-implantation of sexually maturing females caused E2 depletion and led to complete and functional sex inversion *in vivo* (Bhandari *et al.*, in press), whereas E2 compensation preserved the female sex (Bhandari *et al.*, unpublished). Together, present results clearly suggest that a significant drop in endogenous estrogen levels is necessary to trigger the female to male sex change in fish.

In fish, estrogen is believed to act as ovarian inducer. In the gonochoristic Nile tilapia, AI treatments, after the period of gonadal sex differentiation, caused sex reversal of genetically controlled all females into functional males (Nakamura et al., in press), whereas E2 prevented AIinduced sex reversal (Bhandari et al., unpublished). Other than fish, AI inhibition of estrogen synthesis has caused sex reversal of variety of animals, such as Newt (Chardard and Dournon, 1999), reptiles (Dorizzi et al., 1994; Wibbles and Crews, 1994), and chicken (Elbrecht and Smith, 1992; Abinawanto et al., 1996; Burke and Henry, 1999). Moreover, the role of estrogen on female sex differentiation in mammals has been explained by creating estrogen receptor knock-out mice. The ovaries of estrogen receptor  $\alpha$  and  $\beta$ knock-out mice exhibited follicle transdifferentiation to structures resembling seminiferous tubules of the testis, including Sertoli-like cells and expression of mullerian inhibiting substance, sulfated glycoprotein-2, and Sox9 (Couse et al., 1999). These results strongly support the present results that estrogen is inevitable for maintaining female sex in the wide range of vertebrate classes, where depending on the stages of sexual development, any alteration of estrogen balance could result in sex reversal.

In the present study, plasma 11-KT levels were increased in the sex-changed males, while no change was observed for T levels. Although, the levels of  $11\beta$ -hydroxytestosterone (11β-HT) could not be measured in plasma, it is likely that testosterone might have converted into 11β-HT as T to 11B-HT conversion is active during sex change in some hermaphrodite fishes (Idler et al., 1976). In the wild population of honeycomb grouper, 11-KT levels tended to increase with the developmental stages of testicular tissues (Bhandari et al., 2003), while in the wrasse the 11-KT levels increased sharply after the onset of sex change (Nakamura et al., 1989). In the field populations of red grouper, 11-KT levels were high in the individuals possessing inter-sex and male gonads (Johnson et al., 1998). In addition, treatments with 11-KT or methyltestosterone or the mixture of various androgens have resulted in complete sex change of several protogynous hermaphrodite fishes (Tan-Fermin et al., 1996; Kroon and Liley, 2000, Tanaka et al., 2000; Yeh et al., 2003a, b; Higa et al., in press). These results clearly indicate that androgens also play important role in the sex change of hermaphrodite fishes, but yet, there still remain some questions: whether increase in 11-KT levels exert stimulatory effects on spermatogonia proliferation in the ovary or stimulate spermatogenesis after its complete transformation into testicular tissues. Nevertheless, future development of protocol for in vitro sex change would be helpful for elucidating

the cellular mechanism of germ cell proliferation during sex reversal and sex change in fish.

In conclusion, present results support the hypothesis that E2 plays an important role in maintaining female sex of hermaphrodite fishes, and that the AI inhibition of E2 synthesis causes oocyte degeneration and subsequently the spermatogonia proliferation in the ovary leading to functional sex inversion. Moreover, Fadrozole being a potent aromatase inhibitor (Steele *et al.*, 1987; Smith, 1999), could be an important tool for the sex inversion of hermaphrodite fishes and also in the studies to further investigate what determines the fate of a gonocyte to develop into either oogonia or spermatogonia during the onset of sex change or reversal. In the aquaculture point of view, the AI-induction of sex change would be an important tool for the timely production of male brood stocks for breeding purposes and to enhance grouper aquaculture.

# ACKNOWLEDGEMENT

Authors thank Mr. Y. Henna and Mr. Y. Kojima of Sesoko Station for assistance with fish transportation and sampling. The part of this study is the contribution from Japan Science and Technology (JST) Foundation, and Sesoko Station, University of the Ryukyus, Japan. The authers would like to thank Novartis Pharma, Japan for aromatase inhibitor gift.

# REFERENCES

- Abinawanto, Shimada K, Yoshida K, Saito N (1996) Effects of aromatase inhibitor on sex differentiation and levels of P45017 and P450arom messenger ribonucleic acid of gonads in chicken embryos. Gen Comp Endocrinol 102: 241–246
- Afonso LOB, Wassermann GJ, Terezinha de Oliveira R (2001) Sex reversal in Nile tilapia (*Oreochromis niloticus*) using a nonsteroidal aromatase inhibitor. J Exp Zool 290: 177–181
- Bhandari RK, Higa M, Nakamura S, Nakamura M (2003) Aromatase inhibitor induces complete sex change in protogynous honeycomb grouper (*Epinephelus merra*). Mol Reprod Dev 67: 303– 307
- Bhandari RK, Komuro H, Higa M, Nakamura S, Nakamura M (2003) Gonadal restructuring and correlative sex steroid profiles during protogynous sex change in the honeycomb grouper (*Epi-nephelus merra*). Zool Sci 20: 1399–1404
- Burke WH, Henry MH (1999) Gonadal development and growth of chickens and turkeys hatched from eggs injected with an aromatase inhibitor. Poultry Sci 78: 1019–1033
- Chardard D, Dournon C (1999) Sex reversal by aromatase inhibitor treatment in the newt (*Pleurodeles waltl*). J Exp Zool 283: 43– 50
- Couse JF, Curtis Hewitt S, Bunch DO, Sar M, Walker VR, Davis BJ, Korach KS (1999) Postnatal sex reversal of the ovaries in mice lacking estrogen receptor  $\alpha$  and  $\beta$ . Science 286: 2328–2331
- D'Cotta H, Fostier A, Guiguen Y, Govoroun MS, Baroiller JF (2001) Aromatase plays a key role during normal and temperatureinduced sex differentiation of tilapia (*Oreochromis niloticus*). Mol Reprod Dev 59: 265–276
- Dorizzi M, Richard-Mercier N, Desvages G, Pieau C (1994) Masculinization of gonads by aromatase inhibitors in a turtle with temperature-dependent sex determination. Differentiation 58: 1–8
- Elbrecht A, Smith RG (1992) Aromatase activity and sex determination in chickens. Science 255: 467–470

- Guiguen Y, Baroiller JF, Ricordel MJ, Iseki K, Mcmeel OM, Martin, SA, Fostier A (1999) Involvement of estrogens in the process of sex differentiation in two fish species: the rainbow trout (*Oncorhynchus mykiss*) and a tilapia (*Oreochromis niloticus*). Mol Reprod Dev 54: 154–162
- Higa M, Ogasawara K, Sakaguchi A, Nagahama Y, Nakamura M (2003) Roles of steroid hormones in sex change of wrasse. Fish Physiol Biochem (in press)
- Idler DR, Reinboth R, Walsh JM, Truscott B (1976) A comparison of 11-hydroxytestosterone and 11-ketotestosterone in blood of ambisexual and gonochoristic teleosts. Gen Comp Endocrinol 30: 517–521
- Johnson AK, Thomas P, Wilson JR RR (1998) Seasonal cycles of gonadal development and plasma sex steroid levels in *Epinephelus morio*, a protogynous grouper in the eastern Gulf of Mexico. J Fish Biol 52: 502–518
- Kitano T, Takamune K, Nagahama Y, Abe SI (2000) Aromatase inhibitor and 17alpha-methyltestosterone cause sex-reversal from genetical females to phenotypic males and suppression of P450 aromatase gene expression in Japanese flounder (*Paralichthys olivaceus*). Mol Reprod Dev 56: 1–5
- Kroon FJ, Liley NR (2000) The role of steroid hormones in protogynous sex change in the Blackeye goby, *Coryphopterus nicholsii* (Teleostei: Gobiidae). Gen Comp Endocrinol 118: 273–283
- Kwon JY, Haghpanah V, Kogson-Hurtado LM, McAndrew BJ, Penman DJ (2000) Masculinization of genetic female Nile tilapia (*Oreochromis niloticus*) by dietary administration of an aromatase inhibitor during sexual differentiation. J Exp Zool 287: 46–53
- Kwon JY, McAndrew BJ, Penman DJ (2002) Treatment with an aromatase inhibitor suppresses high-temperature feminization of genetic male (YY) Nile tilapia. J Fish Biol 60: 625–636
- Nakamura M, Bhandari RK, Higa M (2003) The role estrogens play during sex differentiation and sex change in fish. Fish Physiol Biochem (review, in press)
- Nakamura M, Hourigan TF, Yamauchi K, Nagahama Y, Grau GE (1989) Histological and ultrastructural evidence for the role of gonadal steroid hormones in sex change in the protogynous wrasse (*Thalassoma duperrey*). Env Biol Fish 24: 117–136
- Nakamura M, Kobayashi T, Chang XT, Nagahama Y (1998) Gonadal sex differentiation in teleost fish. J Exp Zool 281: 362– 372
- Nakamura M, Kobayashi T, Yoshiura Y, Nagahama Y (2000) Role of endogenous steroid hormones on gonadal sex differentiation in fish. In "Proc. 6th Int symp reprod physiol fish", Ed by B Noberg, OS Kjesbu, GL Taranger, E Andersson, SO Stefansson, Institute for Marine Research and University of Bergen, pp 247–249
- Piferrer F, Zanuy S, Carrillo M, Solar II, Devlin RH, Donaldson EM (1994) Brief treatment with an aromatase inhibitor during sex differentiation causes chromosomally female salmon to develop as normal, functional males. J Exp Zool 270: 255–262
- Smith IE (1999) Aromatase inhibitors: a dose-response effect? Endocrine-related Cancer 6: 245–249
- Steele RE, Mellor LB, Sawyer WK, Wasvary JM, Browne LJ (1987) In vitro and in vivo studies demonstrating potent and selective oestrogen inhibition with the nonsteroidal aromatase inhibitor CGS 16949A. Steroids 50: 147–161
- Strüssmann CA, Nakamura M (2002) Morphology, endocrinology, and environmental modulation of gonadal sex differentiation in teleost fishes. Fish Physiol Biochem 26: 13–29
- Tanaka H, Tsuchihashi Y, Kuromiya Y (2000) Induction of sex reversal in the sevenband grouper, *Epinephelus septemfasciatus*. In "Proc. 6th International symposium on the reproductive physiology of fish". Ed by B Norberg, OS Kjesbu, GL Taranger, E Andersson, SO Stefansson, Institute for Marine Research and University of Bergen, pp 423

- Tan-Fermin JD, Garcia-Luis MB, Castillo AR Jr (1994) Induction of sex inversion in juvenile grouper, (*Epinephelus suilus*, Valenciennes) by injections of 17-alpha-methyltestosterone. Jpn J Ichthyol 40: 413–420
- Wibbels T, Crews D (1994) Putative aromatase inhibitor induces male sex determination in a female unisexual lizard and in a turtle with temperature dependent sex determination. J Endocrinol 141: 295–299
- Yeh SL, Kuo CM, Ting YY, Chang CF (2003a) Androgens stimulate sex change in protogynous grouper (*Epinephelus coioides*): spawning performance in sex-changed males. Comp Biochem Physiol 135C: 375–382
- Yeh SL, Kuo CM, Ting YY, Chang CF (2003b) The effects of exogenous androgens on ovarian development and sex change in female orange-spotted protogynous grouper, (*Epinephelus coioides*). Aquaculture 218: 729–739

(Received September 12, 2003 / Accepted November 22, 2003)