



Perymenium vandeenderorum (Asteraceae), a New Species From Sonora, Mexico

Author: Turner, Billie L.

Source: *Lundellia*, 20(1) : 1-3

Published By: The Plant Resources Center, The University of Texas at Austin

URL: <https://doi.org/10.25224/1097-993X-20.1.1>

BioOne Complete (complete.BioOne.org) is a full-text database of 200 subscribed and open-access titles in the biological, ecological, and environmental sciences published by nonprofit societies, associations, museums, institutions, and presses.

Your use of this PDF, the BioOne Complete website, and all posted and associated content indicates your acceptance of BioOne's Terms of Use, available at www.bioone.org/terms-of-use.

Usage of BioOne Complete content is strictly limited to personal, educational, and non - commercial use. Commercial inquiries or rights and permissions requests should be directed to the individual publisher as copyright holder.

BioOne sees sustainable scholarly publishing as an inherently collaborative enterprise connecting authors, nonprofit publishers, academic institutions, research libraries, and research funders in the common goal of maximizing access to critical research.

PERYMENIUM VANDEVENDERORUM (ASTERACEAE), A NEW SPECIES FROM SONORA, MEXICO

Billie L. Turner

Plant Resources Center, The University of Texas at Austin, Main Bldg. 127, 110 Inner Campus Drive, Stop F0404,
Austin, Texas 78712 Austin, TX 78712
Email: billie.turner@austin.utexas.edu

Abstract: A novel species of *Perymenium* from Sonora, Mexico is described. It appears to be closest to *P. nesomii* from southwestern Chihuahua, with which it is compared. A map showing distribution of the two taxa is provided.

Keywords: Asteraceae, *Perymenium*, *P. nesomii*, Mexico, Sonora, Chihuahua

As noted by Panero (2007), *Perymenium* Schrad. is a rather large genus with 50 or more species, most of them located in Mexico, these last treated by Turner (2014), in which five species were recorded for the state of Sonora, or along its boundaries.

***Perymenium vandevenderorum* B.L. Turner, sp. nov.**

PERENNIAL HERBS to 1 m high (roots not examined). MID-STEMS pubescent with upwardly appressed minute hairs. LEAVES ovate-lanceolate, opposite throughout, mostly 6-10 cm long, 1.5-2.5 cm wide, pubescent like the stems, 3-nervate from near the base, their margins entire or nearly so; petioles 5-10 mm long. CAPITULESCENCE a terminal array of 10-30 heads, 4-8 cm across. HEADS ca. 8 mm high, 4 mm wide, their ultimate peduncles 1-3 cm long. INVOLUCRAL BRACTS broadly ovate, 3-4 seriate, rounded to obtuse at apex, the upper margins sparsely ciliate. RECEPTACLE CONVEX. PALES linear, 4-7 mm long. Ray florets, 8-11, pistillate, fertile; ligules yellow, 6-8 mm long, 1-2 mm wide. DISC FLORETS, 15-20 per head; corollas yellow, glabrous, 4-5 mm long; tubes ca. 1.5 mm long; throats ca. 3 mm long; lobes ca. 0.5 mm long. ACHENES of ray and disc florets similar, 2-sided, bearing a pappus of short, readily deciduous, bristles 1-3 mm long.

TYPE: MEXICO. SONORA: Mpio. Bacanora, Restaurant el Torreoncito, 11.8 km (by

air) WSW of Bacanora, 28°56'59"N, 109°31'01"W, 949 m. "Foothills thorn scrub. Common herbaceous perennial on roadside," 30 Jul 2014, T.R. Van Devender et al. 2014-365 (HOLOTYPE: TEX). Fig. 1.

ADDITIONAL SPECIMENS EXAMINED: MEXICO. SONORA: Las Tierras de Jimenez, Sierra de Murrieta, 28°33'33"N, 109°31'33"W, "Open woodland on limestone ridge. Occasional herbaceous perennial" 1 Aug 2014, T.R. Van Devender et al. 2014-331 (TEX).

This novelty is readily recognized by its robust habit and entire leaves, the latter strikingly different from closely related taxa, namely *Perymenium nesomii* B.L. Turner of nearby southwestern Chihuahua that appears to be its nearest relative; indeed, it will key to, or near, that species in my treatment of the genus for Mexico, the taxon endemic to the Cascada de Basaseachic of SW Chihuahua. *Perymenium vandevenderorum* is readily identified by its larger, more branched habit, entire leaves (vs. serrate) and somewhat larger heads. Distribution of the two species is shown in Fig. 2.

The collectors of the plants concerned describe the material as perennial herbs, but they appear to be more nearly like subshrubs, only the upper parts of the collections mounted, as noted above and shown on the type itself.

It's a pleasure to name the novelty for my close friend, Thomas R. van Devender and his wife Ana Lila Reina, who collected both of the above cited specimens.



FIG. 1. The type specimen of *Perymenium vandevenderorum* B.L. Turner located at TEX.




FIG. 2. Distribution of *Perymenium vandevederorum* and *P. nesomii* in Mexico.

ACKNOWLEDGMENTS

Specimens cited are on file at LL-TEX. My editorial assistant, Jana Kos, provided helpful input.

LITERATURE CITED

- Panero, J. L.** 2007. *Perymenium*, p. 456 in Kubitzki, K. (ed.) *The Families and Genera of Vascular Plants*. Vol. 8 *Eudicots: Asterales* J.W. Kadereit and C. Jeffrey (eds). New York, Springer Verlag.
- Turner, B. L.** 2014. *Perymenium* pp. 3–31 in *The Comps of Mexico*. *Phytologia Memoirs* 21.