

## **Genetic evidence of unisexual reproduction in the Moroccan hexaploid barbel *Labeobarbus fritschi***

Authors: Berrebi, Patrick, Retif, Xavier, and Bouhbouh, Saadia

Source: Folia Zoologica, 62(4) : 257-263

Published By: Institute of Vertebrate Biology, Czech Academy of Sciences

URL: <https://doi.org/10.25225/fozo.v62.i4.a2.2013>

---

BioOne Complete ([complete.BioOne.org](https://complete.BioOne.org)) is a full-text database of 200 subscribed and open-access titles in the biological, ecological, and environmental sciences published by nonprofit societies, associations, museums, institutions, and presses.

Your use of this PDF, the BioOne Complete website, and all posted and associated content indicates your acceptance of BioOne's Terms of Use, available at [www.bioone.org/terms-of-use](https://www.bioone.org/terms-of-use).

Usage of BioOne Complete content is strictly limited to personal, educational, and non - commercial use. Commercial inquiries or rights and permissions requests should be directed to the individual publisher as copyright holder.

---

BioOne sees sustainable scholarly publishing as an inherently collaborative enterprise connecting authors, nonprofit publishers, academic institutions, research libraries, and research funders in the common goal of maximizing access to critical research.

# Genetic evidence of unisexual reproduction in the Moroccan hexaploid barbel *Labeobarbus fritschi*

Patrick BERREBI<sup>1\*</sup>, Xavier RETIF<sup>1</sup> and Saadia BOUHBOUH<sup>2</sup>

<sup>1</sup> Institut des Sciences de l'Évolution, UMR 5554 UM2-CNRS-IRD, Université Montpellier II, CC 065, Place E. Bataillon, 34095 Montpellier Cedex 5, France; e-mail: patrick.berrebi@univ-montp2.fr

<sup>2</sup> Université Sidi Mohamed Ben Abdellah, Faculté des Sciences Dhar El Mehraz, Fès, Morocco

Received 7 September 2013; Accepted 15 November 2013

**Abstract.** The hexaploid barbel population of the species *Labeobarbus fritschi* which lives in the Allal El Fassi dam reservoir in the north of Morocco showed an absence of males among a sample of 597 individuals analyzed in a first survey and among the 39 fish analyzed in this study. A possible explanation is that this population may be composed of females only and that they reproduce either by gynogenesis or hybridogenesis, two processes triggered by hybridization with a sexually incompatible species (possibly the sympatric barbel *Luciobarbus setivimensis* in this case). We used molecular markers to collect any information that could explain the phenomenon. To do this, numerous nuclear loci were screened (more than 50) in search of polymorphism. Genetic diversity was low as a possible consequence of clonal reproduction: only three allozymic and two intronic loci presented a sufficient level of polymorphism to be exploited. The existence of several copies of the same multilocus genotypes (1/3 of the fish) – with some showing a significant probability of not being derived from sexual reproduction – provided evidence of unisexual reproduction. Deviations from panmixia as well as linkage equilibrium were also observed. Although preliminary, the data supports the hypothesis of the unisexual reproduction of females by way of gynogenesis rather than hybridogenesis.

**Key words:** Morocco, allozymes, introns, gynogenesis, hybridogenesis

## Introduction

The genus *Barbus sensu lato* (Cuvier & Cloquet, 1816) belongs to the family Cyprinidae, composed of more than 800 species (Howes 1987). This genus, in its *sensu lato* (*s.l.*) acceptation, gathers taxa from Asia, Europe and Africa; it is only diagnosed as possessing two pairs of barbels, with several exceptions (one pair or no barbels).

In this enormous aggregation (Myers 1960), recent taxonomic improvements based on molecular/cytogenetic data made it possible to identify large ploidy lineages already considered as distinct genera (Berrebi 1995, Berrebi et al. 1996, Skelton 2002, Berrebi & Tsigenopoulos 2002, 2003, Berrebi et al. 2013).

*Labeobarbus fritschi* (Günther, 1874) is a hexaploid barbel (El Gharbi 1994, Machordom & Doadrio 2001). Since 1996, the subgenus *Labeobarbus* has become a genus. It is widespread in Africa (Golubtsov & Krysanov 1993, Berrebi et al. 1996, Skelton 2002), particularly in Morocco (Machordom & Doadrio 2001) and in the Middle East (Durand et al. 2002, Tsigenopoulos et al. 2002). *L. fritschi* lives, among others, in the Allal El Fassi dam lake (Morocco), in

the River Zerouka, a tributary of the River Sebu, north of Rabat (Bouhbouh 2002) where it accounts for most of the ichthyofauna with another tetraploid barbel, *Luciobarbus setivimensis* (Valencia, 1842).

Bouhbouh (2002) reported that samples of *L. fritschi* caught in this dam lake included only females (597 females from 597 fish sampled). The absence of males may be explained by two assumptions: (i) the males of this species were absent from the fishing sites since the barbel has a tendency to migrate, (ii) it is a monosex population consisting only of females and the reproductive process behind this phenomenon must be elucidated.

Teleost fish can develop different modes of reproduction including various forms of unisexual reproductions. Several mechanisms were described as self-fecundation in some hermaphrodites, or gynogenesis and hybridogenesis in unisexual females of some populations (Chevassus 1998, Baroiller & Guiguen 2001, Gui & Zhou 2010, Collares-Pereira et al. 2013). Based on the genetic analysis of barbel using nuclear molecular markers, the objective of this study is to understand the relevance of each one of the two

hypotheses: differential migrations between the sexes, or a unisexual population reproducing either by gynogenesis or hybridogenesis.

Material and Methods

Thirty-nine individuals of the species *L. fritschi* were collected in February 2002 using gillnets in the Allal El Fassi reservoir. The fish were frozen at -20 °C immediately after capture and transported to the laboratory in Montpellier, France. The sex of each of the 39 fish was verified by observation of gonads with a microscope after dissection. The molecular markers analyzed were of two types: allozymes and introns (EPIC – Exon Primed Intron-Crossing).

Allozymes

The enzyme extraction was performed on muscle and liver tissue of each fish according to the protocols described by Pasteur et al. (1987). The desired enzyme loci were highlighted by colour-adapted protocols by Berrebi et al. (1993, 1995) and Tsigenopoulos et al. (1999). Ten enzyme systems (IDH, LDH, PGI, MDH, 6PGD, AAT, GDA, AK and PGM) and the soluble muscle proteins (PT), corresponding to 25 presumptive loci, were tested. Table 1a presents details of the retained loci and buffers used (nomenclature according to Shaklee et al. 1990).

Introns

DNA was extracted according to the classical phenol/chloroform protocol (Sambrook et al. 1989). EPIC-

PCR used primers complementary to exon sequences flanking introns. Exon sequences are mostly conserved among the species, providing universal priming zones which hold a practical interest (Atarhouch et al. 2003). A total of 27 primer pairs were tested in this study. Table 1 lists the characteristics of the retained loci for the analysis. PCR reactions were performed using an Eppendorf Mastercycler in a volume of 10 µL for each individual, containing 1 µL of 10X buffer (Promega), 2.5 mM MgCl 2, 0.2 mM of each dNTP (Invitrogen), 0.5 µM of each of the two primers, one of which is labeled with a fluorochrome (Cy3, Cy5 or Fuorescein, MWG-Biotech), 0.3 U of *Taq* polymerase (Sigma) and 1 µL of DNA template at about 150 µg/mL. The PCR program included an initial denaturation at 94 °C for 3 min, then 35 cycles consisting of denaturation for 1 min, binding of the primers (annealing) at the temperature specific to each locus (see Table 1b) for 1 min and extension at 72 °C for 1 min 20 sec; the program ended with a final extension at 72 °C for 10 min. The PCR products were then subjected to electrophoresis on acrylamide denaturing (urea) 8 % gel. This gel was read by a Hitachi FMBIO II scanner. The FMBIO Analysis 8.0 software can accurately measure allele sizes in number of base pairs (bp).

Statistical analyses

The data matrix was statistically processed involving three steps:  
1) Calculation of the parameters commonly used in population genetics (allele frequencies, heterozygosity).

Table 1. Marker descriptions.  
a) Allozymic loci: tissue of extraction, electrophoretic buffers and staining solution compositions.

Enzymatic loci				
Locus	Tissue	Gem buffer	Electrode buffer	Staining
PGI-2*	liver	TCB 8.7	Borate 8.2	5 mL tris Hcl 0.2M ph 8
PGI-3*	liver	TCB 8.7	Borate 8.2	1 mL Mgcl2 0.5M
				1 mL NAD
				0.5 mL NADP
				10 mg fructose 6 P
PGI-4*	muscle	Tris HCl	Borate 8.2	0.5 mL PMS
				0.5 mL MTT
				1 mL NBT
				6 µL G6PDH
				Agar

b) Intronic loci: primer sequences (F = forward, R = reverse), original publication and specific annealing temperature.

Intronic loci			
Locus	Primers	References	Tm
RP	F = TGGCCTCTTCCTTGGCCGTC R = AACTCGTCTGGCTTTTCGCC	Chow & Takeyama 1998	54 °C
Mlc2a	F = CAGCTGTCCCATGGTGGCCA R = GGCCGGTACCTCCAATGTGTTT	Atarhouch et al. 2003	55 °C

2) Calculation of Wright's fixation indices such as  $F_{is}$  (checking panmixia) by the estimator  $f$  of Weir & Cockerham (1984) and linkage disequilibrium (Black & Krafur 1985). The statistical significance of these parameters was estimated by resampling permutations tests (5000 permutations). The GENETIX software (Belkhir et al. 1998) was used for these first two steps. 3) The clonality of genotypes was tested using the GENECLONE software (Arnaud-Haond & Belkhir 2007). This software established the probability of existence for each multilocus genotype taking into account the  $F_{is}$  of the population analyzed, i.e. a deviation from panmixia as described by Young et al. (2002). Next, the probability that  $n$  repetitions of the same genotype can occur in the population by  $n$  independent events of sexual reproduction was estimated (psex; Tibayrenc et al. 1990, Parks & Werth 1993, Arnaud-

matrix which was subjected to the statistical analyzes described above.

1) Table 2 gives the allelic frequencies of enzyme loci and introns, as well as the partial estimated (considering only the polymorphic loci) and observed heterozygosities.

2) The estimate of  $F_{is}$  produces a negative value ( $-0.328$ ) that is highly significant ( $p < 0.001$ ). This negative value indicates an excess of heterozygotes (see Table 2) and thus a deviation from Hardy-Weinberg equilibrium (HWE) in the barbel population. Similarly, the test of linkage disequilibrium between the four loci shows that the pairs of loci PGI-2\*/Mlc2a and PGI-4\*/RP were significantly correlated ( $p < 0.001$ ).

3) The test of clonality was applied to the data set. A group of multi-locus genotypes (those of individuals

**Table 2.** Allelic frequencies and heterozygosities of the five loci analyzed. Ho = observed heterozygosity, He = expected heterozygosity.

Alleles	Intronic loci						Enzymatic loci						Heterozygosities	
	RP		Mlc2a		PGI-2*		PGI-3*				PGI-4*		Ho	He
Allelic frequencies	100	110	100	110	080	100	090	100	110	120	080	100	0.51	0.38

Haond et al. 2005). If  $psex < 0.05$ , repeated genotypes have a significant probability of being propagated by unisexual reproduction.

## Results

Dissection and observation under a binocular microscope of a gonad for each of the 39 fish revealed that they were all females, the gonads containing oocytes that were easily recognizable, at different stages of maturity between individuals.

Among the 10 enzymatic systems tested, 5 showed interpretable zymograms: AK, PGI, MDH, PGM and PT despite hexaploidy, for a total of 8 loci. The loci AK\*, MDH-1\* and MDH-2\* were monomorphic in the sample and uninformative. The loci PT\* and PGM\* showed a presence/absence polymorphism not used for our demonstration.

Only three loci of the PGI system (Phospho-Gluco-isomerase) were polymorphic and usable for our purpose: the loci PGI-2\* and PGI-3\* migrated to the cathode while PGI-4\* migrated to the anode.

Among the twenty-seven intronic systems tested, only two gave usable results with a measurable length polymorphism: the PR and Mlc2a systems. Each had only a single polymorphic locus, in both cases with two alleles. Other systems tested were either unreadable due to overlapping bands (12) or monomorphic (13). Enzymes and introns data were gathered in a single

h4, h14, h25 and h36, see Appendix) showed a significant probability ( $psex < 0.03$ ) of not having been generated by sexual reproduction and therefore generated by clonal reproduction.

## Discussion

### *Polyploidy compensates low polymorphism*

Polyploidy is considered to confer more tolerance to ecological variations because the duplication of their genes provides metabolic flexibility similar to a permanent high genomic heterozygosity (Uyeno & Smith 1972, Otto & Whitton 2000, Jiang et al. 2013). The *Labeobarbus fritschii* Moroccan populations showed low polymorphism and thus low heterozygosity; however their hexaploidy may compensate this lack. Technically speaking, hexaploidy provoked a very low level of usable markers in our study due to complex multiple bands patterns at all nuclear markers.

El Gharbi (1994) analyzed a population of *L. fritschii* from the River Zerouka in her work on the phylogeny of the genus *Barbus s.l.* complex in North Africa. Her sample came from a scientific station located downstream of the Allal El Fassi dam lake analyzed in this study. It showed that the downstream population was in HWE. This author does not seem to have estimated the sex ratio of this river population.

In this 1994 study, only two allozyme loci were polymorphic (AK-2\* and PGI-4\*) out of the 15

analyzed in total. In this study, five loci were evidenced as polymorphic among 8 allozymic and 15 intronic interpretable markers. The polymorphic loci analyzed in the two studies were similarly scarce (respectively 2/15 and 5/23).

#### *Sex migration dimorphism hypothesis*

In agreement with the observations of Bouhbouh (2002), examination of the gonads of 39 barbel fish of 2002 from the dam lake revealed that they were all females. How can this population reproduce with only females in the lake? The two relevant hypotheses that may explain this phenomenon are: migration of male fish with seasonal meetings with females, or unisexual reproduction.

The absence of males as remarked by Bouhbouh (2002) was observed after many gillnet fishing expeditions and during all seasons between 2000 and 2001. Our own observations on a sample of 2002 are consistent with this result. If the meeting of the two sexes was seasonal, multiple fishing opportunities should have revealed some males. It is therefore very likely that no male is living in the lake. This simple observation leads us to consider unisexual reproduction as the best hypothesis.

#### *Unisexual reproduction hypothesis*

The data analysis resulted in some important features pointing in favour of the unisexual reproduction hypothesis. Deviations from HWE are important population features. Firstly, the Fis estimate showed that there was a very significant excess of heterozygotes while most panmictic deviations described in natural populations showed a deficit in heterozygotes, often due to a Wahlund effect (Aurelle et al. 2002). Secondly, we found two pairs of loci (PGI-2\*/Mlc2a and PGI-4\*/RP) that were statistically associated ( $p < 0.001$ ). Analysis of genotypes showed that two thirds of the specimens possessed the same genotype at the five loci studied. Given the low polymorphism at our disposal, it is questionable whether such sets of identical genotypes can be observed in fish populations that reproduce sexually. The test of clonality (GENECLONE) showed that at least one multilocus genotype, repeated four times (four individuals, n°4, 14, 25 and 36, bearing the letter h in the Appendix), had a significant probability of not being propagated by sexual reproduction processes. All these results lead us to the unisexual reproduction hypothesis.

#### *Gynogenesis*

Unisexual female populations are known in fish, often after episodes of hybridization between species of

the same genus (see review in Chevassus 1998). The cases of the genera *Poecilia* and *Poeciliopsis* illustrate two classical patterns of female monosex populations. *Poecilia formosa* lives in unisexual female populations, reproducing by gynogenesis (premeiotic endomitosis) with the production of diploid eggs. The sperm of another species is only used to trigger the embryonic development without participating in the genome of the future embryo as evidenced in several studies (Hubbs & Hubbs 1932, Turner et al. 1980, Castelli 1994, Ryan et al. 1996, Yang et al. 2001).

Reproduction by gynogenesis at the most maintains the initial heterozygosity (Schultz 1977, Bulger & Schultz 1979) but its reduction is expected. This mode of unisexual reproduction does not meet the Mendelian laws and causes major deviations from HWE and statistical linkages among loci.

In the hybridogenesis case of the *Poeciliopsis* complex (Moore & McKay 1971, Chevassus 1998), unisexual female populations of *P. monacha* coexist with bisexual sympatric populations (*P. lucida* and *P. occidentalis*). Thus, two forms of unisexual diploid female populations result from the hybridization between females of *P. monacha* and males of each of the two sympatric bisexual species. In these diploid females, meiosis is not recombinant but reductional and hemiclinal (hybridogenesis), allowing the genome of the mother to be transmitted without recombination: the eggs contain only the maternal genome (*P. monacha*), the paternal genome brought by the males of *P. lucida* or *P. occidentalis* participating in the somatic line only (Schultz 1980). Hemiclinal hybridogenesis maintains their hybrid status (Schultz 1977, Bulger & Schultz 1979), leading to highly polymorphic populations.

In this *Labeobarbus* monosex population, the triggering role of sympatric and even syntopic males of *Luciobarbus setivimensis* can be expected since favorable spawning areas in the dam lake are probably limited, leading to accidental gametes contacts and since *Luciobarbus* genus is a distant lineage (*Labeobarbus* is an African hexaploid lineage while *Luciobarbus* is a North African and European tetraploid one). According to the limited information produced here for the *Labeobarbus* case, gynogenesis is more probable than hybridogenesis which should follow fecundation between very different lineages (tetraploid and hexaploid of a distinct origin, Tsigenopoulos et al. 2002).

Limited nuclear genetic data is mainly due to the hexaploid nature of the *Labeobarbus* genus (Tsigenopoulos et al. 2002). Demonstrating the



mechanism of unisexual reproduction occurring in the lake requires much more information than that produced for this preliminary study. Microsatellites and introns are traditionally the main markers used in this kind of research. Despite the high number of intron system runs (15), only two proved to be interpretable for this study. It is probable that microsatellites would add a more few markers, but genome scan methods

will clearly be necessary (genome scan techniques as AFLP) to disentangle the process.

## Acknowledgements

The authors would like to thank Jean-François Baroiller for his help in elucidating the reproductive process, Sophie Arnaud-Haond for her contribution to statistical questions and Edwige Quillet for a time-consuming critical reading of a preliminary version of this manuscript.

## Literature

- Arnaud-Haond S. & Belkhir K. 2007: GenClone: a computer program to analyse genotypic data, test for clonality and describe spatial clonal organization. *Mol. Ecol. Notes* 7: 15–17.
- Arnaud-Haond S., Alberto F., Procaccini G., Serrao E.A. & Duarte C.M. 2005: Assessing genetic diversity in clonal organisms: low diversity or low resolution? Combining power and cost-efficiency in selecting markers. *J. Hered.* 96: 1–8.
- Atarhouch T., Rami M., Cattaneo-Berrebi G., Ibanez C., Augros S., Boissin E., Dakkak A. & Berrebi P. 2003: New primers for EPIC amplification of intron sequences for fish and other vertebrate population genetic studies. *BioTechniques* 35: 676–682.
- Aurelle D., Cattaneo-Berrebi G. & Berrebi P. 2002: Natural and artificial secondary contact in brown trout (*Salmo trutta*, L.) in the French western Pyrenees assessed by allozymes and microsatellites. *Heredity* 89: 171–183.
- Baroiller J.-F. & Guiguen Y. 2001: Endocrine and environmental aspects of sex differentiation in gonochoristic fish. In: Scherer G. & Schmid M. (eds.), *Genes and mechanisms in vertebrate sex determination*. Birkhäuser Verlag, Basel-Switzerland: 177–201.
- Belkhir K., Borsa P., Goudet J., Chikhi L. & Bonhomme F. 1998: Genetix, logiciel sous Windows™ pour la génétique des populations. *Laboratoire Génome et Populations, CNRS UPR 9060, Université Montpellier II, Montpellier, France*. <http://www.univ-montp2.fr/~genetix/genetix.htm>
- Berrebi P. 1995: Speciation of the genus *Barbus* in the north Mediterranean basin: recent advances from biochemical genetics. *Biol. Conserv.* 72: 237–249.
- Berrebi P. & Tsigenopoulos C. 2002: Contribution des séquences mitochondriales à l'éclatement phylogénétique du genre polyploïde *Barbus* (Téléostéens Cyprinidés). Implications biogéographiques. *Biosystema* 20: 49–56.
- Berrebi P. & Tsigenopoulos C. 2003: Phylogenetic organisation of the genus *Barbus sensu stricto* – a synthesis on the molecular markers data. In: Banarescu P. (ed.), *Freshwater fishes of Europe – Cyprinidae*. Aula-Verlag, Wiebelsheim: 11–22.
- Berrebi P., Cattaneo-Berrebi G. & Le Brun N. 1993: Natural hybridization of two species of tetraploid barbels: *Barbus meridionalis* and *B. barbus* (Osteichthyes, Cyprinidae) in southern France. *Biol. J. Linn. Soc.* 48: 319–333.
- Berrebi P., Chenuil A., Kotlik P., Machordom A. & Tsigenopoulos C.S. 2013: Disentangling the evolutionary history of the genus *Barbus sensu lato*, a twenty years adventure. In: Alves J. (ed.), *Professor Carlos Almaça (1934-2010) – Estado da Arte em Áreas Científicas que Desenvolveu*. Museu Nacional de Historia Natural e da Ciencia, Lisboa.
- Berrebi P., Kottelat M., Skelton P. & Ráb P. 1996: Systematics of *Barbus*: state of the art and heuristic comments. *Folia Zool.* 45: 5–12.
- Berrebi P., Kraiem M.M., Doadrio I., El Gharbi S. & Cattaneo-Berrebi G. 1995: Ecological and genetic differentiation of *Barbus callensis* populations in Tunisia. *J. Fish Biol.* 47: 850–864.
- Black W.C. & Krafus E.S. 1985: A FORTRAN program for the calculation and analysis of two-locus linkage disequilibrium coefficients. *Theor. Appl. Genet.* 70: 491–496.
- Bouhbouh S. 2002: Bio-écologie de *Barbus callensis* (Valencienne 1842) et *Barbus fritschii* (Günther 1874) au niveau du réservoir Allal El Fassi (Maroc). *Thèse de la Faculté des Sciences Dhar El Mehraz, Fes, Maroc*.
- Bulger A.J. & Schultz R.J. 1979: Heterosis and interclonal variation in thermal tolerance in unisexual fishes. *Evolution* 33: 848–859.
- Castelli M. 1994: Study on sex determination in the common barbel (*Barbus barbus* L.) (Pisces, Cyprinidae) using gynogenesis. In: Beaumont A.R. (ed.), *Genetics and evolution of aquatic organisms*. Chapman & Hall, London: 509–519.
- Chevassus B. 1998: Modification du phénotype sexuel et du mode de reproduction chez les poissons Salmonidés: inversion sexuelle hormonale, gynogenèse, hybridation interspécifique et polyploïdisation. *Thèse de l'Université Paris XI-Orsay*: 162.
- Chow S. & Takeyama H. 1998: Intron length variation observed in the creatine kinase and ribosomal protein genes of the swordfish *Xiphias gladius*. *Fish. Sci.* 64: 397–402.
- Collares-Pereira M.J., Matos I., Morgado-Santos M. & Coelho M.M. 2013: Natural pathways towards polyploidy in animals: the *Squalius alburnoides* fish complex as a model system to study genome size and genome reorganization in polyploids. *Cytogenet. Genome Res.* 140: 97–116.
- Durand J.-D., Tsigenopoulos C.S., Unlu E. & Berrebi P. 2002: Phylogeny and biogeography of the family Cyprinidae in the Middle East inferred from Cytochrome *b* DNA evolutionary significance of this region. *Mol. Phylogenet. Evol.* 22: 91–100.
- El Gharbi S. 1994: Biosystématique, évolution et biogéographie dans les interactions hôte-parasite le modèle *Barbus* (Cyprinidés) *Dactylogyrus* (Monogenea). *Thèse d'Etat, Faculté des sciences de Rabat (Maroc)*.
- Golubtsov A.S. & Krysanov E.Y. 1993: Karyological study of some cyprinid species from Ethiopia. The ploidy differences between large and small *Barbus* of Africa. *J. Fish Biol.* 42: 445–455.
- Gui J.F. & Zhou L. 2010: Genetic basis and breeding application of clonal diversity and dual reproduction modes in polyploid *Carassius auratus gibelio* (review). *Sci. China Life Sci.* 53: 409–415.

- Howes G.J. 1987: The phylogenetic position of the Yugoslavian cyprinid fish genus *Aulopyge* Heckel, 1841, with an appraisal of the genus *Barbus* Cuvier & Cloquet, 1816, and the subfamily Cyprininae. *B. Brit. Mus. Nat. Hist. (Zool.)* 52: 165–196.
- Hubbs C.L. & Hubbs L.C. 1932: Apparent parthenogenesis in nature, in a form of fish of hybrid origin. *Science* 76: 628–630.
- Jiang F.F., Wang Z.W., Zhou L., Jiang L., Zhang X.J., Apalikova O.V., Brykov V.A. & Gui J.F. 2013: High male incidence and evolutionary implications of triploid form in northeast Asia *Carassius auratus* complex. *Mol. Phylogenet. Evol.* 66: 350–359.
- Machordom A. & Doadrio I. 2001: Evidence of a Cenozoic Betic-Kabilian connection based on freshwater fish phylogeny (*Luciobarbus*, Cyprinidae). *Mol. Phylogenet. Evol.* 18: 252–263.
- Moore W.S. & McKay F.E. 1971: Coexistence in unisexual-bisexual species complexes of Poeciliopsis (Pisces: Poeciliidae). *Ecology* 52: 791–799.
- Myers G.S. 1960: Preface to any future classification of the cyprinid fishes of the genus *Barbus*. *Stanford Ichthyol. Bull.* 7: 212–215.
- Otto S.P. & Whitton J. 2000: Polyploid incidence and evolution. *Annu. Rev. Genet.* 34: 401–437.
- Parks J.C. & Werth C.R. 1993: A study of spatial features of clones in a population of bracken fern, *Pteridium-Aquilinum* (Dennstaedtiaceae). *Am. J. Bot.* 80: 537–544.
- Pasteur N., Pasteur G., Bonhomme F., Catalan J. & Britton-Davidian J. 1987: Manuel de génétique par électrophorèse des protéines. *Lavoisier, Paris*: 217.
- Ryan M.J., Dries L.A., Batra P. & Hillis D.M. 1996: Male mate preferences in a gynogenetic species complex of Amazon mollies. *Anim. Behav.* 52: 1225–1236.
- Sambrook J., Fritsch E.F. & Maniatis T. 1989: Molecular cloning: a laboratory manual. *Cold Spring Harbor, New York*.
- Schultz R.J. 1977: Evolution and ecology of unisexual fishes. *Evol. Biol.* 10: 277–331.
- Schultz R.J. 1980: Role of polyploidy in the evolution of fishes. In: Lewis W.H. (ed.), Polyploidy: biological relevance. *Plenum Press, New York*: 313–340.
- Shaklee J.B., Allendorf F.W., Morizot D.C. & Whitt G.S. 1990: Gene nomenclature for protein-coding loci in fish. *Trans. Amer. Fish. Soc.* 119: 2–15.
- Skelton P. 2002: Changes to the scientific and common names of southern African freshwater fishes. *Afr. J. Aquat. Sci.* 27: 171–174.
- Tibayrenc M., Kjellberg F. & Ayala F. 1990: A clonal theory of parasitic protozoa: the population structures of *Entamoeba*, *Giardia*, *Leishmania*, *Naegleria*, *Plasmodium*, *Trichomonas* and *Trypanosoma* and their medical and taxonomical consequences. *Proc. Natl. Acad. Sci. USA* 87: 2414–2418.
- Tsigenopoulos C., Karakousis Y. & Berrebi P. 1999: The North Mediterranean *Barbus* lineage: phylogenetic hypotheses and taxonomic implications based on allozyme data. *J. Fish Biol.* 54: 267–286.
- Tsigenopoulos C.S., Ráb P., Naran D. & Berrebi P. 2002: Multiple origins of polyploidy in the phylogeny of southern Africans barbs (Cyprinidae) as inferred from mtDNA marks. *Heredity* 88: 466–473.
- Turner B.J., Brett B.-L.H., Rasch E.M. & Balsano J.S. 1980: Evolutionary genetics of a gynogenetic fish, *Poecilia formosa*, the Amazon molly. *Evolution* 34: 246–258.
- Uyeno T. & Smith G.R. 1972: Tetraploid origin of the karyotype of catostomid fishes. *Science* 175: 644–646.
- Weir B.S. & Cockerham C.C. 1984: Estimating F-statistics for the analysis of population structure. *Evolution* 38: 1358–1370.
- Yang L., Yang S.T., Wei X.H. & Gui J.F. 2001: Genetic diversity among different clones of the gynogenetic silver crucian carp, *Carassius auratus gibelio*, revealed by transferrin and isozyme markers. *Biochem. Genet.* 39: 213–225.
- Young A.G., Hill J.H., Murray B.G. & Peakall R. 2002: Breeding system, genetic diversity and clonal structure in the sub-alpine forb *Rutidosis leiolepis* F. Muell. (Asteraceae). *Biol. Conserv.* 106: 71–78.

**Appendix.** Genotypes of the 39 barbels analyzed at five polymorphic loci. Individuals sharing the same multiloci genotypes are indicated by the same letter in the first column.

Fish numbers	enzyme PGI-2*	enzyme PGI-3*	enzyme PGI-4*	intron RP	intron Mlc2a
1	100100	100120	100080	100100	100110
6	100100	100110	100100	100110	100100
9	100100	100110	100080	100110	100100
10	100100	100100	100080	110110	100110
11	100100	100100	100080	100110	100110
15	100100	100120	100080	100110	100110
20	100080	100110	100080	100100	100100
21	100080	120120	100080	100100	100110
22	100100	100110	100080	100110	100100
28	100100	100090	100100	-	100100
31	100100	100110	100100	100110	100110
33	100100	100120	100100	100100	100110
35	100080	100110	100100	100100	100110
38	100080	100110	100080	100110	100100
a19	100080	100110	100080	100100	100110

a5	100080	100110	100080	100100	100110
b12	100080	100110	100100	100100	100100
b32	100080	100110	100100	100100	-
c24	100100	100110	080080	100100	100100
c30	100100	100110	080080	100100	100100
d17	100100	100110	080080	100100	100110
d8	100100	100110	080080	100100	100110
e23	100100	100110	100080	100100	100100
e29	100100	100110	100080	100100	100100
f13	100100	100110	100080	100100	100110
f18	100100	100110	100080	100100	100110
f2	100100	100110	100080	100100	100110
f3	100100	100110	100080	100100	100110
f34	100100	100110	100080	100100	100110
f37	100100	100110	100080	100100	100110
g16	100100	100110	100080	100110	100110
g27	100100	100110	100080	100110	100110
g39	100100	100110	100080	100110	100110
h14	100100	100110	100100	100100	100100
h25	100100	100110	100100	100100	100100
h36	100100	100110	100100	100100	100100
h4	100100	100110	100100	100100	100100
i26	100100	100120	100080	100100	100100
i7	100100	100120	100080	100100	-

---