

## **An updated checklist of the birds of Rio Grande do Norte, Brazil, with comments on new, rare, and unconfirmed species**

Authors: Sagot-Martin, François, Lima, Rafael Dantas, Pacheco, José Fernando, Irusta, Jorge Bañuelos, Pichorim, Mauro, et al.

Source: Bulletin of the British Ornithologists' Club, 140(3) : 218-298

Published By: British Ornithologists' Club

URL: <https://doi.org/10.25226/bboc.v140i3.2020.a2>

---

BioOne Complete (complete.BioOne.org) is a full-text database of 200 subscribed and open-access titles in the biological, ecological, and environmental sciences published by nonprofit societies, associations, museums, institutions, and presses.

Your use of this PDF, the BioOne Complete website, and all posted and associated content indicates your acceptance of BioOne's Terms of Use, available at [www.bioone.org/terms-of-use](http://www.bioone.org/terms-of-use).

Usage of BioOne Complete content is strictly limited to personal, educational, and non - commercial use. Commercial inquiries or rights and permissions requests should be directed to the individual publisher as copyright holder.

---

BioOne sees sustainable scholarly publishing as an inherently collaborative enterprise connecting authors, nonprofit publishers, academic institutions, research libraries, and research funders in the common goal of maximizing access to critical research.

# An updated checklist of the birds of Rio Grande do Norte, Brazil, with comments on new, rare, and unconfirmed species

by François Sagot-Martin<sup>\*1</sup>, Rafael Dantas Lima\*, José Fernando Pacheco, Jorge Bañuelos Irusta, Mauro Pichorim & David Maurice Hassett

Received 12 November 2019; revised 23 May 2020; published 21 September 2020

<http://zoobank.org/urn:lsid:zoobank.org:pub:9B9E545B-1768-4858-87B8-41BDD1CAD22D>

**SUMMARY.**—We present a detailed bird list for the north-east Brazilian state of Rio Grande do Norte. This is the first inventory of the avifauna of the state that complies with strict criteria as to its compilation and documentation. To achieve this, we reviewed all records published for the state, as well as records in several online databases and museum collections. Our consolidated list is enhanced by our own unpublished records, and encompasses 425 species, 391 of them based on available specified physical evidence. The list includes 12 taxa not previously documented in the state and 35 documented taxa that are considered nationally or globally threatened. Other noteworthy records include three documented range extensions of 1,000 km or more and the first proven breeding by three species in far north-eastern Brazil. The rather diverse avifauna of Rio Grande do Norte reflects its location at the juxtaposition of three major domains: Atlantic Forest, Caatinga, Cerrado, and their ecotones, bordered by a marine environment.

Our knowledge of the birds of the state of Rio Grande do Norte, in the extreme north-east corner of South America, remained impoverished until just two decades ago, when field work began to reveal a hitherto unsuspected diversity. Until the early 20th century, the lack of reliable information was almost absolute, because practically none of the great historical naturalists that traversed the Brazilian north-east visited the state (Pacheco 2004). The sole exception is a single reference in G. Marcgrave's famous *Historia rerum naturalis Brasiliae* to the abundance of Greater Rheas *Rhea americana* in the open zones of Rio Grande do Norte. This seems to be the first mention of birds in the state, from among the descriptions of at least 134 wild bird species found in the nearby states of Pernambuco and Paraíba (Marcgrave 1648, Teixeira 1992). Prior to this, there are only ancient cave paintings showing rheas and flamingos (presumably American Flamingo *Phoenicopterus ruber*) at São Rafael (Souza & Medeiros 1982) and Carnaúba dos Dantas (Macedo 2003), rheas, parrots and toucans at Parelhas (Silva 2003), and macaws, parrots and presumably herons at Apodi (Fundação Amigos do Lajedo de Soledade pers. comm.), all in Rio Grande do Norte. The rhea appeared on the state's coat of arms during the colonial period, initially by order of Prince Mauricio of Nassau (Sick 1997).

Throughout the 20th century, references to the birds of Rio Grande do Norte were still scarce: the work of O. M. O. Pinto included just one reference, to *Rhea*, in the first part of his catalogue (Pinto 1938) and three in the second part (Pinto 1944). D. Lamm occasionally visited southern Rio Grande do Norte and included Grey-breasted Martin *Progne chalybea* in an annotated list of the birds of Paraíba and Pernambuco states (Lamm 1948). Faria (1961) published the results of interviews with hunters about birds in the state's Seridó region.

<sup>1\*</sup> These authors contributed equally to this work.

Coelho (1978) referred to just one specimen, a Savanna Hawk *Heterospizias meridionalis*, collected in Rio Grande do Norte among a list of 273 birds of Pernambuco. Lara-Resende & Leal (1982) provided the first information concerning recoveries of ringed birds for the state. H. Sick, after visiting Rio Grande do Norte for a brief period, mentioned only 13 species in his keynote *Ornitologia brasileira* (Sick 1985). By the end of the century, several specific publications on the state's avifauna had become available: Antas (1987a,b), Sick *et al.* (1988), Azevedo & Antas (1990), Praxedes *et al.* (1997), Varela-Freire (1997), Varela-Freire & Araújo (1997), Varela-Freire *et al.* (1997), Rocha-Neto *et al.* (1998), Silveira & Varela-Freire (1999), the first state bird list by Varela-Freire *et al.* (1999), Azevedo *et al.* (2000), Medeiros *et al.* (2000), Nascimento (2000) and Nascimento & Larrazábal (2000). Some of these contained misidentifications and doubtful records, which have been repeated until the present. Other useful contributions on the birds of the state included Sick (1991), Teixeira *et al.* (1993), Pacheco & Whitney (1995), Cordeiro *et al.* (1996), Silva e Silva (1996) and Whitney *et al.* (2000), but avian inventories were scarce.

In the 21st century, the situation has changed considerably, with numerous surveys and articles on the state's birds, by Silveira *et al.* (2001), Larrazábal *et al.* (2002), Olmos (2003), Araújo *et al.* (2004) and Azevedo *et al.* (2004), among others detailed in the species accounts that follow. These works have resulted in a significant increase in the number of species recorded in the state, resulting in the present overview, wherein we highlight the status of 133 species, detail many noteworthy records, and correct some prior misidentifications. We also comment on aspects of the biogeography and conservation of the region's avifauna.

## Methods

**Study area.**—North-east Brazil lies between the Amazon Basin and the Atlantic Forest, with influences from both regions (Werneck 2011, Ledo & Colli 2017; Fig. 1A). Rio Grande do Norte is bordered by the Atlantic Ocean to the north and east (Fig. 1B). The Brazilian continental shelf reaches its narrowest point near Natal, with the shelf break and its nutrient-rich upwellings just 27 km offshore (Lees *et al.* 2015).

The state can be divided into two main phytogeographic domains (Fig. 1C): (1) the Atlantic Forest, a narrow coastal strip of semi-deciduous forest, now quite fragmented and degraded, and forming part of the Pernambuco Centre of Endemism (Brown 1982, Silva & Castelletti 2003), plus severely modified mangroves along the coast, and (2) the Caatinga, a seasonally dry tropical forest which occupies practically the entire interior as far as the northern border, covering >90% of the state (J. M. C. Silva *et al.* 2017), today appearing more like a scrubby steppe-savanna (Veloso *et al.* 1991).

In addition, some of the southwestern 'serras' (Fig. 1C) are influenced by the dominant humid trade winds and possess vegetation more like the Atlantic Forest (Queiroz *et al.* 2017); they are referred to as 'brejos de altitude' (Andrade-Lima 1982, Tabarelli & Santos 2004).

It is frequently overlooked that on the state's north-east coast there is an isolated patch of Cerrado (Oliveira *et al.* 2012, Pichorim *et al.* 2014a), a generic term for the savanna formations typical of the central uplands of Brazil. Moreover, so-called 'tabuleiros costeiros' (literally, 'coastal tablelands'), occur throughout the littoral—as Cerrado-like stretches mixed with Atlantic Forest on the east coast and Caatinga mixed with Atlantic *restinga* on the north.

According to the Köppen-Geiger climate classification system (Geiger 1961), Rio Grande do Norte lies in a region that varies from 'tropical savanna climate with dry winter' to 'tropical savanna climate with dry summer', Aw / As, and 'hot semi-arid climate', BSh. The state shows (1) a distinctly wetter eastern region, with a 'tropical oriental northeast' climate and rainfall of 900–1,500 mm / year generally concentrated in the austral winter




Figure 1. The state of Rio Grande do Norte in Brazil and South America (A). Far north-east Brazil with the location of the Pernambuco Centre of Endemism (B). Main phytophysionomies in Rio Grande do Norte, with 'serras' indicated by shading (C).



(March–July); and (2) an extensive semi-arid area with a ‘tropical equatorial zone’ climate, rainfall of 400–900 mm / year and a dry season lasting up to nine months (April–December) (Diniz & Pereira 2015).

The state is characterised by mean temperatures of 23.2–26.8°C in the interior—minimum in Serra de Martins—and 25.5–26.5°C on the east coast. Trade winds from the south-east dominate year-round, but are stronger during August–November and less intense in February–March (Amarante *et al.* 2003). Highest mean relative humidity is in Natal, virtually always above 80% with a peak of 84% in June, which is the wettest month (Espírito Santo & Silva 2016), and decreasing from east to west, to 47% in the Caatinga (INMET 2019).

**Methods.**—We compiled bird records from the state’s mainland plus coastal and offshore areas up to the edge of the continental shelf, thus excluding the oceanic Atol das Rocas Reserve and surrounding waters, which belong to Brazil but are not part of Rio Grande do Norte (Federal Decree 83549, 5 June 1979; M. Brito pers. comm.). We reviewed records from the literature and included our own observations up to 31 March 2019.

We evaluated available records in various online databases and museum collections, as follows: Coleção de Aves Heretiano Zenaide, Universidade Federal da Paraíba, João Pessoa (CAHZ), 60 specimens; Coleção Ornitológica da Universidade Federal do Rio Grande do Norte, Natal (COUFRN), 1,020 specimens; Centro de Pesquisas do Parque Estadual das Dunas de Natal (CPPD), two specimens; Fonoteca Neotropical Jacques Vielliard, Campinas (FNJV), one record; the Internet Bird Collection, [www.hbw.com/ibc](http://www.hbw.com/ibc) (IBC), 35 records (now incorporated into ML, below); Louisiana State University Museum of Natural Science, Baton Rouge (LSUMZ), four specimens; Museum of Comparative Zoology, Harvard University, Cambridge, MA (MCZ), four specimens; Macaulay Library, [www.macaulaylibrary.org](http://www.macaulaylibrary.org) (ML), 249 records; Museu Nacional, Universidade Federal do Rio de Janeiro (MNRJ), 93 specimens; Museu Paraense Emílio Goeldi, Belém (MPEG), six specimens; Museu de Zoologia da Universidade de São Paulo (MZUSP), one specimen; San Diego Natural History Museum, CA (SDNHM), one specimen; Coleção de Aves da Universidade Federal de Pernambuco (UFPE), 205 specimens; University of Michigan Museum of Zoology, Ann Arbor, MI (UMMZ), two specimens; National Museum of Natural History, Smithsonian Institution, Washington DC (USNM), one specimen; WikiAves, [www.wikiaves.com.br](http://www.wikiaves.com.br) (WA), 19,680 records; and Xeno-canto, [www.xeno-canto.org](http://www.xeno-canto.org) (XC), 60 records.

We follow the nomenclature and taxonomic sequence of the Brazilian Ornithological Records Committee (Piacentini *et al.* 2015), where the primary and secondary lists form the consolidated list of accepted species for Brazil, with those belonging to the secondary list in square brackets []. The tertiary list and a list of species of probable / possible occurrence in Rio Grande do Norte comprise Appendices 1–2. Our definitions for the three first lists are:

**Primary list.**—Species with at least one unequivocal state record based on physical evidence, available for independent re-assessment (complete or partial specimen, photograph, or audio or video recording) and which unambiguously provides a certain diagnosis of the taxon involved (Carlos *et al.* 2010).

**Secondary list.**—Species with published state records, but without available physical evidence, plus our own records when no others can be cited for inclusion of a species.

**Tertiary list.**—Species with published state records, but for which the available evidence is either questionable or invalid, and whose occurrence in Rio Grande do Norte appears improbable based on current knowledge.

To justify inclusion in the primary list, we present the documentary evidence or reference to support it, giving priority to the oldest available record, unless the first documented record is of low quality or difficult to access.

TABLE 1

Consolidated list of the birds of Rio Grande do Norte, Brazil. Species on the secondary list in square brackets [] ; \* = subspecies based on distribution; National occurrence: R = Resident, VN = seasonal visitor from the Northern Hemisphere, VS = seasonal visitor from southern South America, VA(N) = vagrant from north, E = endemic to Brazil, and # = status not confirmed; Conservation: national (listed first) and global status: CR = Critically Endangered, EN = Endangered, VU = Vulnerable, NT = Near Threatened, LC = Least Concern, DD = Data Deficient, and NE = Not Evaluated (subspecies evaluated only at national level).

| Taxon   | English name                 | National occurrence | National; Global conservation | Specimen    | Photo      | Sound                      |
|---|------------------------------|---------------------|-------------------------------|-------------|------------|----------------------------|
| <b>RHEIFORMES</b>                                     |                              |                     |                               |             |            |                            |
| <b>Rheidae</b>  |                              |                     |                               |             |            |                            |
| <i>Rhea americana</i> (Linnaeus, 1758)                | Greater Rhea                 | R                   | LC; NT                        |             | WA 1624047 |                            |
| <i>R. a. americana</i> (Linnaeus, 1758)*              |                              |                     |                               |             |            |                            |
| <b>TINAMIFORMES</b>                                   |                              |                     |                               |             |            |                            |
| <b>Tinamidae</b>                                      |                              |                     |                               |             |            |                            |
| <i>Crypturellus noctivagus</i> (Wied, 1820)           | Yellow-legged Tinamou        | R, E                | NT; NT                        |             |            | WA 3217181                 |
| <i>C. n. zabele</i> (Spix, 1825)*                     |                              |                     |                               |             |            |                            |
|   |                              |                     | VU                            |             |            |                            |
| <i>Crypturellus parvirostris</i> (Wagler, 1827)       | Small-billed Tinamou         | R                   | LC; LC                        | COUFRN 1194 |            | WA 1254365                 |
| <i>Crypturellus tataupa</i> (Temminck, 1815)          | Tataupa Tinamou              | R                   | LC; LC                        | COUFRN 1116 | WA 2222231 | WA 1254765                 |
| <i>C. t. lepidotus</i> (Swainson, 1837)*              |                              |                     |                               |             |            |                            |
| [ <i>Rhynchotus rufescens</i> (Temminck, 1815)]       | Red-winged Tinamou           |                     | LC; LC                        |             |            |                            |
| <i>Nothura boraquira</i> (Spix, 1825)                 | White-bellied Nothura        | R                   | LC; LC                        |             | WA 342523  | WA 3300847                 |
| <i>Nothura maculosa</i> (Temminck, 1815)              | Spotted Nothura              | R                   | LC; LC                        | COUFRN 1189 | WA 324818  | Minns <i>et al.</i> (2010) |
| <i>N. m. cearensis</i> Naumburg, 1932*                |                              |                     |                               |             |            |                            |
| <b>ANSERIFORMES</b>                                   |                              |                     |                               |             |            |                            |
| <b>Anatidae</b>                                       |                              |                     |                               |             |            |                            |
| <i>Dendrocygna bicolor</i> (Vieillot, 1816)           | Fulvous Whistling Duck       | R                   | LC; LC                        |             | WA 2168387 |                            |
| <i>Dendrocygna viduata</i> (Linnaeus, 1766)           | White-faced Whistling Duck   | R                   | LC; LC                        | COUFRN 1011 | WA 343450  | WA 2187215                 |
| <i>Dendrocygna autumnalis</i> (Linnaeus, 1758)        | Black-bellied Whistling Duck | R                   | LC; LC                        |             | WA 3032864 |                            |
| <i>D. a. autumnalis</i> (Linnaeus, 1758)*             |                              |                     |                               |             |            |                            |
| <i>Cairina moschata</i> (Linnaeus, 1758)              | Muscovy Duck                 | R                   | LC; LC                        |             | WA 2888883 |                            |
| <i>Sarkidiornis sylvicola</i> Ihering & Ihering, 1907 | Comb Duck                    | R                   | LC; LC                        |             | WA 2120536 |                            |
| <i>Amazonetta brasiliensis</i> (J. F. Gmelin, 1789)   | Brazilian Teal               | R                   | LC; LC                        |             | WA 336213  |                            |
| <i>A. b. brasiliensis</i> (J. F. Gmelin, 1789)*       |                              |                     |                               |             |            |                            |



| Taxon   | English name              | National occurrence | National; Global conservation | Specimen    | Photo      | Sound      |  |  |  |
|---|---------------------------|---------------------|-------------------------------|-------------|------------|------------|--|--|--|
| <i>Anas bahamensis</i><br>Linnaeus, 1758                      | White-cheeked Pintail     | R                   | LC; LC                        |             | WA 238944  |            |  |  |  |
| <i>A. b. bahamensis</i> Linnaeus, 1758*                       |                           |                     |                               |             |            |            |  |  |  |
| [ <i>Anas discors</i> Linnaeus, 1766]                         | Blue-winged Teal          | VA(N)               | LC; LC                        |             |            |            |  |  |  |
| <i>Netta erythrophthalma</i><br>(Wied, 1833)                  | Southern Pochard          | R                   | LC; LC                        | MNRJ 34409  | WA 2495205 |            |  |  |  |
| <i>N. e. erythrophthalma</i> (Wied, 1833)*                    |                           |                     |                               |             |            |            |  |  |  |
| <i>Nomonyx dominicus</i><br>(Linnaeus, 1766)                  | Masked Duck               | R                   | LC; LC                        |             | WA 973135  |            |  |  |  |
| GALLIFORMES   |                           |                     |                               |             |            |            |  |  |  |
| Cracidae  |                           |                     |                               |             |            |            |  |  |  |
| <i>Penelope supercilialis</i><br>Temminck, 1815               | Rusty-margined Guan       | R                   | LC; LC                        | COUFRN 1196 | WA 515077  |            |  |  |  |
| <i>P. s. ochromitra</i> Spix, 1825*                           |                           |                     | DD                            |             |            |            |  |  |  |
| <i>P. s. alagoensis</i> Nardelli, 1993                        |                           |                     | CR                            |             |            |            |  |  |  |
| <i>Penelope jacucaca</i> Spix, 1825                           | White-browed Guan         | R, E                | VU; VU                        |             | WA 2016795 |            |  |  |  |
| <i>Ortalis araucuan</i> (Spix, 1825)                          | East Brazilian Chachalaca | R, E                | LC; LC                        |             | WA 609213  | XC 430030  |  |  |  |
| Podicipedidae   |                           |                     |                               |             |            |            |  |  |  |
| <i>Tachybaptus dominicus</i><br>(Linnaeus, 1766)              | Least Grebe               | R                   | LC; LC                        |             | WA 340531  | WA 2982539 |  |  |  |
| <i>T. d. brachyrhynchus</i> (Chapman, 1899)*                  |                           |                     |                               |             |            |            |  |  |  |
| <i>Podilymbus podiceps</i><br>(Linnaeus, 1758)                | Pied-billed Grebe         | R                   | LC; LC                        |             | WA 240767  |            |  |  |  |
| <i>P. p. antarcticus</i> (Lesson, 1842)*                      |                           |                     |                               |             |            |            |  |  |  |
| Phoenicopteridae  |                           |                     |                               |             |            |            |  |  |  |
| [ <i>Phoenicopterus ruber</i><br>Linnaeus, 1758]              | American Flamingo         | R#                  | DD; LC                        |             |            |            |  |  |  |
| Spheniscidae  |                           |                     |                               |             |            |            |  |  |  |
| [ <i>Spheniscus magellanicus</i><br>(J. R. Forster, 1781)]    | Magellanic Penguin        | VS                  | NT; NT                        |             |            |            |  |  |  |
| PROCELLARIIFORMES   |                           |                     |                               |             |            |            |  |  |  |
| Diomedidae  |                           |                     |                               |             |            |            |  |  |  |
| [ <i>Thalassarche chlororhynchos</i><br>(J. F. Gmelin, 1789)] | Yellow-nosed Albatross    | VS                  | EN; EN                        |             |            |            |  |  |  |
| Procellariidae  |                           |                     |                               |             |            |            |  |  |  |
| <i>Fulmarus glacialis</i><br>(A. Smith, 1840)                 | Southern Fulmar           | VS                  | LC; LC                        | MCZ 186306  |            |            |  |  |  |
| <i>Daption capense</i><br>(Linnaeus, 1758)                    | Cape Petrel               | VS                  | LC; LC                        | MCZ 226730  |            |            |  |  |  |
| <i>D. c. capense</i> (Linnaeus, 1758)*                        |                           |                     |                               |             |            |            |  |  |  |
| [ <i>Pterodroma madeira</i><br>Mathews, 1934]                 | Zino’s Petrel             |                     | EN; EN                        |             |            |            |  |  |  |

| Taxon   | English name               | National occurrence | National; Global conservation | Specimen    | Photo        | Sound |
|---|----------------------------|---------------------|-------------------------------|-------------|--------------|-------|
| [ <i>Pterodroma deserta</i> Mathews, 1934]                  | Desertas Petrel            |                     | CR; EN                        |             |              |       |
| <i>Pterodroma arminjoniana</i> (Giglioli & Salvadori, 1869) | Trindade Petrel            | R                   | CR; VU                        |             | WA 1592035   |       |
| [ <i>Bulweria bulwerii</i> (Jardine & Selby, 1828)]         | Bulwer's Petrel            | VN#                 | NE; LC                        |             |              |       |
| <i>Calonectris borealis</i> (Cory, 1881)                    | Cory's Shearwater          | VN                  | LC; LC                        | COUFRN 948  | WA 2518923   |       |
| <i>Puffinus griseus</i> (J. F. Gmelin, 1789)                | Sooty Shearwater           | VS                  | LC; NT                        |             | ML 77958651  |       |
| <i>Puffinus gravis</i> (O'Reilly, 1818)                     | Great Shearwater           | VS                  | LC; LC                        |             | WA 2813943   |       |
| <i>Puffinus puffinus</i> (Brünnich, 1764)                   | Manx Shearwater            | VN                  | LC; LC                        | COUFRN 831  | WA 1593190   |       |
| [ <i>Puffinus boydi</i> Mathews, 1912]                      | Boyd's Shearwater          | VN                  | ; NE                          |             |              |       |
| <b>Hydrobatidae</b>   |                            |                     |                               |             |              |       |
| <i>Fregetta tropica</i> (Gould, 1844)                       | Black-bellied Storm Petrel | VS#                 | LC; LC                        | COUFRN 120  |              |       |
| <i>F. t. tropica</i> (Gould, 1844)                          |                            |                     |                               |             |              |       |
| <i>Oceanites oceanicus</i> (Kuhl, 1820)                     | Wilson's Storm Petrel      | VS                  | LC; LC                        |             | WA 1593174   |       |
| <i>Oceanodroma leucorhoa</i> (Vieillot, 1818)               | Leach's Storm Petrel       | VN                  | LC; VU                        | COUFRN 728  | WA 1593164   |       |
| <i>O. l. leucorhoa</i> (Vieillot, 1818)*                    |                            |                     |                               |             |              |       |
| <b>CICONIIFORMES</b>  |                            |                     |                               |             |              |       |
| <b>Ciconiidae</b>   |                            |                     |                               |             |              |       |
| <i>Jabiru mycteria</i> (M. H. C. Lichtenstein, 1819)        | Jabiru                     | R                   | LC; LC                        |             | Fig. 3A      |       |
| <i>Mycteria americana</i> Linnaeus, 1758                    | Wood Stork                 | R                   | LC; LC                        |             | WA 1945521   |       |
| <b>SULIFORMES</b>   |                            |                     |                               |             |              |       |
| <b>Fregatidae</b>   |                            |                     |                               |             |              |       |
| <i>Fregata magnificens</i> Mathews, 1914                    | Magnificent Frigatebird    | R                   | LC; LC                        |             | WA 236127    |       |
| <b>Sulidae</b>  |                            |                     |                               |             |              |       |
| <i>Sula dactylatra</i> Lesson, 1831                         | Masked Booby               | R                   | LC; LC                        | COUFRN 1253 | ML 127037771 |       |
| <i>S. d. dactylatra</i> Lesson, 1831*                       |                            |                     |                               |             |              |       |
| <i>Sula sula</i> (Linnaeus, 1766)                           | Red-footed Booby           |                     | EN; LC                        | COUFRN 1275 |              |       |
| <i>S. s. sula</i> (Linnaeus, 1766)*                         |                            |                     |                               |             |              |       |
| [ <i>Sula leucogaster</i> (Boddaert, 1783)]                 | Brown Booby                | R                   | LC; LC                        |             |              |       |
| <b>Phalacrocoracidae</b>                                    |                            |                     |                               |             |              |       |
| <i>Nannopterum brasilianus</i> (J. F. Gmelin, 1789)         | Neotropic Cormorant        | R                   | LC; LC                        |             | WA 283214    |       |
| <i>N. b. brasilianus</i> (J. F. Gmelin, 1789)*              |                            |                     |                               |             |              |       |

| Taxon   | English name               | National occurrence | National; Global conservation | Specimen    | Photo      | Sound      |
|---|----------------------------|---------------------|-------------------------------|-------------|------------|------------|
| <b>Anhingidae</b>                               |                            |                     |                               |             |            |            |
| <i>Anhinga anhinga</i> (Linnaeus, 1766)         | Anhinga                    | R                   | LC; LC                        |             | WA 1366323 |            |
| <i>A. a. anhinga</i> (Linnaeus, 1766)*          |                            |                     |                               |             |            |            |
| <b>PELECANIFORMES</b>                           |                            |                     |                               |             |            |            |
| <b>Ardeidae</b>                                 |                            |                     |                               |             |            |            |
| <i>Tigrisoma lineatum</i> (Boddaert, 1783)      | Rufescent Tiger Heron      | R                   | LC; LC                        |             | WA 2712033 |            |
| <i>T. l. marmoratum</i> (Vieillot, 1817)*       |                            |                     |                               |             |            |            |
| <i>Botaurus pinnatus</i> (Wagler, 1829)         | Pinnated Bittern           | R                   | LC; LC                        |             | WA 622996  |            |
| <i>B. p. pinnatus</i> (Wagler, 1829)*           |                            |                     |                               |             |            |            |
| <i>Ixobrychus exilis</i> (J.F. Gmelin, 1789)    | Least Bittern              | R                   | LC; LC                        |             | WA 2411673 |            |
| <i>I. e. erythromelas</i> (Vieillot, 1817)*     |                            |                     |                               |             |            |            |
| <i>Ixobrychus involucris</i> (Vieillot, 1823)   | Stripe-backed Bittern      | R                   | LC; LC                        |             | WA 3065361 |            |
| <i>Nycticorax nycticorax</i> (Linnaeus, 1758)   | Black-crowned Night Heron  | R                   | LC; LC                        | USNM 330093 | WA 2723524 |            |
| <i>N. n. hoactli</i> (J. F. Gmelin, 1789)       |                            |                     |                               |             |            |            |
| <i>Nyctanassa violacea</i> (Linnaeus, 1758)     | Yellow-crowned Night Heron | R                   | LC; LC                        |             | WA 149007  |            |
| <i>N. v. cayennensis</i> (J. F. Gmelin, 1789)*  |                            |                     |                               |             |            |            |
| <i>Butorides striata</i> (Linnaeus, 1758)       | Striated Heron             | R                   | LC; LC                        | COUFRN 742  | WA 653335  |            |
| <i>B. s. striata</i> (Linnaeus, 1758)*          |                            |                     |                               |             |            |            |
| <i>Bubulcus ibis</i> (Linnaeus, 1758)           | Cattle Egret               | R                   | LC; LC                        |             | WA 269132  |            |
| <i>B. i. ibis</i> (Linnaeus, 1758)*             |                            |                     |                               |             |            |            |
| <i>Ardea cocoi</i> Linnaeus, 1766               | Cocoi Heron                | R                   | LC; LC                        |             | WA 295058  |            |
| <i>Ardea alba</i> Linnaeus, 1758                | Great Egret                | R                   | LC; LC                        |             | WA 625168  | WA 2605407 |
| <i>A. a. egretta</i> J.F. Gmelin, 1789*         |                            |                     |                               |             |            |            |
| [ <i>Pilherodius pileatus</i> (Boddaert, 1783)] | Capped Heron               | R                   | LC; LC                        |             |            |            |
| <i>Egretta thula</i> (Molina, 1782)             | Snowy Egret                | R                   | LC; LC                        |             | WA 266283  | WA 3300838 |
| <i>E. t. thula</i> (Molina, 1782)*              |                            |                     |                               |             |            |            |
| <i>Egretta caerulea</i> (Linnaeus, 1758)        | Little Blue Heron          | R                   | LC; LC                        |             | WA 92412   |            |
| <b>Threskiornithidae</b>                        |                            |                     |                               |             |            |            |
| [ <i>Theristicus caudatus</i> (Boddaert, 1783)] | Buff-necked Ibis           | R                   | LC; LC                        |             |            |            |
| <i>Platalea ajaja</i> Linnaeus, 1758            | Roseate Spoonbill          | R                   | LC; LC                        | SDNHM 35485 |            |            |



| Taxon  | English name                 | National occurrence | National; Global conservation | Specimen   | Photo      | Sound      |
|--|------------------------------|---------------------|-------------------------------|------------|------------|------------|
| <b>Cathartidae</b>                                     |                              |                     |                               |            |            |            |
| <i>Cathartes aura</i> (Linnaeus, 1758)                 | Turkey Vulture               | R                   | LC; LC                        |            | WA 248533  |            |
| <i>C. a. ruficollis</i> Spix, 1824                     |                              |                     |                               |            |            |            |
| <i>Cathartes burrovianus</i> Cassin, 1845              | Lesser Yellow-headed Vulture | R                   | LC; LC                        |            | WA 239455  |            |
| <i>C. b. urubutinga</i> Pelzeln, 1861*                 |                              |                     |                               |            |            |            |
| <i>Coragyps atratus</i> (Bechstein, 1793)              | Black Vulture                | R                   | LC; LC                        |            | WA 724431  |            |
| <i>C. a. brasiliensis</i> (Bonaparte, 1850)*           |                              |                     |                               |            |            |            |
| <i>Sarcoramphus papa</i> (Linnaeus, 1758)              | King Vulture                 | R                   | NT; LC                        |            | WA 1649096 |            |
| <b>ACCIPITRIFORMES</b>                                 |                              |                     |                               |            |            |            |
| <b>Pandionidae</b>                                     |                              |                     |                               |            |            |            |
| <i>Pandion haliaetus</i> (Linnaeus, 1758)              | Osprey                       | VN                  | LC; LC                        |            | WA 252946  |            |
| <i>P. h. carolinensis</i> (J. F. Gmelin, 1788)         |                              |                     |                               |            |            |            |
| <b>Accipitridae</b>                                    |                              |                     |                               |            |            |            |
| <i>Leptodon forbesi</i> (Swann, 1922)                  | White-collared Kite          | R, E                | EN; CR                        |            | WA 2263455 |            |
| <i>Chondrohierax uncinatus</i> (Temminck, 1822)        | Hook-billed Kite             | R                   | LC; LC                        |            | WA 1351596 |            |
| <i>C. u. uncinatus</i> (Temminck, 1822)*               |                              |                     |                               |            |            |            |
| [ <i>Elanoides forficatus</i> (Linnaeus, 1758)]        | Swallow-tailed Kite          | R                   | LC; LC                        |            |            |            |
| <i>Gampsonyx swainsonii</i> Vigors, 1825               | Pearl Kite                   | R                   | LC; LC                        |            | WA 324802  | WA 3267846 |
| <i>G. s. swainsonii</i> Vigors, 1825*                  |                              |                     |                               |            |            |            |
| <i>Elanus leucurus</i> (Vieillot, 1818)                | White-tailed Kite            | R                   | LC; LC                        |            | WA 249718  |            |
| <i>E. l. leucurus</i> (Vieillot, 1818)*                |                              |                     |                               |            |            |            |
| <i>Accipiter striatus</i> Vieillot, 1808               | Sharp-shinned Hawk           | R                   | LC; LC                        |            | WA 2837626 |            |
| <i>A. s. erythronemius</i> (Kaup, 1850)                |                              |                     | ; NE                          |            |            |            |
| <i>Accipiter bicolor</i> (Vieillot, 1817)              | Bicolored Hawk               | R                   | LC; LC                        |            | WA 3089927 |            |
| <i>A. b. pileatus</i> (Temminck, 1823)                 |                              |                     |                               |            |            |            |
| <i>Ictinia plumbea</i> (J. F. Gmelin, 1788)            | Plumbeous Kite               | R                   | LC; LC                        |            | WA 249719  |            |
| <i>Rostrhamus sociabilis</i> (Vieillot, 1817)          | Snail Kite                   | R                   | LC; LC                        | MNRJ 34415 | WA 338222  | WA 2980216 |
| <i>R. s. sociabilis</i> (Vieillot, 1817)*              |                              |                     |                               |            |            |            |
| <i>Geranospiza caerulescens</i> (Vieillot, 1817)       | Crane Hawk                   | R                   | LC; LC                        |            | WA 2712040 |            |
| <i>G. c. gracilis</i> (Temminck, 1821)*                |                              |                     |                               |            |            |            |
| <i>Buteogallus aequinoctialis</i> (J. F. Gmelin, 1788) | Rufous Crab Hawk             | R                   | NT; NT                        |            | WA 2874182 | WA 2868593 |
| <i>Heterospizias meridionalis</i> (Latham, 1790)       | Savanna Hawk                 | R                   | LC; LC                        | UFPE 382   | WA 238946  | WA 1230486 |

| Taxon  | English name                | National occurrence | National; Global conservation | Specimen   | Photo      | Sound                      |
|--|-----------------------------|---------------------|-------------------------------|------------|------------|----------------------------|
| <i>Urubitinga urubitinga</i><br>(J. F. Gmelin, 1788)   | Great Black Hawk            | R                   | LC; LC                        |            | WA 1011184 |                            |
| <i>U. u. urubitinga</i> (J. F. Gmelin, 1788)*          |                             |                     |                               |            |            |                            |
| <i>Rupornis magnirostris</i><br>(J. F. Gmelin, 1788)   | Roadside Hawk               | R                   | LC; LC                        | COUFRN 802 | WA 1365656 | XC 430044                  |
| <i>R. m. nattereri</i> (P. L. Sclater & Salvin, 1869)* |                             |                     |                               |            |            |                            |
| <i>Parabuteo unicinctus</i><br>(Temminck, 1824)        | Harris's Hawk               | R                   | LC; LC                        |            | WA 2518913 | WA 3202039                 |
| <i>P. u. unicinctus</i> (Temminck, 1824)*              |                             |                     |                               |            |            |                            |
| <i>Geranoaetus albicaudatus</i><br>(Vieillot, 1816)    | White-tailed Hawk           | R                   | LC; LC                        |            | WA 47525   |                            |
| <i>G. a. albicaudatus</i> (Vieillot, 1816)*            |                             |                     |                               |            |            |                            |
| <i>Geranoaetus melanoleucus</i><br>(Vieillot, 1819)    | Black-chested Buzzard-Eagle | R                   | LC; LC                        |            | WA 1331947 | XC 4838                    |
| <i>G. m. melanoleucus</i> (Vieillot, 1819)*            |                             |                     |                               |            |            |                            |
| <i>Buteo nitidus</i> (Latham, 1790)                    | Grey-lined Hawk             | R                   | LC; LC                        |            | WA 3065367 | WA 2577547                 |
| <i>B. n. pallidus</i> (Todd, 1915)*                    |                             |                     |                               |            |            |                            |
| <i>Buteo brachyurus</i> Vieillot, 1816                 | Short-tailed Hawk           | R                   | LC; LC                        | COUFRN 369 | WA 249715  |                            |
| <i>B. b. brachyurus</i> Vieillot, 1816*                |                             |                     |                               |            |            |                            |
| <i>Buteo albonotatus</i> Kaup, 1847                    | Zone-tailed Hawk            | R                   | LC; LC                        |            | WA 2829659 |                            |
| <b>GRUIFORMES</b>                                      |                             |                     |                               |            |            |                            |
| <b>Aramidae</b>  |                             |                     |                               |            |            |                            |
| <i>Aramus guarauna</i><br>(Linnaeus, 1766)             | Limpkin                     | R                   | LC; LC                        |            | WA 198235  | WA 3066368                 |
| <i>A. g. guarauna</i> (Linnaeus, 1766)*                |                             |                     |                               |            |            |                            |
| <b>Rallidae</b>  |                             |                     |                               |            |            |                            |
| <i>Rallus longirostris</i><br>Boddaert, 1783           | Mangrove Rail               | R                   | LC; LC                        |            | WA 322480  | WA 3300848                 |
| <i>R. l. crassirostris</i> Lawrence, 1871*             |                             |                     |                               |            |            |                            |
| <i>Aramides mangle</i> (Spix, 1825)                    | Little Wood Rail            | R                   | LC; LC                        |            | WA 385434  | Minns <i>et al.</i> (2010) |
| <i>Aramides cajaneus</i><br>(Statius Muller, 1776)     | Grey-necked Wood Rail       | R                   | LC; LC                        |            | WA 2574570 | WA 912029                  |
| <i>A. c. cajaneus</i> (Statius Muller, 1776)*          |                             |                     |                               |            |            |                            |
| <i>Amaurolimnas concolor</i><br>(Gosse, 1847)          | Uniform Crake               | R                   | LC; LC                        |            |            | WA 2117154                 |
| <i>A. c. castaneus</i> (Pucheran, 1851)*               |                             |                     |                               |            |            |                            |
| <i>Laterallus viridis</i><br>(Statius Muller, 1776)    | Russet-crowned Crake        | R                   | LC; LC                        |            | WA 2689621 | WA 1994114                 |
| <i>L. v. viridis</i> (Statius Muller, 1776)*           |                             |                     |                               |            |            |                            |
| <i>Laterallus melanophaius</i><br>(Vieillot, 1819)     | Rufous-sided Crake          | R                   | LC; LC                        |            | WA 2832972 | WA 3300851                 |
| <i>L. m. melanophaius</i> (Vieillot, 1819)*            |                             |                     |                               |            |            |                            |

| Taxon  | English name           | National occurrence | National; Global conservation | Specimen    | Photo      | Sound      |
|--|------------------------|---------------------|-------------------------------|-------------|------------|------------|
| <i>Laterallus exilis</i> (Temminck, 1831)              | Grey-breasted Crake    | R                   | LC; LC                        |             |            | WA 2324424 |
| <i>Porzana flaviventer</i> (Boddaert, 1783)            | Yellow-breasted Crake  | R                   | DD; LC                        |             | WA 2336075 |            |
| <i>P. f. flaviventer</i> (Boddaert, 1783)*             |                        |                     |                               |             |            |            |
| <i>Mustelirallus albicollis</i> (Vieillot, 1819)       | Ash-throated Crake     | R                   | LC; LC                        |             | WA 2333492 | WA 3300909 |
| <i>M. a. albicollis</i> (Vieillot, 1819)*              |                        |                     |                               |             |            |            |
| <i>Neocrex erythrops</i> (P. L. Sclater, 1867)         | Paint-billed Crake     | R                   | LC; LC                        | COUFRN 1224 | WA 343447  | WA 2992265 |
| <i>N. e. olivascens</i> Chubb, 1917*                   |                        |                     |                               |             |            |            |
| <i>Pardirallus maculatus</i> (Boddaert, 1783)          | Spotted Rail           | R                   | LC; LC                        |             | WA 2408428 |            |
| <i>P. m. maculatus</i> (Boddaert, 1783)*               |                        |                     |                               |             |            |            |
| <i>Gallinula galeata</i> (M. H. C. Lichtenstein, 1818) | Common Gallinule       | R                   | LC; LC                        | UFPE 3869   | WA 295455  | WA 3009806 |
| <i>G. g. galeata</i> (M. H. C. Lichtenstein, 1818)*    |                        |                     |                               |             |            |            |
| <i>Porphyriops melanops</i> (Vieillot, 1819)           | Spot-flanked Gallinule | R                   | LC; LC                        |             | WA 3089884 |            |
| <i>P. m. melanops</i> (Vieillot, 1819)*                |                        |                     |                               |             |            |            |
| <i>Porphyrio martinicus</i> (Linnaeus, 1766)           | Purple Gallinule       | R                   | LC; LC                        | COUFRN 1191 | WA 344504  | WA 2117494 |
| <i>Porphyrio flavirostris</i> (J. F. Gmelin, 1789)     | Azure Gallinule        | R                   | LC; LC                        |             | WA 609215  |            |
| <b>Heliornithidae</b>                                  |                        |                     |                               |             |            |            |
| [ <i>Heliornis fulica</i> (Boddaert, 1783)]            | Sungrebe               | R                   | LC; LC                        |             |            |            |
| <b>Charadriidae</b>                                    |                        |                     |                               |             |            |            |
| <i>Vanellus cayanus</i> (Latham, 1790)                 | Pied Lapwing           | R                   | LC; LC                        | MNRJ 34431  | WA 1344089 |            |
| <i>Vanellus chilensis</i> (Molina, 1782)               | Southern Lapwing       | R                   | LC; LC                        | COUFRN 1230 | WA 249749  | WA 728917  |
| <i>V. c. lampronotus</i> (Wagler, 1827)*               |                        |                     |                               |             |            |            |
| <i>Pluvialis dominica</i> (Statius Muller, 1776)       | American Golden-Plover | VN                  | DD; LC                        |             | WA 893159  |            |
| <i>Pluvialis squatarola</i> (Linnaeus, 1758)           | Black-bellied Plover   | VN                  | LC; LC                        | UFPE 3568   | WA 1342260 | WA 2669029 |
| <i>P. s. cynosurae</i> (Thayer & Bangs, 1914)*         |                        |                     |                               |             |            |            |
| <i>Charadrius semipalmatus</i> Bonaparte, 1825         | Semipalmated Plover    | VN                  | LC; LC                        | UFPE 3138   | WA 1342258 | WA 2669026 |
| <i>Charadrius wilsonia</i> Ord, 1814                   | Wilson's Plover        | R                   | VU; LC                        |             | WA 250851  | WA 1254343 |
| <i>C. w. crassirostris</i> Spix, 1825*                 |                        |                     | VU                            |             |            |            |
| <i>Charadrius collaris</i> Vieillot, 1818              | Collared Plover        | R                   | LC; LC                        | UFPE 3964   | WA 250844  |            |

| Taxon   | English name           | National occurrence | National; Global conservation | Specimen  | Photo      | Sound                      |
|---|------------------------|---------------------|-------------------------------|-----------|------------|----------------------------|
| <b>Haematopodidae</b>                                 |                        |                     |                               |           |            |                            |
| <i>Haematopus palliatus</i><br>Temminck, 1820         | American Oystercatcher | R                   | NT; LC                        |           | WA 253376  | WA 1433694                 |
| <i>H. p. palliatus</i> Temminck, 1820                 |                        |                     |                               |           |            |                            |
| <b>Recurvirostridae</b>                               |                        |                     |                               |           |            |                            |
| <i>Himantopus mexicanus</i><br>(Statius Muller, 1776) | Black-necked Stilt     | R                   | LC; NE                        | UFPE 3975 | WA 339866  | Minns <i>et al.</i> (2010) |
| <b>Scolopacidae</b>                                   |                        |                     |                               |           |            |                            |
| <i>Gallinago paraguayae</i><br>(Vieillot, 1816)       | South American Snipe   | R                   | LC; LC                        |           | WA 3065363 | WA 2179162                 |
| <i>Gallinago p. paraguayae</i> (Vieillot, 1816)*      |                        |                     |                               |           |            |                            |
| <i>Limnodromus griseus</i><br>(J. F. Gmelin, 1789)    | Short-billed Dowitcher | VN                  | CR; LC                        |           | WA 247845  | WA 2669028                 |
| <i>L. g. griseus</i> (J. F. Gmelin 1789)*             |                        |                     |                               |           |            |                            |
| [ <i>Limosa haemastica</i><br>(Linnaeus, 1758)]       | Hudsonian Godwit       | VN                  | LC; LC                        |           |            |                            |
| <i>Limosa lapponica</i><br>(Linnaeus, 1758)           | Bar-tailed Godwit      | VA(N)               | NE; NT                        |           | WA 3089895 |                            |
| <i>L. l. lapponica</i> (Linnaeus, 1758)*              |                        |                     |                               |           |            |                            |
| <i>Numenius hudsonicus</i><br>Latham, 1790            | American Whimbrel      | VN                  | NT; NE                        | UFPE 3673 | WA 3089895 | WA 1433695                 |
| <i>N. h. hudsonicus</i> Latham, 1790*                 |                        |                     |                               |           |            |                            |
| [ <i>Numenius phaeopus</i><br>(Linnaeus, 1758)]       | Eurasian Whimbrel      | VA(N)               | NE; LC                        |           |            |                            |
| <i>Actitis macularius</i><br>(Linnaeus, 1766)         | Spotted Sandpiper      | VN                  | LC; LC                        |           | WA 2655752 | WA 2581737                 |
| <i>Tringa solitaria</i> Wilson, 1813                  | Solitary Sandpiper     | VN                  | LC; LC                        |           | WA 579049  |                            |
| <i>T. s. solitaria</i> Wilson, 1813*                  |                        |                     |                               |           |            |                            |
| <i>Tringa melanoleuca</i><br>(J. F. Gmelin, 1789)     | Greater Yellowlegs     | VN                  | LC; LC                        |           | WA 198284  | WA 1433699                 |
| <i>Tringa semipalmata</i><br>(J. F. Gmelin, 1789)     | Willet                 | VN                  | LC; LC                        |           | WA 1522100 | WA 2581745                 |
| <i>T. s. semipalmata</i> (J. F. Gmelin, 1789)         |                        |                     |                               |           |            |                            |
| <i>Tringa flavipes</i><br>(J. F. Gmelin, 1789)        | Lesser Yellowlegs      | VN                  | LC; LC                        | UFPE 3596 | WA 1367974 | WA 1433697                 |
| <i>Arenaria interpres</i><br>(Linnaeus, 1758)         | Ruddy Turnstone        | VN                  | NT; LC                        | UFPE 3569 | WA 1342259 | WA 2669025                 |
| <i>A. i. morinella</i> (Linnaeus, 1766)*              |                        |                     |                               |           |            |                            |
| <i>Calidris canutus</i><br>(Linnaeus, 1758)           | Red Knot               | VN                  | CR; NT                        | UFPE 3595 | WA 2040564 | WA 1450088                 |
| <i>C. c. rufa</i> (Wilson, 1813)*                     |                        |                     |                               |           |            |                            |
| <i>Calidris alba</i> (Pallas, 1764)                   | Sanderling             | VN                  | LC; LC                        | UFPE 3961 | WA 2655751 | WA 1433693                 |
| <i>C. a. rubida</i> (J. F. Gmelin, 1789)*             |                        |                     |                               |           |            |                            |
| <i>Calidris pusilla</i> (Linnaeus, 1766)              | Semipalmated Sandpiper | VN                  | EN; NT                        | UFPE 3067 | WA 2655753 | WA 2581740                 |

| Taxon   | English name           | National occurrence | National; Global conservation | Specimen  | Photo      | Sound      |
|---|------------------------|---------------------|-------------------------------|-----------|------------|------------|
| <i>Calidris minutilla</i> (Vieillot, 1819)            | Least Sandpiper        | VN                  | DD; LC                        | UFPE 3959 | WA 108627  |            |
| <i>Calidris fuscicollis</i> (Vieillot, 1819)          | White-rumped Sandpiper | VN                  | LC; LC                        | UFPE 3657 | WA 636372  |            |
| <i>Calidris melanotos</i> (Vieillot, 1819)            | Pectoral Sandpiper     | VN                  | LC; LC                        |           | ML27817531 |            |
| <i>Calidris himantopus</i> (Bonaparte, 1826)          | Stilt Sandpiper        | VN                  | LC; LC                        |           | WA 95041   |            |
| <b>Jacaniidae</b>                                     |                        |                     |                               |           |            |            |
| <i>Jacana jacana</i> (Linnaeus, 1766)                 | Wattled Jacana         | R                   | LC; LC                        |           | WA 2722891 |            |
| <i>J. j. jacana</i> (Linnaeus, 1766)*                 |                        |                     |                               |           |            |            |
| <b>Stercorariidae</b>                                 |                        |                     |                               |           |            |            |
| <i>Stercorarius maccormicki</i> Saunders, 1893        | South Polar Skua       | VS                  | LC; LC                        |           | WA 791473  |            |
| <i>Stercorarius pomarinus</i> (Temminck, 1815)        | Pomarine Jaeger        | VN                  | LC; LC                        |           | WA 1593176 |            |
| <i>Stercorarius parasiticus</i> (Linnaeus, 1758)      | Parasitic Jaeger       | VN                  | LC; LC                        |           | ML77958391 |            |
| <b>Laridae</b>  |                        |                     |                               |           |            |            |
| <i>Chroicocephalus cirrocephalus</i> (Vieillot, 1818) | Grey-hooded Gull       | R                   | LC; LC                        | UFPE 3070 | WA 322479  | WA 1433713 |
| <i>C. c. cirrocephalus</i> (Vieillot, 1818)*          |                        |                     |                               |           |            |            |
| <i>Leucophaeus atricilla</i> (Linnaeus, 1758)         | Laughing Gull          | VN                  | LC; LC                        |           | WA 287755  |            |
| <i>L. a. atricilla</i> (Linnaeus, 1758)*              |                        |                     |                               |           |            |            |
| <i>Larus dominicanus</i> M. H. C. Lichtenstein, 1823  | Kelp Gull              | R                   | LC; LC                        |           | WA 2016695 |            |
| <i>L. d. dominicanus</i> M. H. C. Lichtenstein, 1823* |                        |                     |                               |           |            |            |
| <b>Sternidae</b>                                      |                        |                     |                               |           |            |            |
| <i>Anous stolidus</i> (Linnaeus, 1758)                | Brown Noddy            | R                   | LC; LC                        |           | WA 1728204 |            |
| <i>A. s. stolidus</i> (Linnaeus, 1758)*               |                        |                     |                               |           |            |            |
| <i>Onychoprion fuscatus</i> (Linnaeus, 1766)          | Sooty Tern             | R                   | LC; LC                        |           | WA 1593177 |            |
| <i>O. f. fuscatus</i> (Linnaeus, 1766)*               |                        |                     |                               |           |            |            |
| <i>Sternula antillarum</i> Lesson, 1847               | Least Tern             | R                   | LC; LC                        | UFPE 3139 | WA 1359913 | WA 3300839 |
| <i>S. a. antillarum</i> Lesson, 1847*                 |                        |                     |                               |           |            |            |
| <i>Sternula supercilii</i> (Vieillot, 1819)           | Yellow-billed Tern     | R                   | LC; LC                        |           | WA 611256  |            |
| <i>Phaetusa simplex</i> (J. F. Gmelin, 1789)          | Large-billed Tern      | R                   | LC; LC                        |           | WA 2503157 |            |
| <i>P. s. simplex</i> (J. F. Gmelin, 1789)*            |                        |                     |                               |           |            |            |



| Taxon  | English name               | National occurrence | National; Global conservation | Specimen    | Photo      | Sound                      |
|--|----------------------------|---------------------|-------------------------------|-------------|------------|----------------------------|
| <i>Gelochelidon nilotica</i><br>(J. F. Gmelin, 1789)   | Gull-billed Tern           | R                   | LC; LC                        | UFPE 3072   | WA 11254   | WA 1433714                 |
| <i>G. n. gronvoldi</i> (Mathews, 1912)*                |                            |                     |                               |             |            |                            |
| [ <i>Chlidonias niger</i><br>(Linnaeus, 1758)]         | Black Tern                 | VN                  | NE; LC                        |             |            |                            |
| <i>Sterna hirundo</i> Linnaeus, 1758                   | Common Tern                | VN                  | LC; LC                        |             | WA 11253   |                            |
| <i>S. h. hirundo</i> Linnaeus, 1758*                   |                            |                     |                               |             |            |                            |
| <i>Sterna dougallii</i> Montagu, 1813                  | Roseate Tern               | VN                  | VU; LC                        |             | WA 1359911 |                            |
| <i>S. d. dougallii</i> Montagu, 1813*                  |                            |                     |                               |             |            |                            |
| <i>Sterna paradisaea</i><br>Pontoppidan, 1763          | Arctic Tern                | VN                  | LC; LC                        |             | WA 612272  |                            |
| <i>Thalasseus acuflavidus</i><br>(Cabot, 1847)         | Cabot's Tern               | R                   | LC; LC                        |             | WA 236785  |                            |
| <i>T. a. acuflavidus</i> (Cabot, 1847)*                |                            |                     |                               |             |            |                            |
| <i>T. a. eurygnathus</i> (Saunders, 1876)              |                            |                     |                               |             |            |                            |
| <b>Rynchopidae</b>                                     |                            |                     |                               |             |            |                            |
| <i>Rynchops niger</i> Linnaeus, 1758                   | Black Skimmer              | R                   | LC; LC                        |             | WA 283229  | WA 2987184                 |
| <i>R. n. intercedens</i> Saunders, 1895*               |                            |                     |                               |             |            |                            |
| <b>COLUMBIFORMES</b>                                   |                            |                     |                               |             |            |                            |
| <b>Columbidae</b>                                      |                            |                     |                               |             |            |                            |
| <i>Columbina passerina</i><br>(Linnaeus, 1758)         | Common Ground Dove         | R                   | LC; LC                        | UFPE 3865   | WA 241789  | XC 430053                  |
| <i>C. p. griseola</i> Spix, 1825*                      |                            |                     |                               |             |            |                            |
| <i>Columbina minuta</i><br>(Linnaeus, 1766)            | Plain-breasted Ground Dove | R                   | LC; LC                        | COUFRN 75   | WA 347676  | WA 1254340                 |
| <i>C. m. minuta</i> (Linnaeus, 1766)*                  |                            |                     |                               |             |            |                            |
| <i>Columbina talpacoti</i><br>(Temminck, 1810)         | Ruddy Ground Dove          | R                   | LC; LC                        | COUFRN 782  | WA 354629  | WA 3300854                 |
| <i>C. t. talpacoti</i> (Temminck, 1810)*               |                            |                     |                               |             |            |                            |
| <i>Columbina squammata</i><br>(Lesson, 1831)           | Scaled Dove                | R                   | LC; LC                        | COUFRN 1206 | WA 357485  | Minns <i>et al.</i> (2010) |
| <i>C. s. squammata</i> (Lesson, 1831)*                 |                            |                     |                               |             |            |                            |
| <i>Columbina picui</i><br>(Temminck, 1813)             | Picui Ground Dove          | R                   | LC; LC                        | COUFRN 2    | WA 47101   | WA 1254504                 |
| <i>C. p. strepitans</i> Spix, 1825*                    |                            |                     |                               |             |            |                            |
| <i>Claravis pretiosa</i><br>(Ferrari-Perez, 1886)      | Blue Ground Dove           | R                   | LC; LC                        | COUFRN 953  | WA 2557623 | WA 2557967                 |
| <i>Columba livia</i> J. F. Gmelin, 1789                | Rock Pigeon                | R                   | NE; LC                        |             | WA 724444  | WA 3003975                 |
| <i>C. l. livia</i> J. F. Gmelin, 1789*                 |                            |                     |                               |             |            |                            |
| [ <i>Patagioenas speciosa</i><br>(J. F. Gmelin, 1789)] | Scaled Pigeon              | R                   | LC; LC                        |             |            |                            |
| <i>Patagioenas picazuro</i><br>(Temminck, 1813)        | Picazuro Pigeon            | R                   | LC; LC                        |             | WA 279033  |                            |
| <i>P. p. marginalis</i> (Naumburg, 1932)*              |                            |                     |                               |             |            |                            |

| Taxon  | English name           | National occurrence | National; Global conservation | Specimen              | Photo      | Sound                      |
|--|------------------------|---------------------|-------------------------------|-----------------------|------------|----------------------------|
| <i>Patagioenas cayennensis</i> (Bonnaterre, 1792)      | Pale-vented Pigeon     | R                   | LC; LC                        |                       | WA 1615429 | WA 2673184                 |
| <i>P. c. sylvestris</i> (Vieillot, 1818)*              |                        |                     |                               |                       |            |                            |
| <i>Zenaidura auriculata</i> (Des Murs, 1847)           | Eared Dove             | R                   | LC; LC                        | MNRJ 36288            | WA 342788  | WA 2928515                 |
| <i>Z. a. noronha</i> Sharpe, 1890*                     |                        |                     |                               |                       |            |                            |
| <i>Leptotila verreauxi</i> Bonaparte, 1855             | White-tipped Dove      | R                   | LC; LC                        | COUFRN 803            | WA 346986  | WA 3865184                 |
| <i>L. v. approximans</i> Cory, 1917*                   |                        |                     |                               |                       |            |                            |
| <i>Leptotila rufaxilla</i> (Richard & Bernard, 1792)   | Grey-fronted Dove      | R                   | LC; LC                        |                       |            | WA 2569992                 |
| <i>L. r. bahiae</i> Berlepsch, 1885*                   |                        |                     |                               |                       |            |                            |
| [ <i>Geotrygon montana</i> (Linnaeus, 1758)]           | Ruddy Quail-Dove       | R                   | LC; LC                        |                       |            |                            |
| <b>CUCULIFORMES</b>                                    |                        |                     |                               |                       |            |                            |
| <b>Cuculidae</b>                                       |                        |                     |                               |                       |            |                            |
| <i>Micrococcyx cinereus</i> (Vieillot, 1817)           | Ash-coloured Cuckoo    | R                   | LC; LC                        |                       | WA 1716483 |                            |
| <i>Piaya cayana</i> (Linnaeus, 1766)                   | Squirrel Cuckoo        | R                   | LC; LC                        | COUFRN 179            | WA 249743  | WA 3300857                 |
| <i>P. c. pallescens</i> (Cabanis & Heine, 1863)*       |                        |                     |                               |                       |            |                            |
| <i>Coccyzus melacoryphus</i> Vieillot, 1817            | Dark-billed Cuckoo     | R                   | LC; LC                        | COUFRN 837            | WA 344900  | WA 3300857                 |
| <i>Coccyzus americanus</i> (Linnaeus, 1758)            | Yellow-billed Cuckoo   | VN                  | LC; LC                        | COUFRN 131            | WA 2499860 |                            |
| <i>C. a. americanus</i> (Linnaeus, 1758)*              |                        |                     |                               |                       |            |                            |
| <i>Coccyzus euleri</i> Cabanis, 1873                   | Pearly-breasted Cuckoo | R                   | LC; LC                        |                       | WA 249004  |                            |
| <i>Crotophaga major</i> J. F. Gmelin, 1788             | Greater Ani            | R                   | LC; LC                        |                       | WA 249751  | WA 3300858                 |
| <i>Crotophaga ani</i> Linnaeus, 1758                   | Smooth-billed Ani      | R                   | LC; LC                        | COUFRN 87             | WA 249750  | WA 1254713                 |
| <i>Guirra guirra</i> (J. F. Gmelin, 1788)              | Guirra Cuckoo          | R                   | LC; LC                        | COUFRN 618            | WA 725015  | Minns <i>et al.</i> (2010) |
| <i>Tapera naevia</i> (Linnaeus, 1766)                  | Striped Cuckoo         | R                   | LC; LC                        |                       | WA 251688  | WA 3300859                 |
| <i>T. n. naevia</i> (Linnaeus, 1766)*                  |                        |                     |                               |                       |            |                            |
| <b>STRIGIFORMES</b>                                    |                        |                     |                               |                       |            |                            |
| <b>Tytonidae</b>                                       |                        |                     |                               |                       |            |                            |
| <i>Tyto furcata</i> (Temminck, 1827)                   | American Barn Owl      | R                   | LC; NE                        | CPPD (not catalogued) | WA 2016708 | WA 2926836 (Online)        |
| <i>T. f. tuidara</i> (J. E. Gray, 1828)*               |                        |                     |                               |                       |            |                            |
| <b>Strigidae</b>                                       |                        |                     |                               |                       |            |                            |
| <i>Megascops choliba</i> (Vieillot, 1817)              | Tropical Screech Owl   | R                   | LC; LC                        | MNRJ 36295            | WA 344491  | WA 488604                  |
| <i>M. c. decussatus</i> (M. H. C. Lichtenstein, 1823)* |                        |                     |                               |                       |            |                            |

| Taxon  | English name          | National occurrence | National; Global conservation | Specimen              | Photo      | Sound      |
|--|-----------------------|---------------------|-------------------------------|-----------------------|------------|------------|
| [ <i>Pulsatrix perspicillata</i> (Latham, 1790)]   | Spectacled Owl        | R                   | LC; LC                        |                       |            |            |
| <i>Bubo virginianus</i> (J. F. Gmelin, 1788)       | Great Horned Owl      | R                   | LC; LC                        |                       | WA 2853518 | WA 316723  |
| <i>B. v. deserti</i> Reiser, 1905*                 |                       |                     |                               |                       |            |            |
| [ <i>Strix virgata</i> (Cassin, 1849)]             | Mottled Owl           | R                   | LC; LC                        |                       |            |            |
| <i>Glaucidium brasilianum</i> (J. F. Gmelin, 1788) | Ferruginous Pygmy Owl | R                   | LC; LC                        | MNRJ 34465            | WA 87779   | WA 257119  |
| <i>G. b. brasilianum</i> (J. F. Gmelin, 1788)*     |                       |                     |                               |                       |            |            |
| <i>Athene cunicularia</i> (Molina, 1782)           | Burrowing Owl         | R                   | LC; LC                        | COUFRN 1218           | WA 37073   | WA 2577098 |
| <i>A. c. grallaria</i> (Temminck, 1822)*           |                       |                     |                               |                       |            |            |
| [ <i>Aegolius harrisii</i> (Cassin, 1849)]         | Buff-fronted Owl      | R                   | LC; LC                        |                       |            |            |
| <i>Asio clamator</i> (Vieillot, 1808)              | Striped Owl           | R                   | LC; LC                        | CPPD (not catalogued) | WA 2035087 |            |
| <i>A. c. clamator</i> (Vieillot, 1808)*            |                       |                     |                               |                       |            |            |
| <i>Asio flammeus</i> (Pontoppidan, 1763)           | Short-eared Owl       | R                   | LC; LC                        |                       | WA 3065349 |            |
| <i>A. f. suinda</i> (Vieillot, 1817)*              |                       |                     |                               |                       |            |            |

## NYCTIBIIFORMES

### Nyctibiidae

|   |              |   |        |  |            |            |
|---|--------------|---|--------|--|------------|------------|
| <i>Nyctibius griseus</i> (J. F. Gmelin, 1789) | Common Potoo | R | LC; LC |  | WA 1361294 | WA 2016103 |
| <i>N. g. griseus</i> (J. F. Gmelin, 1789)*    |              |   |        |  |            |            |

### Caprimulgidae

|   |                        |      |        |            |            |            |
|---|------------------------|------|--------|------------|------------|------------|
| <i>Antrostomus rufus</i> (Boddaert, 1783)           | Rufous Nightjar        | R    | LC; LC |            |            | XC 430029  |
| <i>A. r. rufus</i> (Boddaert, 1783)*                |                        |      |        |            |            |            |
| <i>Lurocalis semitorquatus</i> (J. F. Gmelin, 1789) | Short-tailed Nighthawk | R    | LC; LC |            | WA 2615000 |            |
| <i>L. s. nattereri</i> (Temminck, 1822)*            |                        |      |        |            |            |            |
| <i>Nyctidromus albicollis</i> (J. F. Gmelin, 1789)  | Common Pauraque        | R    | LC; LC | COUFRN 644 | WA 1270179 | WA 3300862 |
| <i>N. a. albicollis</i> (J. F. Gmelin, 1789)*       |                        |      |        |            |            |            |
| <i>Nyctidromus hirundinaceus</i> (Spix, 1825)       | Pygmy Nightjar         | R, E | DD; LC | MNRJ 34469 | WA 1359959 | WA 3300864 |
| <i>N. h. cearae</i> (Cory, 1917)*                   |                        |      |        |            |            |            |
| <i>Hydropsalis parvula</i> (Gould, 1837)            | Little Nightjar        | R    | LC; LC | UFPE 3674  | WA 864255  | WA 3300866 |
| <i>Hydropsalis longirostris</i> (Bonaparte, 1825)   | Band-winged Nightjar   | R    | LC; LC |            | WA 3219414 | WA 3220109 |
| <i>H. l. longirostris</i> (Bonaparte, 1825)*        |                        |      |        |            |            |            |

| Taxon   | English name               | National occurrence | National; Global conservation | Specimen    | Photo      | Sound      |
|---|----------------------------|---------------------|-------------------------------|-------------|------------|------------|
| <i>Hydropsalis torquata</i><br>(J. F. Gmelin, 1789)     | Scissor-tailed Nightjar    | R                   | LC; LC                        | COUFRN 69   | WA 317704  |            |
| <i>H. t. torquata</i> (J. F. Gmelin, 1789)*             |                            |                     |                               |             |            |            |
| <i>Nannochordeiles pusillus</i><br>(Gould, 1861)        | Least Nighthawk            | R                   | LC; LC                        | MNRJ 34468  | WA 1355616 |            |
| <i>N. p. xerophilus</i> (Dickerman, 1988)*              |                            |                     |                               |             |            |            |
| <i>Podager nacunda</i> (Vieillot, 1817)                 | Nacunda Nighthawk          | R                   | LC; LC                        | COUFRN 1228 | WA 2985934 |            |
| <i>P. n. nacunda</i> (Vieillot, 1817)*                  |                            |                     |                               |             |            |            |
| [ <i>Chordeiles minor</i><br>(J.R. Forster, 1771)]      | Common Nighthawk           | VN                  | LC; LC                        |             |            |            |
| <i>Chordeiles acutipennis</i><br>(Hermann, 1783)        | Lesser Nighthawk           | R                   | LC; LC                        |             | WA 3089210 |            |
| <i>C. a. acutipennis</i> (Hermann, 1783)*               |                            |                     |                               |             |            |            |
| <b>APODIFORMES</b>                                      |                            |                     |                               |             |            |            |
| <b>Apodidae</b>   |                            |                     |                               |             |            |            |
| <i>Streptoprocne biscutata</i><br>(P. L. Sclater, 1866) | Biscutate Swift            | R                   | LC; LC                        | MNRJ 36897  | WA 313921  | WA 3300867 |
| <i>S. b. seridoensis</i> Sick, 1991                     |                            |                     |                               |             |            |            |
| [ <i>Chaetura meridionalis</i><br>Hellmayr, 1907]       | Sick's Swift               | R                   | LC; LC                        |             |            |            |
| <i>Tachornis squamata</i><br>(Cassin, 1853)             | Fork-tailed Palm Swift     | R                   | LC; LC                        |             | WA 238953  | WA 3184658 |
| <i>T. s. squamata</i> (Cassin, 1853)*                   |                            |                     |                               |             |            |            |
| <b>Trochilidae</b>                                      |                            |                     |                               |             |            |            |
| <i>Glaucis hirsutus</i> (J. F. Gmelin, 1788)            | Rufous-breasted Hermit     | R                   | LC; LC                        |             | WA 2688509 |            |
| <i>G. h. hirsutus</i> (J. F. Gmelin, 1788)*             |                            |                     |                               |             |            |            |
| <i>Anopetia gounellei</i><br>(Boucard, 1891)            | Broad-tipped Hermit        | R, E                | LC; LC                        | COUFRN 627  | WA 347675  | WA 2967669 |
| <i>Phaethornis ruber</i><br>(Linnaeus, 1758)            | Reddish Hermit             | R                   | LC; LC                        | COUFRN 508  | WA 244642  | XC 430053  |
| <i>P. r. pygmaeus</i> (Spix, 1825)*                     |                            |                     |                               |             |            |            |
| <i>Phaethornis pretrei</i><br>(Lesson & Delattre, 1839) | Planalto Hermit            | R                   | LC; LC                        | COUFRN 506  | WA 2670787 |            |
| <i>Eupetomena macroura</i><br>(J. F. Gmelin, 1788)      | Swallow-tailed Hummingbird | R                   | LC; LC                        | COUFRN 726  | WA 631517  | XC 430036  |
| <i>E. m. simoni</i> Hellmayr, 1929*                     |                            |                     |                               |             |            |            |
| <i>Anthracothorax nigricollis</i><br>(Vieillot, 1817)   | Black-throated Mango       | R                   | LC; LC                        |             | WA 2929048 |            |
| <i>Chrysolampis mosquitos</i><br>(Linnaeus, 1758)       | Ruby-topaz Hummingbird     | R                   | LC; LC                        | UFPE 3446   | WA 239581  |            |
| <i>Chlorestes notata</i> (Reich, 1793)                  | Blue-chinned Sapphire      | R                   | LC; LC                        | COUFRN 686  | WA 1151475 |            |
| <i>C. n. notata</i> (Reich, 1793)*                      |                            |                     |                               |             |            |            |

| Taxon   | English name                | National occurrence | National; Global conservation | Specimen   | Photo      | Sound      |
|---|-----------------------------|---------------------|-------------------------------|------------|------------|------------|
| <i>Chlorostilbon lucidus</i><br>(Shaw, 1812)        | Glittering-bellied Emerald  | R                   | LC; LC                        | COUFRN 732 | WA 2715988 |            |
| <i>C. l. pucherani</i> (Bourcier & Mulsant, 1848)*  |                             |                     |                               |            |            |            |
| [ <i>Hylocharis cyanus</i><br>(Vieillot, 1818)]     | White-chinned Sapphire      | R                   | LC; LC                        |            |            |            |
| <i>Polytmus guainumbi</i><br>(Pallas, 1764)         | White-tailed Goldenthrout   | R                   | LC; LC                        | UFPE 3690  | WA 623011  |            |
| <i>P. g. thaumantias</i> (Linnaeus, 1766)*          |                             |                     |                               |            |            |            |
| <i>Amazilia leucogaster</i><br>(J. F. Gmelin, 1788) | Plain-bellied Emerald       | R                   | LC; LC                        | COUFRN 255 | WA 817119  | WA 2819810 |
| <i>A. l. bahiae</i> (E. Hartert, 1899)*             |                             |                     |                               |            |            |            |
| [ <i>Amazilia versicolor</i><br>(Vieillot, 1818)]   | Versicoloured Emerald       | R                   | LC; LC                        |            |            |            |
| <i>Amazilia fimbriata</i><br>(J.F. Gmelin, 1788)    | Glittering-throated Emerald | R                   | LC; LC                        | COUFRN 779 | WA 480164  |            |
| <i>A. f. nigricauda</i> (Elliot, 1878)*             |                             |                     |                               |            |            |            |
| <i>Helictin bilophus</i><br>(Temminck, 1820)        | Horned Sungem               | R                   | LC; LC                        |            | WA 2682924 | WA 2984447 |
| <i>Helimaster squamosus</i><br>(Temminck, 1823)     | Stripe-breasted Starthroat  | R, E                | LC; LC                        |            | WA 534458  |            |
| <i>Calliphlox amethystina</i><br>(Boddaert, 1783)   | Amethyst Woodstar           | R                   | LC; LC                        |            | WA 2472156 |            |

## TROGONIFORMES

### Trogonidae

|   |                     |   |        |            |           |                               |
|---|---------------------|---|--------|------------|-----------|-------------------------------|
| <i>Trogon curucui</i><br>Linnaeus, 1766 | Blue-crowned Trogon | R | LC; LC | COUFRN 498 | WA 250915 | Minns <i>et al.</i><br>(2010) |
| <i>T. c. curucui</i> Linnaeus, 1766*    |                     |   |        |            |           |                               |

## CORACIIFORMES

### Alcedinidae

|   |                   |   |        |  |            |            |
|---|-------------------|---|--------|--|------------|------------|
| <i>Megaceryle torquata</i><br>(Linnaeus, 1766)        | Ringed Kingfisher | R | LC; LC |  | WA 2715841 | WA 2895442 |
| <i>M. t. torquata</i> (Linnaeus, 1766)*               |                   |   |        |  |            |            |
| <i>Chloroceryle amazona</i><br>(Latham, 1790)         | Amazon Kingfisher | R | LC; LC |  | WA 615812  | WA 3039036 |
| <i>Chloroceryle americana</i><br>(J. F. Gmelin, 1788) | Green Kingfisher  | R | LC; LC |  | WA 250857  |            |
| <i>C. a. americana</i> (J. F. Gmelin, 1788)*          |                   |   |        |  |            |            |

### Momotidae

|   |                  |   |        |  |            |  |
|---|------------------|---|--------|--|------------|--|
| <i>Momotus momota</i><br>(Linnaeus, 1766)         | Amazonian Motmot | R | LC; LC |  | WA 2510629 |  |
| <i>M. m. marcgravianus</i> Pinto & Camargo, 1961* |                  |   | EN     |  |            |  |



| Taxon  | English name               | National occurrence | National; Global conservation | Specimen   | Photo      | Sound                      |
|--|----------------------------|---------------------|-------------------------------|------------|------------|----------------------------|
| <b>GALBULIFORMES</b>                                 |                            |                     |                               |            |            |                            |
| <b>Galbulidae</b>                                    |                            |                     |                               |            |            |                            |
| <i>Galbula ruficauda</i> Cuvier, 1816                | Rufous-tailed Jacamar      | R                   | LC; LC                        | COUFRN 791 | WA 244631  | WA 3300873                 |
| <i>G. r. rufoviridis</i> Cabanis, 1851               |                            |                     |                               |            |            |                            |
| <b>Bucconidae</b>                                    |                            |                     |                               |            |            |                            |
| <i>Nystalus maculatus</i> (J. F. Gmelin, 1788)       | Spot-backed Puffbird       | R, E                | LC; LC                        | COUFRN 141 | WA 1365618 | Minns <i>et al.</i> (2010) |
| <b>PICIFORMES</b>                                    |                            |                     |                               |            |            |                            |
| <b>Picidae</b>                                       |                            |                     |                               |            |            |                            |
| <i>Picumnus fulvescens</i> Stager, 1961              | Tawny Piculet              | R, E                | LC; NT                        | COUFRN 47  | WA 1343769 |                            |
| <i>Picumnus liniae</i> Snethlage, 1924               | Ochraceous Piculet         | R, E                | LC; LC                        | CAHZ 705   | WA 235428  | WA 253695                  |
| <i>Melanerpes candidus</i> (Otto, 1796)              | White Woodpecker           | R                   | LC; LC                        |            | WA 342790  | XC 206440                  |
| <i>Veniliornis passerinus</i> (Linnaeus, 1766)       | Little Woodpecker          | R                   | LC; LC                        | COUFRN 799 | WA 248472  | Minns <i>et al.</i> (2010) |
| <i>V. p. taenionotus</i> (Reichenbach, 1854)*        |                            |                     |                               |            |            |                            |
| <i>Piculus chrysochloros</i> (Vieillot, 1818)        | Golden-green Woodpecker    | R                   | LC; LC                        |            | WA 34096   |                            |
| <i>Colaptes melanochloros</i> (J. F. Gmelin, 1788)   | Green-barred Woodpecker    | R                   | LC; LC                        | CAHZ 692   | WA 47058   | WA 1254721                 |
| <i>C. m. nattereri</i> (Malherbe, 1845)*             |                            |                     |                               |            |            |                            |
| <i>Colaptes campestris</i> (Vieillot, 1818)          | Campo Flicker              | R                   | LC; LC                        |            | WA 2221867 | WA 2886788                 |
| <i>C. c. campestris</i> (Vieillot, 1818)             |                            |                     |                               |            |            |                            |
| <i>Celeus ochraceus</i> (Spix, 1824)                 | Ochre-backed Woodpecker    | R, E                | LC; LC                        |            | WA 1441573 | WA 3300878                 |
| <i>Dryocopus lineatus</i> (Linnaeus, 1766)           | Lineated Woodpecker        | R                   | LC; LC                        |            | WA 1609673 |                            |
| <i>D. l. lineatus</i> (Linnaeus, 1766)*              |                            |                     |                               |            |            |                            |
| <i>Campephilus melanoleucos</i> (J. F. Gmelin, 1788) | Crimson-crested Woodpecker | R                   | LC; LC                        |            | WA 239452  |                            |
| <i>C. m. cearae</i> (Cory, 1915)*                    |                            |                     |                               |            |            |                            |
| <b>CARIAMIFORMES</b>                                 |                            |                     |                               |            |            |                            |
| <b>Cariamidae</b>                                    |                            |                     |                               |            |            |                            |
| <i>Cariama cristata</i> (Linnaeus, 1766)             | Red-legged Seriema         | R                   | LC; LC                        |            | WA 249742  | WA 2088256                 |
| <b>FALCONIFORMES</b>                                 |                            |                     |                               |            |            |                            |
| <b>Falconidae</b>                                    |                            |                     |                               |            |            |                            |
| <i>Caracara plancus</i> (J. F. Miller, 1777)         | Southern Caracara          | R                   | LC; LC                        |            | WA 6413    | WA 3300879                 |
| <i>Milvago chimachima</i> (Vieillot, 1816)           | Yellow-headed Caracara     | R                   | LC; LC                        |            | WA 287762  | WA 3300845                 |
| <i>M. c. chimachima</i> (Vieillot, 1816)*            |                            |                     |                               |            |            |                            |

| Taxon  | English name           | National occurrence | National; Global conservation | Specimen | Photo     | Sound      |
|--|------------------------|---------------------|-------------------------------|----------|-----------|------------|
| <i>Herpetotheres cachinnans</i> (Linnaeus, 1758) | Laughing Falcon        | R                   | LC; LC                        |          | WA 348725 | WA 1736786 |
| <i>H. c. cachinnans</i> (Linnaeus, 1758)*        |                        |                     |                               |          |           |            |
| [ <i>Microstur ruficollis</i> (Vieillot, 1817)]  | Barred Forest Falcon   | R                   | LC; LC                        |          |           |            |
| <i>Microstur semitorquatus</i> (Vieillot, 1817)  | Collared Forest Falcon | R                   | LC; LC                        |          |           | WA 3390273 |
| <i>M. s. semitorquatus</i> (Vieillot, 1817)*     |                        |                     |                               |          |           |            |
| <i>Falco sparverius</i> Linnaeus, 1758           | American Kestrel       | R                   | LC; LC                        |          | WA 80474  |            |
| <i>F. s. cearae</i> (Cory, 1915)*                |                        |                     |                               |          |           |            |
| <i>Falco femoralis</i> Temminck, 1822            | Aplomado Falcon        | R                   | LC; LC                        |          | WA 47056  |            |
| <i>F. f. femoralis</i> Temminck, 1822*           |                        |                     |                               |          |           |            |
| <i>Falco peregrinus</i> Tunstall, 1771           | Peregrine Falcon       | VN                  | LC; LC                        |          | WA 287761 |            |
| <i>F. p. tundrius</i> C. M. White, 1968*         |                        |                     |                               |          |           |            |
| <i>F. p. anatum</i> Bonaparte, 1838*             |                        |                     |                               |          |           |            |

## PSITTACIFORMES

### Psittacidae

|  |                          |      |        |            |            |            |
|--|--------------------------|------|--------|------------|------------|------------|
| <i>Primolius maracana</i> (Vieillot, 1816)             | Blue-winged Macaw        | R    | NT; NT |            | WA 1333526 | WA 2614632 |
| <i>Diopsittaca nobilis</i> (Linnaeus, 1758)            | Red-shouldered Macaw     | R    | LC; LC |            |            | WA 3233556 |
| <i>D. n. cumanensis</i> (M. H. C. Lichtenstein, 1823)* |                          |      |        |            |            |            |
| <i>Aratinga jandaya</i> (J. F. Gmelin, 1788)           | Jandaya Parakeet         | R, E | LC; LC |            | WA 81727   | WA 1779831 |
| <i>Eupsittula aurea</i> (J. F. Gmelin, 1788)           | Peach-fronted Parakeet   | R    | LC; LC |            | WA 480165  | WA 2591407 |
| <i>Eupsittula cactorum</i> (Kuhl, 1820)                | Cactus Parakeet          | R, E | LC; LC |            | WA 47014   | WA 1254747 |
| <i>E. c. caixana</i> (Spix, 1824)*                     |                          |      |        |            |            |            |
| <i>Forpus xanthopterygius</i> (Spix, 1824)             | Blue-winged Parrotlet    | R    | LC; LC | COUFRN 875 | WA 2716847 | WA 1254361 |
| <i>F. x. flavissimus</i> Hellmayr, 1929*               |                          |      |        |            |            |            |
| <i>Amazona amazonica</i> (Linnaeus, 1766)              | Orange-winged Parrot     | R    | LC; LC |            | WA 619977  | WA 2063543 |
| <i>Amazona aestiva</i> (Linnaeus, 1758)                | Turquoise-fronted Parrot | R    | NT; LC |            | WA 2614629 | WA 2650544 |
| <i>A. a. aestiva</i> (Linnaeus, 1758)                  |                          |      |        |            |            |            |

## PASSERIFORMES

### Thamnophilidae

|  |                       |   |        |           |           |            |
|--|-----------------------|---|--------|-----------|-----------|------------|
| <i>Myrmorchilus strigilatus</i> (Wied, 1831) | Stripe-backed Antbird | R | LC; LC | COUFRN 52 | WA 347677 | WA 3300914 |
| <i>M. s. strigilatus</i> (Wied, 1831)        |                       |   |        |           |           |            |

| Taxon  | English name              | National occurrence | National; Global conservation | Specimen    | Photo      | Sound                      |
|--|---------------------------|---------------------|-------------------------------|-------------|------------|----------------------------|
| <i>Myrmotherula axillaris</i> (Vieillot, 1817)       | White-flanked Antwren     | R                   | LC; LC                        | COUFRN 27   | WA 1359465 |                            |
| <i>M. a. luctuosa</i> Pelzeln, 1868                  |                           |                     | LC                            |             |            |                            |
| <i>Formicivora grisea</i> (Boddaert, 1783)           | White-fringed Antwren     | R                   | LC; LC                        | LSUMZ 10422 | WA 90135   | Minns <i>et al.</i> (2010) |
| <i>F. g. grisea</i> (Boddaert, 1783)*                |                           |                     |                               |             |            |                            |
| <i>Formicivora melanogaster</i> Pelzeln, 1868        | Black-bellied Antwren     | R                   | LC; LC                        | COUFRN 80   | WA 346640  | WA 1254742                 |
| <i>F. m. bahiae</i> Hellmayr, 1909*                  |                           |                     |                               |             |            |                            |
| <i>Formicivora rufa</i> (Wied, 1831)                 | Rusty-backed Antwren      | R                   | LC; LC                        |             | WA 247663; | XC 430046                  |
| <i>F. r. rufa</i> (Wied, 1831)*                      |                           |                     |                               |             |            |                            |
| <i>Dysithamnus mentalis</i> (Temminck, 1823)         | Plain Antvireo            | R                   | LC; LC                        | MNRJ 8092   | WA 452161  | WA 2577546                 |
| <i>D. m. emiliae</i> Hellmayr, 1912*                 |                           |                     |                               |             |            |                            |
| <i>Herpsilochmus sellowi</i> Whitney & Pacheco, 2000 | Caatinga Antwren          | R, E                | LC; LC                        | COUFRN 94   | WA 250894  | Minns <i>et al.</i> (2010) |
| <i>Herpsilochmus atricapillus</i> Pelzeln, 1868      | Black-capped Antwren      | R                   | LC; LC                        |             | WA 1650539 | Minns <i>et al.</i> (2010) |
| <i>Herpsilochmus pectoralis</i> P. L. Sclater, 1857  | Pectoral Antwren          | R, E                | LC; VU                        | COUFRN 1    | WA 452125  | Minns <i>et al.</i> (2010) |
| <i>Herpsilochmus rufimarginatus</i> (Temminck, 1822) | Rufous-winged Antwren     | R                   | LC; LC                        | COUFRN 101  | WA 1876728 | WA 253723                  |
| <i>H. r. scapularis</i> (Wied, 1831)*                |                           |                     |                               |             |            |                            |
| <i>Sakesphorus cristatus</i> (Wied,1831)             | Silvery-cheeked Antshrike | R, E                | LC; LC                        | COUFRN 252  | WA 343961  | WA 3300881                 |
| <i>Thamnophilus capistratus</i> Lesson, 1840         | Caatinga Antshrike        | R, E                | LC; NE                        | COUFRN 108  | WA 348848  | WA 1254752                 |
| <i>Thamnophilus torquatus</i> Swainson, 1825         | Rufous-winged Antshrike   | R                   | LC; LC                        | COUFRN 96   | WA 2016715 | XC 430049                  |
| <i>Thamnophilus pelzelni</i> Hellmayr, 1924          | Planalto Slaty Antshrike  | R, E                | LC; LC                        | COUFRN 34   | WA 72129   | WA 2063563                 |
| <i>Taraba major</i> (Vieillot, 1816)                 | Great Antshrike           | R                   | LC; LC                        | COUFRN 31   | WA 193676  | XC 430050                  |
| <i>T. m. stagurus</i> (M. H. C. Lichtenstein, 1823)* |                           |                     |                               |             |            |                            |
| <i>Cercomacroides laeta</i> (Todd, 1920)             | Willis's Antbird          | R, E                | LC; LC                        |             |            | WA 2660533                 |
| <i>C. l. sabinoi</i> (Pinto, 1939)*                  |                           |                     | NE                            |             |            |                            |
| <b>Conopophagidae</b>                                |                           |                     |                               |             |            |                            |
| <i>Conopophaga cearae</i> Cory, 1916                 | Ceara Gnat eater          | R, E                | EN; LC                        |             | WA 2159276 | WA 3128684                 |
| <i>Conopophaga melanops</i> (Vieillot, 1818)         | Black-cheeked Gnat eater  | R, E                | LC; LC                        |             | WA 2304026 | WA 2236800                 |
| <i>C. m. nigrifrons</i> Pinto, 1954*                 |                           |                     | VU                            |             |            |                            |
| <b>Dendrocolaptidae</b>                              |                           |                     |                               |             |            |                            |
| <i>Sittasomus griseicapillus</i> (Vieillot, 1818)    | Olivaceous Woodcreeper    | R                   | LC; LC                        | COUFRN 1172 | WA 2603253 | XC 388588                  |
| <i>S. g. reiseri</i> Hellmayr, 1917*                 |                           |                     |                               |             |            |                            |



| Taxon  | English name                | National occurrence | National; Global conservation | Specimen    | Photo      | Sound                      |
|--|-----------------------------|---------------------|-------------------------------|-------------|------------|----------------------------|
| <i>Xiphorhynchus guttatus</i> (M. H. C. Lichtenstein, 1820)          | Buff-throated Woodcreeper   | R                   | LC; LC                        |             | WA 619978  | WA 2064613                 |
| <i>X. g. guttatus</i> (M. H. C. Lichtenstein, 1820)*                 |                             |                     |                               |             |            |                            |
| <i>Campylorhamphus trochilirostris</i> (M. H. C. Lichtenstein, 1820) | Red-billed Scythebill       | R                   | LC; LC                        | COUFRN 1107 | WA 1549462 | WA 2966049                 |
| <i>C. t. major</i> Ridgway, 1911*                                    |                             |                     |                               |             |            |                            |
| <i>Dendroplex picus</i> (J. F. Gmelin, 1788)                         | Straight-billed Woodcreeper | R                   | LC; LC                        | COUFRN 61   | WA 250856  | WA 3300882                 |
| <i>D. p. picus</i> (J. F. Gmelin, 1788)*                             |                             |                     |                               |             |            |                            |
| <i>Lepidocolaptes angustirostris</i> (Vieillot, 1818)                | Narrow-billed Woodcreeper   | R                   | LC; LC                        | COUFRN 74   | WA 47542   | WA 3300884                 |
| <b>Xenopidae</b>   |                             |                     |                               |             |            |                            |
| <i>Xenops minutus</i> (Sparman, 1788)                                | Plain Xenops                | R                   | LC; LC                        | COUFRN 16   | WA 2114850 | WA 2063601                 |
| <i>X. m. alagoanus</i> Pinto, 1954*                                  |                             |                     | VU                            |             |            |                            |
| [ <i>Xenops rutilans</i> Temminck, 1821]                             | Streaked Xenops             | R                   | LC; NE                        |             |            |                            |
| <b>Furnariidae</b>   |                             |                     |                               |             |            |                            |
| <i>Furnarius figulus</i> (M. H. C. Lichtenstein, 1823)               | Wing-banded Hornero         | R, E                | LC; LC                        |             | WA 324819  | WA 2617452                 |
| <i>F. f. figulus</i> (M. H. C. Lichtenstein, 1823)*                  |                             |                     |                               |             |            |                            |
| <i>Furnarius leucopus</i> Swainson, 1838                             | Pale-legged Hornero         | R                   | LC; LC                        | MNRJ 34527  | WA 249194  | Minns <i>et al.</i> (2010) |
| <i>F. l. assimilis</i> Cabanis & Heine, 1859                         |                             |                     |                               |             |            |                            |
| <i>Pseudoseisura cristata</i> (Spix, 1824)                           | Caatinga Cacholote          | R, E                | LC; LC                        |             | WA 249192  | WA 1254342                 |
| <i>Phacellodomus rufifrons</i> (Wied, 1821)                          | Rufous-fronted Thornbird    | R                   | LC; LC                        |             | WA 1851438 |                            |
| <i>P. r. rufifrons</i> (Wied, 1821)*                                 |                             |                     |                               |             |            |                            |
| <i>Certhiaxis cinnamomeus</i> (J. F. Gmelin, 1788)                   | Yellow-chinned Spinetail    | R                   | LC; LC                        | CAHZ 1245   | WA 249146  | WA 1649739                 |
| <i>C. c. cearensis</i> (Cory, 1916)*                                 |                             |                     |                               |             |            |                            |
| <i>Synallaxis hellmayri</i> Reiser, 1905                             | Red-shouldered Spinetail    | R, E                | DD; NT                        | COUFRN 244  | WA 236135  | WA 3300886                 |
| <i>Synallaxis frontalis</i> Pelzel, 1859                             | Sooty-fronted Spinetail     | R                   | LC; LC                        | MNRJ 36903  | WA 571728  | WA 2266747                 |
| <i>Synallaxis albescens</i> Temminck, 1823                           | Pale-breasted Spinetail     | R                   | LC; LC                        | COUFRN 922  | WA 2089572 | WA 2088131                 |
| <i>S. a. albescens</i> Temminck, 1823*                               |                             |                     |                               |             |            |                            |
| <i>Synallaxis scutata</i> P. L. Sclater, 1859                        | Ochre-cheeked Spinetail     | R                   | LC; LC                        | COUFRN 127  | WA 452167  | WA 3065184                 |
| <i>S. s. scutata</i> P. L. Sclater, 1859*                            |                             |                     |                               |             |            |                            |
| <b>Pipridae</b>  |                             |                     |                               |             |            |                            |
| <i>Neopelma pallescens</i> (Lafresnaye, 1853)                        | Pale-bellied Tyrant-Manakin | R                   | LC; LC                        | COUFRN 59   | WA 2016745 | Minns <i>et al.</i> (2010) |

| Taxon   | English name               | National occurrence | National; Global conservation | Specimen    | Photo      | Sound                      |
|---|----------------------------|---------------------|-------------------------------|-------------|------------|----------------------------|
| [ <i>Ceratopipra rubrocapilla</i> (Temminck, 1821)]       | Red-headed Manakin         | R                   | LC; LC                        |             |            |                            |
| <i>Manacus manacus</i> (Linnaeus, 1766)                   | White-bearded Manakin      | R                   | LC; LC                        |             | WA 2016730 | WA 2457367                 |
| <i>M. m. gutturosus</i> (Desmarest, 1806)*                |                            |                     |                               |             |            |                            |
| <i>Chiroxiphia pareola</i> (Linnaeus, 1766)               | Blue-backed Manakin        | R                   | LC; LC                        | COUFRN 345  | WA 37071   | Minns <i>et al.</i> (2010) |
| <i>C. p. pareola</i> (Linnaeus, 1766)*                    |                            |                     |                               |             |            |                            |
| <b>Onychorhynchidae</b>                                   |                            |                     |                               |             |            |                            |
| [ <i>Myiobius barbatus</i> (J. F. Gmelin, 1789)]          | Whiskered Flycatcher       | R                   | LC; LC                        |             |            |                            |
| <b>Tityridae</b>  |                            |                     |                               |             |            |                            |
| <i>Pachyramphus viridis</i> (Vieillot, 1816)              | Green-backed Becard        | R                   | LC; LC                        | COUFRN 1054 | WA 1364274 |                            |
| <i>P. v. viridis</i> (Vieillot, 1816)*                    |                            |                     |                               |             |            |                            |
| <i>Pachyramphus polychopterus</i> (Vieillot, 1818)        | White-winged Becard        | R                   | LC; LC                        | UFPE 3437   | WA 283179  | Minns <i>et al.</i> (2010) |
| <i>P. p. polychopterus</i> (Vieillot, 1818)*              |                            |                     |                               |             |            |                            |
| <i>Pachyramphus validus</i> (M. H. C. Lichtenstein, 1823) | Crested Becard             | R                   | LC; LC                        |             | WA 47060   | WA 1672890                 |
| <i>P. v. validus</i> (M. H. C. Lichtenstein, 1823)*       |                            |                     |                               |             |            |                            |
| <i>Xenopsaris albinucha</i> (Burmeister, 1869)            | White-naped Xenopsaris     | R                   | LC; LC                        | MNRJ 36939  | WA 186702  | WA 3070452                 |
| <i>X. a. albinucha</i> (Burmeister, 1869)*                |                            |                     |                               |             |            |                            |
| <b>Cotingidae</b>   |                            |                     |                               |             |            |                            |
| <i>Procnias averano</i> (Hermann, 1783)                   | Bearded Bellbird           | R                   | LC; LC                        |             | WA 1724942 |                            |
| <i>P. a. averano</i> (Hermann, 1783)*                     |                            |                     |                               |             |            |                            |
| <b>Platyrinchidae</b>                                     |                            |                     |                               |             |            |                            |
| <i>Platyrinchus mystaceus</i> Vieillot, 1818              | White-throated Spadebill   | R                   | LC; LC                        |             | WA 295438  | WA 2598721                 |
| <i>P. m. niveigularis</i> Pinto, 1954*                    |                            |                     |                               |             |            |                            |
| VU  |                            |                     |                               |             |            |                            |
| <b>Rhynchocyclidae</b>                                    |                            |                     |                               |             |            |                            |
| <i>Leptopogon amaurocephalus</i> Tschudi, 1846            | Sepia-capped Flycatcher    | R                   | LC; LC                        |             | WA 2513911 |                            |
| <i>L. a. amaurocephalus</i> Tschudi, 1846*                |                            |                     |                               |             |            |                            |
| <i>Tolmomyias flaviventris</i> (Wied, 1831)               | Yellow-breasted Flycatcher | R                   | LC; LC                        | UFPE 3435   | WA 280125  | WA 3300888                 |
| <i>T. f. flaviventris</i> (Wied, 1831)*                   |                            |                     |                               |             |            |                            |
| <i>Todirostrum cinereum</i> (Linnaeus, 1766)              | Common Tody-Flycatcher     | R                   | LC; LC                        | COUFRN 322  | WA 193675  | WA 1254334                 |
| <i>T. c. cearae</i> Cory, 1916*                           |                            |                     |                               |             |            |                            |



| Taxon  | English name                  | National occurrence | National; Global conservation | Specimen    | Photo      | Sound                      |
|--|-------------------------------|---------------------|-------------------------------|-------------|------------|----------------------------|
| <i>Poecilotriccus fumifrons</i> (Hartlaub, 1853)                     | Smoky-fronted Tody-Flycatcher | R                   | LC; LC                        |             | WA 2270071 | Minns <i>et al.</i> (2010) |
| <i>P. f. fumifrons</i> (Hartlaub, 1853)*                             |                               |                     |                               |             |            |                            |
| <i>Hemitriccus griseipectus</i> (E. Snethlage, 1907)                 | White-bellied Tody-Tyrant     | R                   | LC; LC                        |             | WA 1359463 | WA 2011736                 |
| <i>H. g. naumburgae</i> (J. T. Zimmer, 1945)*                        |                               |                     | VU                            |             |            |                            |
| <i>Hemitriccus striaticollis</i> (Lafresnaye, 1853)                  | Stripe-necked Tody-Tyrant     | R                   | LC; LC                        | COUFRN 44   | WA 587100  | Minns <i>et al.</i> (2010) |
| <i>H. s. striaticollis</i> (Lafresnaye, 1853)*                       |                               |                     |                               |             |            |                            |
| <i>Hemitriccus margaritaceiventer</i> (d'Orbigny & Lafresnaye, 1837) | Pearly-vented Tody-Tyrant     | R                   | LC; LC                        | UFPE 3440   | WA 344899  | WA 1254332                 |
| <i>H. m. wuchereri</i> (Sclater & Salvin, 1873)                      |                               |                     |                               |             |            |                            |
| <b>Tyrannidae</b>  |                               |                     |                               |             |            |                            |
| <i>Hirundinea ferruginea</i> (J. F. Gmelin, 1788)                    | Cliff Flycatcher              | R                   | LC; LC                        |             | WA 269104  | WA 3300889                 |
| <i>H. f. bellicosa</i> (Vieillot, 1819)*                             |                               |                     |                               |             |            |                            |
| <i>Stigmatura napensis</i> Chapman, 1926                             | Lesser Wagtail-Tyrant         | R                   | LC; LC                        | COUFRN 149  | WA 1689483 | WA 1254725                 |
| <i>S. n. bahiae</i> Chapman, 1926*                                   |                               |                     | LC                            |             |            |                            |
| <i>Euscarthmus meloryphus</i> Wied, 1831                             | Tawny-crowned Pygmy Tyrant    | R                   | LC; LC                        | COUFRN 148  | WA 47062   | Minns <i>et al.</i> (2010) |
| <i>E. m. meloryphus</i> Wied, 1831*                                  |                               |                     |                               |             |            |                            |
| <i>Ornithion inermis</i> Hartlaub, 1853                              | White-lored Tyrannulet        | R                   | LC; LC                        |             | WA 3257427 | WA 3300921                 |
| <i>Camptostoma obsoletum</i> (Temminck, 1824)                        | Southern Beardless Tyrannulet | R                   | LC; LC                        | COUFRN 708  | WA 198231  | WA 3300890                 |
| <i>C. o. cinerascens</i> (Wied, 1831)*                               |                               |                     |                               |             |            |                            |
| <i>Elaenia flavogaster</i> (Thunberg, 1822)                          | Yellow-bellied Elaenia        | R                   | LC; LC                        | UFPE 3436   | WA 452134  | Minns <i>et al.</i> (2010) |
| <i>E. f. flavogaster</i> (Thunberg, 1822)*                           |                               |                     |                               |             |            |                            |
| <i>Elaenia spectabilis</i> Pelzelin, 1868                            | Large Elaenia                 | R                   | LC; LC                        | COUFRN 432  | WA 47193   | WA 1254733                 |
| <i>Elaenia chilensis</i> Hellmayr, 1927                              | Chilean Elaenia               | VS                  | LC; NE                        | COUFRN 64   | WA 2111209 | WA 2756392                 |
| <i>Elaenia cristata</i> Pelzelin, 1868                               | Plain-crested Elaenia         | R                   | LC; LC                        | LSUMZ 10420 | WA 253384  | WA 1570871                 |
| <i>E. c. cristata</i> Pelzelin, 1868*                                |                               |                     |                               |             |            |                            |
| <i>Elaenia chiriquensis</i> Lawrence, 1865                           | Lesser Elaenia                | R                   | LC; LC                        | COUFRN 115  | WA 2030096 | XC 430033                  |
| <i>E. c. albivertex</i> Pelzelin, 1868*                              |                               |                     |                               |             |            |                            |
| <i>Suiriri suiriri</i> (Vieillot, 1818)                              | Suiriri Flycatcher            | R                   | LC; LC                        |             | WA 2273569 | WA 2931035                 |
| <i>S. s. bahiae</i> (Berlepsch, 1893)*                               |                               |                     |                               |             |            |                            |
| [ <i>Myiopagis caniceps</i> (Swainson, 1835)]                        | Grey Elaenia                  | R                   | LC; LC                        |             |            |                            |

| Taxon  | English name              | National occurrence | National; Global conservation | Specimen    | Photo      | Sound                      |
|--|---------------------------|---------------------|-------------------------------|-------------|------------|----------------------------|
| <i>Myiopagis viridicata</i> (Vieillot, 1817)             | Greenish Elaenia          | R                   | LC; LC                        | COUFRN 60   | WA 730141  | WA 2905285                 |
| <i>M. v. viridicata</i> (Vieillot, 1817)*                |                           |                     |                               |             |            |                            |
| <i>Capsiempis flaveola</i> (M. H. C. Lichtenstein, 1823) | Yellow Tyrannulet         | R                   | LC; LC                        |             | WA 2657045 | WA 2655815                 |
| <i>C. f. flaveola</i> (M. H. C. Lichtenstein, 1823)*     |                           |                     |                               |             |            |                            |
| <i>Phaeomyias murina</i> (Spix, 1825)                    | Mouse-coloured Tyrannulet | R                   | LC; LC                        | COUFRN 146  | WA 1366321 | XC 430051                  |
| <i>P. m. murina</i> (Spix, 1825)*                        |                           |                     |                               |             |            |                            |
| <i>Serpophaga subcristata</i> (Vieillot, 1817)           | White-crested Tyrannulet  | R                   | LC; LC                        |             | WA 2620697 |                            |
| <i>S. s. straminea</i> (Temminck, 1822)*                 |                           |                     |                               |             |            |                            |
| <i>Myiarchus swainsoni</i> Cabanis & Heine, 1859         | Swainson's Flycatcher     | R                   | LC; LC                        | COUFRN 1088 | WA 2932951 | WA 3300894                 |
| <i>M. s. swainsoni</i> Cabanis & Heine, 1859*            |                           |                     |                               |             |            |                            |
| <i>Myiarchus ferox</i> (J. F. Gmelin, 1789)              | Short-crested Flycatcher  | R                   | LC; LC                        |             |            | WA 2984460                 |
| <i>M. f. ferox</i> (J. F. Gmelin, 1789)*                 |                           |                     |                               |             |            |                            |
| <i>Myiarchus tyrannulus</i> (Statius Muller, 1776)       | Brown-crested Flycatcher  | R                   | LC; LC                        | CAHZ 698    | WA 34098   | WA 3300892                 |
| <i>M. t. bahiae</i> Berlepsch & Leverkühn, 1890*         |                           |                     |                               |             |            |                            |
| <i>Casiornis fuscus</i> Sclater & Salvin, 1873           | Ash-throated Casiornis    | R, E                | LC; LC                        | COUFRN 293  | WA 34103   | Minns <i>et al.</i> (2010) |
| <i>Pitangus sulphuratus</i> (Linnaeus, 1766)             | Great Kiskadee            | R                   | LC; LC                        | COUFRN 86   | WA 55170   | Minns <i>et al.</i> (2010) |
| <i>P. s. maximiliani</i> (Cabanis & Heine, 1859)*        |                           |                     |                               |             |            |                            |
| <i>Machetornis rixosa</i> (Vieillot, 1819)               | Cattle Tyrant             | R                   | LC; LC                        |             | WA 251715  |                            |
| <i>M. r. rixosa</i> (Vieillot, 1819)*                    |                           |                     |                               |             |            |                            |
| <i>Myiodynastes maculatus</i> (Statius Muller, 1776)     | Streaked Flycatcher       | R                   | LC; LC                        | COUFRN 947  | WA 198225  | WA 3300904                 |
| <i>M. m. solitarius</i> (Vieillot, 1819)*                |                           |                     |                               |             |            |                            |
| <i>Megarynchus pitangua</i> (Linnaeus, 1766)             | Boat-billed Flycatcher    | R                   | LC; LC                        |             | WA 250907  | XC 366782                  |
| <i>M. p. pitangua</i> (Linnaeus, 1766)*                  |                           |                     |                               |             |            |                            |
| <i>Myiozetetes similis</i> (Spix, 1825)                  | Social Flycatcher         | R                   | LC; LC                        |             | WA 269146  | XC 430058                  |
| <i>M. s. pallidiventr</i> Pinto, 1935*                   |                           |                     |                               |             |            |                            |
| <i>Tyrannus melancholicus</i> Vieillot, 1819             | Tropical Kingbird         | R                   | LC; LC                        | UMMZ 113608 | WA 98160   | WA 1254723                 |
| <i>T. m. despotes</i> (M. H. C. Lichtenstein, 1823)*     |                           |                     |                               |             |            |                            |
| <i>Tyrannus savana</i> Daudin, 1802                      | Fork-tailed Flycatcher    | R                   | LC; LC                        |             | WA 241803  |                            |
| <i>T. s. savana</i> Daudin, 1802*                        |                           |                     |                               |             |            |                            |

| Taxon  | English name               | National occurrence | National; Global conservation | Specimen    | Photo      | Sound                      |
|--|----------------------------|---------------------|-------------------------------|-------------|------------|----------------------------|
| <i>Empidonomus varius</i> (Vieillot, 1818)               | Variegated Flycatcher      | R                   | LC; LC                        |             | WA 342787  |                            |
| <i>E. v. rufinus</i> (Spix, 1825)*                       |                            |                     |                               |             |            |                            |
| <i>Myiophobus fasciatus</i> (Statius Muller, 1776)       | Bran-coloured Flycatcher   | R                   | LC; LC                        |             | WA 198229  | Minns <i>et al.</i> (2010) |
| <i>M. f. flammiceps</i> (Temminck, 1822)*                |                            |                     |                               |             |            |                            |
| <i>Sublegatus modestus</i> (Wied, 1831)                  | Southern Scrub Flycatcher  | R                   | LC; LC                        | COUFRN 107  | WA 690623  | WA 326535                  |
| <i>S. m. modestus</i> (Wied, 1831)*                      |                            |                     |                               |             |            |                            |
| <i>Fluvicola albiventer</i> (Spix, 1825)                 | Black-backed Water Tyrant  | R                   | LC; LC                        |             | WA 346984  |                            |
| <i>Fluvicola nengeta</i> (Linnaeus, 1766)                | Masked Water Tyrant        | R                   | LC; LC                        | COUFRN 709  | WA 1079213 | WA 2762132                 |
| <i>F. n. nengeta</i> (Linnaeus, 1766)*                   |                            |                     |                               |             |            |                            |
| <i>Arundinicola leucocephala</i> (Linnaeus, 1764)        | White-headed Marsh Tyrant  | R                   | LC; LC                        |             | WA 269130  |                            |
| <i>Cnemotriccus fuscatus</i> (Wied, 1831)                | Fuscous Flycatcher         | R                   | LC; LC                        | COUFRN 50   | WA 638569  |                            |
| <i>C. f. bimaculatus</i> (d'Orbigny & Lafresnaye, 1837)* |                            |                     |                               |             |            |                            |
| [ <i>Lathrotriccus euleri</i> (Cabanis, 1868)]           | Euler's Flycatcher         | R                   | LC; LC                        |             |            |                            |
| <i>Knipolegus nigerrimus</i> (Vieillot, 1818)            | Velvety Black Tyrant       | R, E                | LC; LC                        | MNRJ 2078   | WA 2725455 |                            |
| <i>K. n. nigerrimus</i> (Vieillot, 1818)                 |                            |                     |                               |             |            |                            |
| <i>Hymenops perspicillatus</i> (J. F. Gmelin, 1789)      | Spectacled Tyrant          | R                   | LC; LC                        |             | WA 3065355 |                            |
| <i>H. p. perspicillatus</i> (J. F. Gmelin, 1789)*        |                            |                     |                               |             |            |                            |
| <i>Satrapa icterophrys</i> (Vieillot, 1818)              | Yellow-browed Tyrant       | R                   | LC; LC                        |             | WA 649286  |                            |
| <i>Xolmis cinereus</i> (Vieillot, 1816)                  | Grey Monjita               | R                   | LC; LC                        |             | WA 1689513 |                            |
| <i>X. c. cinereus</i> (Vieillot, 1816)*                  |                            |                     |                               |             |            |                            |
| <i>Xolmis irupero</i> (Vieillot, 1823)                   | White Monjita              | R                   | LC; LC                        |             | WA 2712042 |                            |
| <i>X. i. niveus</i> (Spix, 1825)*                        |                            |                     |                               |             |            |                            |
| <b>Vireonidae</b>  |                            |                     |                               |             |            |                            |
| <i>Cyclarhis gujanensis</i> (J. F. Gmelin, 1789)         | Rufous-browed Peppershrike | R                   | LC; LC                        | UFPE 3434   | WA 2697924 | Minns <i>et al.</i> (2010) |
| <i>C. g. cearensis</i> Baird, 1866*                      |                            |                     |                               |             |            |                            |
| <i>Hylophilus amaurocephalus</i> (Nordmann, 1835)        | Grey-eyed Greenlet         | R, E                | LC; LC                        | COUFRN 45   | WA 295439  | Minns <i>et al.</i> (2010) |
| <i>Vireo chivi</i> (Vieillot, 1817)                      | Chivi Vireo                | R                   | LC; NE                        | COUFRN 207  | WA 866015  | Minns <i>et al.</i> (2010) |
| <i>V. c. agilis</i> (M. H. C. Lichtenstein, 1823)*       |                            |                     |                               |             |            |                            |
| <b>Corvidae</b>  |                            |                     |                               |             |            |                            |
| <i>Cyanocorax cyanopogon</i> (Wied, 1821)                | White-naped Jay            | R, E                | LC; LC                        | COUFRN 1159 | WA 47469   | XC 429375                  |

| Taxon   | English name                  | National occurrence | National; Global conservation | Specimen    | Photo      | Sound                      |
|---|-------------------------------|---------------------|-------------------------------|-------------|------------|----------------------------|
| <b>Hirundinidae</b>                                     |                               |                     |                               |             |            |                            |
| <i>Pygochelidon cyanoleuca</i> (Vieillot, 1817)         | Blue-and-white Swallow        |                     | LC; LC                        |             | WA 1777794 |                            |
| <i>P. c. patagonica</i> (d'Orbigny & Lafresnaye, 1837)* |                               |                     |                               |             |            |                            |
| <i>Stelgidopteryx ruficollis</i> (Vieillot, 1817)       | Southern Rough-winged Swallow | R                   | LC; LC                        |             | WA 196389  |                            |
| <i>S. r. ruficollis</i> (Vieillot, 1817)*               |                               |                     |                               |             |            |                            |
| <i>Progne tapera</i> (Vieillot, 1817)                   | Brown-chested Martin          | R                   | LC; LC                        | LSUMZ 10421 | WA 2634379 |                            |
| <i>P. t. tapera</i> (Linnaeus, 1766)                    |                               |                     |                               |             |            |                            |
| <i>P. t. fusca</i> (Vieillot, 1817)                     |                               |                     |                               |             |            |                            |
| <i>Progne chalybea</i> (Gmelin, 1789)                   | Grey-breasted Martin          | R                   | LC; LC                        | COUFRN 552  | WA 2723512 | WA 2982109                 |
| <i>P. c. macrorhamphus</i> Brooke, 1974*                |                               |                     |                               |             |            |                            |
| <i>Tachycineta albiventer</i> (Boddaert, 1783)          | White-winged Swallow          | R                   | LC; LC                        | COUFRN 991  | WA 725017  |                            |
| <i>Tachycineta leucorrhoa</i> (Vieillot, 1817)          | White-rumped Swallow          | R                   | LC; LC                        |             | Fig. 15A   |                            |
| <i>Tachycineta leucopyga</i> (Meyen, 1834)              | Chilean Swallow               | VS                  | LC; LC                        |             | Fig. 16    |                            |
| <i>Hirundo rustica</i> Linnaeus, 1758                   | Barn Swallow                  | VN                  | LC; LC                        |             | WA 99051   |                            |
| <i>H. r. erythrogaster</i> Boddaert, 1783               |                               |                     |                               |             |            |                            |
| <b>Troglodytidae</b>                                    |                               |                     |                               |             |            |                            |
| <i>Troglodytes musculus</i> Naumann, 1823               | Southern House Wren           | R                   | LC; NE                        | COUFRN 22   | WA 343960  | WA 1254734                 |
| <i>T. m. musculus</i> Naumann, 1823*                    |                               |                     |                               |             |            |                            |
| <i>Pheugopedius genibarbis</i> (Swainson, 1838)         | Moustached Wren               | R                   | LC; LC                        |             | WA 2015201 | WA 3300896                 |
| <i>P. g. genibarbis</i> (Swainson, 1838)*               |                               |                     |                               |             |            |                            |
| <i>Cantorchilus longirostris</i> (Vieillot, 1819)       | Long-billed Wren              | R, E                | LC; LC                        | COUFRN 56   | WA 247605  | XC 429380                  |
| <i>C. l. bahiae</i> (Hellmayr, 1903)*                   |                               |                     |                               |             |            |                            |
| <b>Donacobiidae</b>                                     |                               |                     |                               |             |            |                            |
| <i>Donacobius atricapilla</i> (Linnaeus, 1766)          | Black-capped Donacobius       | R                   | LC; LC                        | COUFRN 14   | WA 279016  |                            |
| <i>D. a. atricapilla</i> (Linnaeus, 1766)*              |                               |                     |                               |             |            |                            |
| <b>Poliophtilidae</b>                                   |                               |                     |                               |             |            |                            |
| <i>Ramphocaenus melanurus</i> Vieillot, 1819            | Long-billed Gnatwren          | R                   | LC; LC                        |             | WA 2214719 | WA 2270254                 |
| <i>R. m. melanurus</i> Vieillot, 1819*                  |                               |                     |                               |             |            |                            |
| <i>Poliophtila plumbea</i> (J. F. Gmelin, 1788)         | Tropical Gnatcatcher          | R                   | LC; LC                        | COUFRN 72   | WA 248474  | Minns <i>et al.</i> (2010) |
| <i>P. p. atricapilla</i> (Swainson, 1831)*              |                               |                     |                               |             |            |                            |
| <b>Turdidae</b>   |                               |                     |                               |             |            |                            |
| <i>Turdus flavipes</i> Vieillot, 1818                   | Yellow-legged Thrush          | R                   | LC; LC                        | COUFRN 812  | WA 1662001 |                            |
| <i>T. f. flavipes</i> Vieillot, 1818                    |                               |                     |                               |             |            |                            |

| Taxon   | English name             | National occurrence | National; Global conservation | Specimen   | Photo      | Sound                      |
|---|--------------------------|---------------------|-------------------------------|------------|------------|----------------------------|
| <i>Turdus leucomelas</i> Vieillot, 1818               | Pale-breasted Thrush     | R                   | LC; LC                        | COUFRN 218 | WA 657407  | Minns <i>et al.</i> (2010) |
| <i>T. l. albiventer</i> Spix, 1824*                   |                          |                     |                               |            |            |                            |
| <i>Turdus rufiventris</i> Vieillot, 1818              | Rufous-bellied Thrush    | R                   | LC; LC                        | COUFRN 112 | WA 2713910 | WA 349956                  |
| <i>T. r. juensis</i> (Cory, 1916)*                    |                          |                     |                               |            |            |                            |
| <i>Turdus amaurochalinus</i> Cabanis, 1850            | Creamy-bellied Thrush    | R                   | LC; LC                        | COUFRN 144 | WA 198226  | Minns <i>et al.</i> (2010) |
| <b>Mimidae</b>  |                          |                     |                               |            |            |                            |
| <i>Mimus gilvus</i> (Vieillot, 1807)                  | Tropical Mockingbird     | R                   | LC; LC                        |            | WA 1343747 | WA 1254333                 |
| <i>M. g. antelius</i> Oberholser, 1919*               |                          |                     |                               |            |            |                            |
| <i>Mimus saturninus</i> (M. H. C. Lichtenstein, 1823) | Chalk-browed Mockingbird | R                   | LC; LC                        | COUFRN 863 | WA 193679  | WA 1254366                 |
| <i>M. s. arenaceus</i> Chapman, 1890*                 |                          |                     |                               |            |            |                            |
| <b>Motacillidae</b>                                   |                          |                     |                               |            |            |                            |
| <i>Anthus lutescens</i> Pucheran, 1855                | Yellowish Pipit          | R                   | LC; LC                        |            | WA 80471   |                            |
| <i>A. l. lutescens</i> Pucheran, 1855*                |                          |                     |                               |            |            |                            |
| <b>Passerellidae</b>                                  |                          |                     |                               |            |            |                            |
| <i>Zonotrichia capensis</i> (Statius Muller, 1776)    | Rufous-collared Sparrow  | R                   | LC; LC                        | UFPE 3655  | WA 348722  | FNJV 7522                  |
| <i>Z. c. matutina</i> (M. H. C. Lichtenstein, 1823)*  |                          |                     |                               |            |            |                            |
| <i>Ammodramus humeralis</i> (Bosc, 1792)              | Grassland Sparrow        | R                   | LC; LC                        | UFPE 3398  | WA 249001  | Minns <i>et al.</i> (2010) |
| <i>A. h. humeralis</i> (Bosc, 1792)*                  |                          |                     |                               |            |            |                            |
| <i>Arremon taciturnus</i> (Hermann, 1783)             | Pectoral Sparrow         | R                   | LC; LC                        |            | WA 283184  | Minns <i>et al.</i> (2010) |
| <i>A. t. taciturnus</i> (Hermann, 1783)*              |                          |                     |                               |            |            |                            |
| <b>Parulidae</b>                                      |                          |                     |                               |            |            |                            |
| <i>Setophaga pitiayumi</i> (Vieillot, 1817)           | Tropical Parula          | R                   | LC; LC                        |            | WA 2250869 |                            |
| <i>S. p. pitiayumi</i> (Vieillot, 1817)*              |                          |                     |                               |            |            |                            |
| <i>Basileuterus culicivorus</i> (Deppe, 1830)         | Golden-crowned Warbler   | R                   | LC; LC                        | COUFRN 43  | WA 283196  | WA 2615240                 |
| <i>B. c. auricapilla</i> (Swainson, 1838)             |                          |                     |                               |            |            |                            |
| <i>Myiothlypis flaveola</i> Baird, 1865               | Flavescent Warbler       | R                   | LC; LC                        | COUFRN 19  | WA 283217  | Minns <i>et al.</i> (2010) |
| <i>M. f. flaveola</i> Baird, 1865*                    |                          |                     |                               |            |            |                            |
| <b>Icteridae</b>                                      |                          |                     |                               |            |            |                            |
| <i>Procacicus solitarius</i> (Vieillot, 1816)         | Solitary Black Cacique   | R                   | LC; LC                        | COUFRN 219 | WA 251714  | Minns <i>et al.</i> (2010) |
| <i>Cacicus cela</i> (Linnaeus, 1758)                  | Yellow-rumped Cacique    | R                   | LC; LC                        |            | WA 1688477 | WA 2266654                 |
| <i>C. c. cela</i> (Linnaeus, 1758)*                   |                          |                     |                               |            |            |                            |

| Taxon  | English name              | National occurrence | National; Global conservation | Specimen    | Photo      | Sound      |
|--|---------------------------|---------------------|-------------------------------|-------------|------------|------------|
| <i>Icterus pyrrhopterus</i> (Vieillot, 1819)                 | Variable Oriole           | R                   | LC; LC                        |             | WA 251716  | WA 247146  |
| <i>I. p. tibialis</i> Swainson, 1838                         |                           |                     |                               |             |            |            |
| <i>Icterus jamacaii</i> (J. F. Gmelin, 1788)                 | Campo Troupial            | R, E                | LC; LC                        | COUFRN 1009 | WA 47528   | WA 1254350 |
| <i>Gnorimopsar chopi</i> (Vieillot, 1819)                    | Chopi Blackbird           | R                   | LC; LC                        |             | WA 864267  | WA 1254730 |
| <i>G. c. sulcirostris</i> (Spix, 1824)*                      |                           |                     |                               |             |            |            |
| <i>Chrysomus ruficapillus</i> (Vieillot, 1819)               | Chestnut-capped Blackbird | R                   | LC; LC                        |             | WA 726026  | WA 2050548 |
| <i>C. r. frontalis</i> (Vieillot, 1819)*                     |                           |                     |                               |             |            |            |
| <i>Agelaioides fringillarius</i> (Spix, 1824)                | Pale Baywing              | R, E                | LC; LC                        |             | WA 339277  | WA 3300898 |
| <i>Molothrus rufoaxillaris</i> Cassin, 1866                  | Screaming Cowbird         | R                   | LC; LC                        |             | WA 2164752 |            |
| <i>Molothrus bonariensis</i> (J. F. Gmelin, 1789)            | Shiny Cowbird             | R                   | LC; LC                        |             | WA 2716855 | WA 1254336 |
| <i>M. b. bonariensis</i> (J. F. Gmelin, 1789)*               |                           |                     |                               |             |            |            |
| <i>Sturnella supercilialis</i> (Bonaparte, 1850)             | White-browed Meadowlark   | R                   | LC; LC                        |             | WA 47055   | WA 3244419 |
| <b>Thraupidae</b>  |                           |                     |                               |             |            |            |
| <i>Porphyrospiza caerulescens</i> (Wied, 1830)               | Blue Finch                | R                   | LC; NT                        | COUFRN 81   | WA 1689382 |            |
| <i>Schistochlamys melanopis</i> (Latham, 1790)               | Black-faced Tanager       | R                   | LC; LC                        |             | WA 228669  |            |
| <i>S. m. melanopis</i> (Latham, 1790)*                       |                           |                     |                               |             |            |            |
| <i>Schistochlamys ruficapillus</i> (Vieillot, 1817)          | Cinnamon Tanager          | R                   | LC; LC                        | UMMZ 113609 | WA 940543  | WA 283905  |
| <i>Paroaria dominicana</i> (Linnaeus, 1758)                  | Red-cowled Cardinal       | R, E                | LC; LC                        | CAHZ 686    | WA 47186   | WA 1254373 |
| <i>Tangara sayaca</i> (Linnaeus, 1766)                       | Sayaca Tanager            | R                   | LC; LC                        | COUFRN 165  | WA 348726  |            |
| <i>T. s. sayaca</i> (Linnaeus, 1766)*                        |                           |                     |                               |             |            |            |
| <i>Tangara palmarum</i> (Wied, 1821)                         | Palm Tanager              | R                   | LC; LC                        | COUFRN 130  | WA 81822   | WA 2762141 |
| <i>T. p. palmarum</i> (Wied, 1821)*                          |                           |                     |                               |             |            |            |
| <i>Tangara cayana</i> (Linnaeus, 1766)                       | Burnished-buff Tanager    | R                   | LC; LC                        | COUFRN 102  | WA 452006  | XC 430039  |
| <i>T. c. flava</i> (J. F. Gmelin, 1789)                      |                           |                     |                               |             |            |            |
| <i>Nemosia pileata</i> (Boddaert, 1783)                      | Hooded Tanager            | R                   | LC; LC                        |             | WA 198256  |            |
| <i>N. p. caerulea</i> (Wied, 1831)*                          |                           |                     |                               |             |            |            |
| <i>Compsothraupis loricata</i> (M. H. C. Lichtenstein, 1819) | Scarlet-throated Tanager  | R, E                | LC; LC                        |             | WA 338567  | WA 3300900 |
| <i>Conirostrum speciosum</i> (Temminck, 1824)                | Chestnut-vented Conebill  | R                   | LC; LC                        |             | WA 2519005 | WA 2892173 |
| <i>C. s. speciosum</i> (Temminck, 1824)*                     |                           |                     |                               |             |            |            |

| Taxon  | English name             | National occurrence | National; Global conservation | Specimen    | Photo      | Sound                      |
|--|--------------------------|---------------------|-------------------------------|-------------|------------|----------------------------|
| <i>Conirostrum bicolor</i> (Vieillot, 1809)<br><i>C. b. bicolor</i> (Vieillot, 1809)*  | Bicoloured Conebill      | R                   | LC; NT                        |             | WA 385417  | WA 2581741                 |
| <i>Sicalis flaveola</i> (Linnaeus, 1766)<br><i>S. f. brasiliensis</i> (J. F. Gmelin, 1789)   | Saffron Finch            | R                   | LC; LC                        |             | WA 1613389 |                            |
| <i>Sicalis luteola</i> (Sparman, 1789)<br><i>S. l. luteiventris</i> (Meyen, 1834)*   | Grassland Yellow Finch   | R                   | LC; LC                        | UFPE 3140   | WA 249159  | WA 1254714                 |
| <i>Volatinia jacarina</i> (Linnaeus, 1766)<br><i>V. j. jacarina</i> (Linnaeus, 1766)*  | Blue-black Grassquit     | R                   | LC; LC                        | CAHZ 689    | WA 269106  | WA 1254764                 |
| <i>Coryphospingus pileatus</i> (Wied, 1821)<br><i>C. p. pileatus</i> (Wied, 1821)*   | Pileated Finch           | R                   | LC; LC                        | UFPE 3443   | WA 116045  | WA 1254731                 |
| <i>Lanio cristatus</i> (Linnaeus, 1766)<br><i>L. c. brunneus</i> (Spix, 1825)*   | Flame-crested Tanager    | R                   | LC; LC                        |             | WA 228661  |                            |
| <i>Tachyphonus rufus</i> (Boddaert, 1783)<br><i>Ramphocelus bresilius</i> (Linnaeus, 1766)<br><i>R. b. bresilius</i> (Linnaeus, 1766)*     | White-lined Tanager      | R                   | LC; LC                        | LSUMZ 10423 | WA 82978   | Minns <i>et al.</i> (2010) |
| <i>Charitospiza eucosma</i> Oberholser, 1905<br><i>Tersina viridis</i> (Illiger, 1811)<br><i>T. v. viridis</i> (Illiger, 1811)*            | Brazilian Tanager        | R, E                | LC; LC                        |             | WA 2761950 |                            |
| <i>Cyanerpes cyaneus</i> (Linnaeus, 1766)<br><i>C. c. holti</i> Parkes, 1977*  | Coal-crested Finch       | R                   | LC; NT                        | COUFRN 815  | WA 1689389 |                            |
| <i>Dacnis cayana</i> (Linnaeus, 1766)<br><i>D. c. paraguayensis</i> Chubb, 1910*   | Swallow Tanager          | R                   | LC; LC                        |             | WA 287757  |                            |
| <i>Coereba flaveola</i> (Linnaeus, 1758)<br><i>C. f. chloropyga</i> (Cabanis, 1850)*   | Red-legged Honeycreeper  | R                   | LC; LC                        |             | WA 1571335 |                            |
| <i>Sporophila lineola</i> (Linnaeus, 1758)<br><i>Sporophila nigricollis</i> (Vieillot, 1823)<br><i>S. n. nigricollis</i> (Vieillot, 1823)* | Blue Dacnis              | R                   | LC; LC                        | COUFRN 605  | WA 5200    | XC 429377                  |
| <i>Sporophila albogularis</i> (Spix, 1825)<br><i>Sporophila leucoptera</i> (Vieillot, 1817)<br><i>S. l. cinereola</i> (Temminck, 1820)     | Bananaquit               | R                   | LC; LC                        | COUFRN 92   | WA 193677  | WA 1254349                 |
|  | Lined Seedeater          | R                   | LC; LC                        |             | WA 336521  | WA 2925073                 |
|  | Yellow-bellied Seedeater | R                   | LC; LC                        |             | WA 237326  |                            |
|  | White-throated Seedeater | R, E                | LC; LC                        | UFPE 3144   | WA 47522   | WA 1254722                 |
|  | White-bellied Seedeater  | R                   | LC; LC                        |             | WA 2676962 |                            |



| Taxon   | English name             | National occurrence | National; Global conservation | Specimen    | Photo      | Sound      |
|---|--------------------------|---------------------|-------------------------------|-------------|------------|------------|
| <i>Sporophila bouvreuil</i> (Statius Muller, 1776)        | Copper Seedeater         | R                   | LC; LC                        | UFPE 3675   | WA 315239  |            |
| <i>Coryphaspiza melanotis</i> (Temminck, 1822)            | Black-masked Finch       | R                   | EN; VU                        | COUFRN 824  | WA 1689384 |            |
| <i>C. m. melanotis</i> (Temminck, 1822)*                  |                          |                     |                               |             |            |            |
| <i>Emberizoides herbicola</i> (Vieillot, 1817)            | Wedge-tailed Grass-Finch | R                   | LC; LC                        |             | WA 2220061 | WA 2051446 |
| <i>E. h. herbicola</i> (Vieillot, 1817)*                  |                          |                     |                               |             |            |            |
| <i>Saltatricula atricollis</i> (Vieillot, 1817)           | Black-throated Saltator  | R                   | LC; LC                        |             | WA 2658215 | WA 2909323 |
| <i>Saltator maximus</i> (Statius Muller, 1776)            | Buff-throated Saltator   | R                   | LC; LC                        |             | WA 2251953 | WA 2263320 |
| <i>S. m. maximus</i> (Statius Muller, 1776)*              |                          |                     |                               |             |            |            |
| [ <i>Saltator similis</i> d'Orbigny & Lafresnaye, 1837]   | Green-winged Saltator    | R                   | LC; LC                        |             |            |            |
| <i>Thlypopsis sordida</i> (d'Orbigny & Lafresnaye, 1837)  | Orange-headed Tanager    | R                   | LC; LC                        | COUFRN 145  | WA 256608  |            |
| <i>T. s. sordida</i> (d'Orbigny & Lafresnaye, 1837)*      |                          |                     |                               |             |            |            |
| <i>Cypsnagra hirundinacea</i> (Lesson, 1831)              | White-rumped Tanager     | R                   | LC; LC                        | COUFRN 902  | WA 2432542 |            |
| <i>C. h. pallidigula</i> Hellmayr, 1907                   |                          |                     |                               |             |            |            |
| <b>Cardinalidae</b>                                       |                          |                     |                               |             |            |            |
| <i>Piranga flava</i> (Vieillot, 1822)                     | Hepatic Tanager          | R                   | LC; LC                        |             | WA 2365584 |            |
| <i>P. f. saira</i> (Spix, 1825)*                          |                          |                     |                               |             |            |            |
| <i>Cyanoloxia brissonii</i> (M. H. C. Lichtenstein, 1823) | Ultramarine Grosbeak     | R                   | LC; LC                        | UFPE 3441   | WA 251708  |            |
| <i>C. b. brissonii</i> (M. H. C. Lichtenstein, 1823)*     |                          |                     |                               |             |            |            |
| <b>Fringillidae</b>                                       |                          |                     |                               |             |            |            |
| [ <i>Spinus yarrellii</i> (Audubon, 1839)]                | Yellow-faced Siskin      | R                   | VU; VU                        |             |            |            |
| <i>Euphonia chlorotica</i> (Linnaeus, 1766)               | Purple-throated Euphonia | R                   | LC; LC                        |             | WA 251718  | WA 3300907 |
| <i>E. c. chlorotica</i> (Linnaeus, 1766)*                 |                          |                     |                               |             |            |            |
| <i>Euphonia violacea</i> (Linnaeus, 1758)                 | Violaceous Euphonia      | R                   | LC; LC                        |             | WA 3257429 | WA 3257457 |
| <i>E. v. aurantiicollis</i> Bertoni, 1901*                |                          |                     |                               |             |            |            |
| <b>Estrildidae</b>  |                          |                     |                               |             |            |            |
| <i>Estrilda astrild</i> (Linnaeus, 1758)                  | Common Waxbill           | R                   | NE; LC                        | COUFRN 89   | WA 725795  | WA 2952512 |
| <i>E. astrild</i> ssp.                                    |                          |                     |                               |             |            |            |
| <b>Passeridae</b>   |                          |                     |                               |             |            |            |
| <i>Passer domesticus</i> (Linnaeus, 1758)                 | House Sparrow            | R                   | NE; LC                        | COUFRN 1089 | WA 561857  |            |
| <i>P. d. domesticus</i> (Linnaeus, 1758)*                 |                          |                     |                               |             |            |            |

Following Piacentini *et al.* (2015) we have mentioned subspecies, but these should not be taken as a taxonomic validation. Entries with an asterisk (\*) are based only on the current known distribution; therefore, these subspecies should not be used for biogeographical purposes. Those without an asterisk were specifically identified to taxon by us. In the light of some recent taxonomic reviews, we discuss a few issues in the species accounts, but do not specifically propose any changes to the list in Piacentini *et al.* (2015).

Conservation status refers both to national (ICMBio / MMA 2018) and global levels (IUCN 2019). National status and endemism follow Piacentini *et al.* (2015), but only for species on the Brazilian primary list.

We provide complementary information on some vouchers, details and comments on single records, and explanatory notes for all species on the secondary list. We have mentioned the few recoveries, recaptures and resightings of ringed birds of polytypic species encountered in the state, to indicate subspecies and to provide phenological data. We offer more detailed information on seabirds, a group less well known to most ornithologists, including available specimens, records of lost specimens, recoveries, other documented records and sight records.

We have given a strict definition to some terms frequently used in this text: 'specimen' = a skin or any part of a bird preserved in a collection; 'voucher' = unequivocal proof of identification of a taxon; 'recovery' = reading of any distinctive marking on a dead bird; 'recapture' = a ringed bird trapped and released again; and 'resighting' = reading of a ring, flag or tag on a live bird that was not recaptured.

## Results

We have compiled records of 425 bird species for the state of Rio Grande do Norte, 391 for the primary list—with available physical evidence—and 34 for the secondary list, with no documentary evidence. Subspecies are proposed for 272 species on the primary list, 27 of them identified, but the other 245 based on distribution. Four species are represented by two subspecies: Rusty-margined Guan *Penelope superciliaris*, Cabot's Tern *Thalasseus acuflavidus*, Peregrine Falcon *Falco peregrinus* and Brown-chested Martin *Progne tapera*. This results in a total of 428 valid or potentially valid distinct taxa (species and subspecies) recorded in the state (Table 1).

A total of 370 bird species that occur in the state are known or presumed to breed in Brazil, 35 of them country endemics (Piacentini *et al.* 2015). Another 51 are migrants from the Northern Hemisphere (36) or southern South America (11), the remaining four being accidental visitors or vagrants from the north. The former status of one species (*Phoenicopterus ruber*) is unknown. Three species introduced from the Old World have established self-sustaining populations: Rock Pigeon *Columba livia*, Common Waxbill *Estrilda astrild* and House Sparrow *Passer domesticus*.

We offer documentation for 12 species new to the state: Black-bellied Storm Petrel *Fregetta tropica*, Jabiru *Jabiru mycteria*, Stripe-backed Bittern *Ixobrychus involucris*, Roseate Spoonbill *Platalea ajaja*, Rufous Crab Hawk *Buteogallus aequinoctialis*, Pectoral Sandpiper *Calidris melanotos*, Short-eared Owl *Asio flammeus*, Lesser Nighthawk *Chordeiles acutipennis*, White-lored Tyrannulet *Ornithion inerme*, Spectacled Tyrant *Hymenops perspicillatus*, Chilean Swallow *Tachycineta leucopyga* and Violaceous Euphonia *Euphonia violacea*.

Noteworthy documented records include three significant range extensions: *Tachycineta leucopyga* (c.1,900 km north), *Hymenops perspicillatus* (c.1,400 km north) and *Asio flammeus* (c.1,000 km north-east). We also present the first proven breeding by Least Tern *Sternula antillarum*, *Thalasseus acuflavidus* and Creamy-bellied Thrush *Turdus amaurochalinus* in extreme north-eastern Brazil.

On the primary list, there are 19 globally threatened and Near Threatened species (4.9% of the list), one-third of them passerines, of which one is Critically Endangered, five Vulnerable, and 13 Near Threatened (IUCN 2019).

At the national level (ICMBio / MMA 2018), several taxa not evaluated by IUCN (2019) are of conservation concern, such as Yellow-legged Tinamou *Crypturellus noctivagus zabele* (VU), *Penelope supercilialis alagoensis* (CR), Wilson's Plover *Charadrius wilsonia crassirostris* (VU), American Whimbrel *Numenius hudsonicus* (NT), Amazonian Motmot *Momotus momota marcgravianus* (EN), Black-cheeked Gnatcatcher *Conopophaga melanops nigrifrons* (VU), Plain Xenops *Xenops minutus alagoanus* (VU), White-throated Spadebill *Platyrrinchus mystaceus niveigularis* (VU) and White-bellied Tody-Tyrant *Hemitriccus griseipectus naumburgae* (VU). In addition, several species, including a Brazilian endemic, have a poorer national status than the global designation, including Trindade Petrel *Pterodroma arminjoniana* (CR vs. VU), Red-footed Booby *Sula sula* (EN vs. LC), King Vulture *Sarcoramphus papa* (NT vs. LC), American Oystercatcher *Haematopus palliatus* (NT vs. LC), Short-billed Dowitcher *Limnodromus griseus* (CR vs. LC), Ruddy Turnstone *Arenaria interpres* (NT vs. LC), Red Knot *Calidris canutus* (CR vs. NT), Semipalmated Sandpiper *C. pusilla* (EN vs. NT), Roseate Tern *Sterna dougallii* (VU vs. LC), Turquoise-fronted Parrot *Amazona aestiva* (NT vs. LC), Ceara Gnatcatcher *Conopophaga cearae* (EN vs. LC) and Black-masked Finch *Coryphospiza melanotis* (EN vs. VU). On the other hand, the conservation prospects of some species, including four Brazilian endemics, are considered of less concern in a national setting, e.g., Sooty Shearwater *Puffinus griseus* (LC vs. NT), Leach's Storm Petrel *Oceanodroma leucorhoa* (LC vs. VU), White-collared Kite *Leptodon forbesi* (EN vs. CR), Tawny Piculet *Picumnus fulvescens* (LC vs. NT), Pectoral Antwren *Herpsilochmus pectoralis* (LC vs. VU), Red-shouldered Spinetail *Synallaxis hellmayri* (DD vs. NT), Blue Finch *Porphyrospiza caerulescens* (LC vs. NT), Bicoloured Conebill *Conirostrum bicolor* (LC vs. NT) and Coal-crested Finch *Charitospiza eucosma* (LC vs. NT). The Near Threatened Bar-tailed Godwit *Limosa lapponica* (IUCN 2019) has not been evaluated at national level. Moreover, in addition to *Synallaxis hellmayri*, five taxa are considered Data Deficient at national level, *Penelope supercilialis ochromitra*, Yellow-breasted Crake *Porzana flaviventer*, American Golden Plover *Pluvialis dominica*, Least Sandpiper *Calidris minutilla* and Pygmy Nighthawk *Nyctidromus hirundinaceus*.

## Annotated species on the Primary List

All localities are in Rio Grande do Norte, unless otherwise stated.

### GREATER RHEA *Rhea americana*

Greater Rhea was once common (Marcgrave 1648), but the native population, mentioned from many historical localities, became extinct sometime between 1913 (Alves 1977) and 1959 (Faria 1961). The name *R. a. macrorhyncha* P. L. Sclater, 1860, from north-east Brazil (Sick 1993, 1997), is not generally accepted by recent works (Piacentini *et al.* 2015, ICMBio / MMA 2018, Gill & Donsker 2019), but taxonomic and distributional limits of the various subspecies require revision (Piacentini *et al.* 2015). An unidentified taxon was reintroduced in the 1990s at Seridó Ecological Station (ESEC Seridó) (06°35'S, 37°15'W; 200–385 m), Serra Negra do Norte, where a small feral population persists, and subsequently in several private and protected areas in the state, where some still thrive.

### YELLOW-LEGGED TINAMOU *Crypturellus noctivagus*

Previously considered probably extinct in the state based on interviews conducted by FSM & DMH with local people in Goianinha, who referred to it as 'zabelê', this tinamou has recently been 'rediscovered'. The first proof of its continued presence was obtained

in a forest fragment (05°48'S, 35°21'W) near Natal International Airport, São Gonçalo do Amarante, on 19 December 2018 (WA 3217181; R. Araújo). This locality appears too small and isolated to have conservation potential. Although we follow Piacentini *et al.* (2015), a recent reassessment recommended that *C. n. zabele* be treated at species level (Tomotani & Silveira 2016).

#### **FULVOUS WHISTLING DUCK** *Dendrocygna bicolor*

A pair introduced in ESEC Seridó, Serra Negra do Norte, was seen breeding in 2002 by FSM, but not in 2012–16 (MP pers. obs.). Previously mentioned for the state (Silveira & Varela-Freire 1999, Azevedo *et al.* 2004, Silva *et al.* 2012), but not documented until a group was photographed at Augusto Severo on 4 September 2015 (WA 2168387; C. Brito) and 13 December 2015 (WA 1951196; J. van den Bosch). Partially migratory in Brazil (Antas 1994, Sick 1997), rare or occasional in many places (Sick 1997), but resident in nearby Ceará (Somenzari *et al.* 2018). Rio Grande do Norte records probably involve southern migrants, but it is also a potential breeder.

#### **RUSTY-MARGINED GUAN** *Penelope superciliaris*

Probably two taxa of the *P. superciliaris* complex occur in RN: the Critically Endangered *P. s. alagoensis* (ICMBio / MMA 2018) in the Atlantic Forest and *restinga* (L. F. Silveira pers. comm.; WA 515077; E. Silva), with perhaps the Data Deficient *P. s. ochromitra* Neumann, 1933 (not Spix, 1825 as in Piacentini *et al.* 2015), in the interior. The latter is not confirmed, but *Penelope* flight feathers collected by FSM at Serra das Queimadas, Parelhas, and Serra do Salobro, Luís Gomes, could be identified at MZUSP as only “not *P. jacucaca*” (L. F. Silveira pers. comm.). A recent review of the complex suggests that these taxa should be treated as full species (Vargas 2017).

#### **WHITE-BROWED GUAN** *Penelope jacucaca*

Just two photo-documented records of this Vulnerable species: at Tibau on 17 June 2010 (WA 2016795; B. França) and Ruy Barbosa on 5 March 2017 (WA 2491752; A. Barbosa). The status of this Caatinga endemic is very poorly known in Rio Grande do Norte; interviews with local people by FSM suggested some local extinctions, but these do not certainly refer to *P. jacucaca*, as two species of guans appear to frequent the Caatinga (see above).

#### **SOUTHERN FULMAR** *Fulmarus glacialis*

Three specimens (MCZ 186306, 186307 and 226729) collected by G. H. Mackay at Cabo São Roque, c.3 km north of the Maxaranguape River mouth (05°29'S, 35°16'W), Maxaranguape, on 9 June 1863 (Carlos *et al.* 2004) are the only state records.

#### **CAPE PETREL** *Daption capense*

An adult *D. c. capense* (MCZ 226730), collected by G. H. Mackay at Cabo São Roque near the mouth of the Maxaranguape River (05°29'S, 35°16'W), Maxaranguape, on the same date as the previous species, another austral migrant (Carlos *et al.* 2004), is the sole state record.

#### **CORY'S SHEARWATER** *Calonectris borealis*

Two recoveries of birds originally ringed in Macaronesia: a nestling (CEMPA L41317) on Selvagem Grande, Madeira, on 17 September 1991, found by P. H. C. Cordeiro at Pirangi beach, Parnamirim, on 11 July 1994, c.4,490 km from the banding site; and one, banded (Spain 6123288) in the Canary Islands on 2 November 2000, found at Extremoz (05°40'S, 35°W) on 12 January 2001 (Olmos 2002). Two other dead birds, identified using Lima *et al.*

(2004), were found by FSM, of which the mummified remains of one at Canto do Amaro (05°05'S, 36°20'W), Guamaré, on 17 December 2010, will be deposited at COUFRN. Live birds: several photographed off the east coast between 24 November and 22 December 2006 (L. França; confirmed by F. Olmos), one close to the north coast at Guamaré on 6 January 2012 (WA 2518923; MP), another at Galinhos on 31 December 2014 (WA 1569868; F. Torres), and 20 seen 27 km east-northeast off Natal (05°43'S, 34°57'W; depth 200 m) on 24 January 2015 (WA 1593241; N. Moura; Lees *et al.* 2015). Also sporadically observed flying over the mangroves of the north coast, at Salina Diamante Branco (05°05'S, 36°17'W), Galinhos (Azevedo *et al.* 2004), and Canto do Amaro on 19 December 2012 (FSM). It appears that *C. borealis* is regular offshore during the boreal winter, with the immature recovered in July. This is consistent with birds from Macaronesia and Mediterranean wintering in the western Atlantic, where the species has been found year-round (Lima *et al.* 2004, Somenzari *et al.* 2018).

### SOOTY SHEARWATER *Puffinus griseus*

One photographed offshore (05°48'S, 34°56'W) on 9 December 2017 (ML 77958651, 77958661; F. Olmos) was the first state record, although this transequatorial migrant was expected given the discovery of one beached on the main island of Fernando de Noronha National Park in 1995 (MZUSP 75463) (Schulz-Neto 2004a, Silva e Silva 2008). Many seen with Manx Shearwaters *P. puffinus* off Trairi, Ceará, on 14 September 2018 (WA 3111847; C. E. Moura). These three records in extreme north-east Brazil are all beyond the species' regular northern limit off Brazil, the state of Bahia (Somenzari *et al.* 2018).

### GREAT SHEARWATER *Puffinus gravis*

Of two beached individuals found by JBI at Senador Georgino Avelino in December 2004, the fresher bird was deposited at COUFRN, as was one found at Via Costeira beach, Natal, donated by the environmental agency IBAMA / Natal prior to 2008; however, both frozen corpses were effectively destroyed by an electricity outage at COUFRN on 1 February 2009. The mummified remains of one found by FSM beached at Chapadão de Pipa, Tibau do Sul, on 31 December 2007, will be deposited at COUFRN. Two undated specimens, donated by Aquário-Natal, lack precise localities: COUFRN 838, and female, COUFRN 949, dated 24 September 2012, although it was probably found earlier than this. Recently photographed at sea off Rio Grande do Norte: adults (WA 2813943; J. Quental, WA 2844237; E. Ramirez) c.20 km off Natal on 10 December 2017, and another on 13 May 2018 (WA 2971931; M. Farah). *P. gravis* breeds on islands in the South Atlantic in November–April, moves north of the equator between March and September, and visits waters off north-east Brazil in March–December (Lima *et al.* 2004, Somenzari *et al.* 2018), which is consistent with the six dated records here.

### MANX SHEARWATER *Puffinus puffinus*

First known from the discovery of beached individuals: one at Barra do Cunhaú, Tibau do Sul, on 24 December 2002; one at Cotovelo, Parnamirim, on 11 December 2004 (FSM); COUFRN 831, found by R. Gondim at Forte dos Reis Magos, Natal, on 23 September 2005; one at Chapadão de Pipa, Tibau do Sul, on 1 January 2008 (FSM); an undated female at Via Costeira, Natal, donated by IBAMA / Natal in 2008, but effectively destroyed by an electricity outage at COUFRN; and eight along 1.5 km of beach at Búzios, Nísia Floresta, on 29 May 2012 (C. Ferreira, identified by FSM). There is an unpublished recovery of a fully grown individual, ringed (EW 77936) at Mallaig (57°N, 05°49'W) in western Scotland by A. D. K. Ramsay on 20 September 2009, found dying by V. Andrade on Cacimbinha



beach, Tibau do Sul, on 15 January 2013, c.7,525 km from the banding site. Hundreds were observed (some photographed) at the shelf break, 27 km east-northeast off Natal, on 25 January 2015 (Lees *et al.* 2015). One flying south 200 m offshore at Santa Rita, Extremoz, on 1 November 2009 (FSM). This species frequents Brazilian and Argentine coasts during the boreal winter (December–March), and breeds in the north-east Atlantic, with isolated records from July and November in Bahia (Somenzari *et al.* 2018). Our data show that *P. puffinus* frequents eastern coastal waters between September and January, with one record in May, thus, is more common off Rio Grande do Norte than previously supposed (Mestre *et al.* 2010, Somenzari *et al.* 2018).

### BLACK-BELLIED STORM PETREL *Fregetta tropica*

A beached female with just one leg (COUFRN 120; Fig. 2) was found by the environmental police at Ponta Negra, Natal, on 3 August 2009. The large black hood and its ragged border, which are the best features to distinguish this species from White-bellied Storm Petrel *F. grallaria* (Howell 2010), identify the specimen as *F. tropica*, while the spotted underparts, rather than pure white belly of *F. t. melanoleuca*, indicates the nominate subspecies (Howell 2010, Robertson *et al.* 2016). Note that toe projection and the presence of white feathers on the chin and throat are not necessarily useful for species identification, while the black leading edge of the underwing, narrower in *F. grallaria* (Howell 2010), is not fully visible in our specimen. The only state record, and the fourth specimen of *F. tropica* for Brazil (Lima *et al.* 2004). This austral migrant had already been reported from far north-east Brazil (Olmos 2000): during the 19th century c.30 were observed around a ship some 290 km from Fernando de Noronha archipelago (Silva e Silva 2008). *F. t. tropica* presumably frequents Brazilian waters mainly during the austral winter (Lima *et al.* 2004, Somenzari *et al.* 2018); a record from Bahia in January (Olmos 2000), i.e. the austral summer, may have involved an immature.



Figure 2. Black-bellied Storm Petrel *Fregetta t. tropica*, COUFRN 120, found in Natal, Rio Grande do Norte, Brazil, August 2009 (Rafael Dantas Lima)

### WILSON'S STORM PETREL *Oceanites oceanicus*

The first documentation of this austral migrant in state waters was published by Lees *et al.* (2015) from over the slope (05°43'S, 34°53'W; depth 2,400 m), which is outside the boundary we have defined but very close to the shelf break (05°43'S, 34°57'W; depth 200 m). Another photo also taken on 24 January 2015, is from the same general area, 27 km east-northeast off Natal (WA 1593174; A. Lees). Previously, one was observed <20 km off Natal during a trip to Atol das Rocas in 1994 (A. Schulz-Neto). It is unknown which of the two subspecies occurs: the nominate, which breeds on subantarctic islands, or *O. o. exasperatus* from Antarctica. Both occur in Brazil (Piacentini *et al.* 2015), where previously recorded between March and

November (Lima *et al.* 2004, Somenzari *et al.* 2018); the January records from Rio Grande do Norte extend its temporal presence in the country's waters.

### LEACH'S STORM PETREL *Oceanodroma leucorhoa*

A specimen of this Vulnerable species (COUFRN 728), donated by IBAMA / Natal, was found on the state's north coast, either on a petroleum platform or a beach, and prepared as a specimen on 6 September 2012 by MP. Two were photographed by A. Lees (WA 1593164) and F. Olmos at the shelf break (05°43'S, 34°57'W; depth 200 m) on 24 January 2015 (Lees *et al.* 2015). Sick (1985) cited a March record for the state, albeit perhaps from Atol das Rocas, which is sometimes considered part of Rio Grande do Norte, but it was not mentioned in Sick (1993, 1997). Part of the Northern Hemisphere population winters off Brazil, with records from Amapá south to São Paulo, mainly in January–February (Somenzari *et al.* 2018).

### JABIRU *Jabiru mycteria*

The presence of Jabiru in the state was based on an undocumented old record until 2012. Sr. Raimundo, when 73 years old, informed FSM on 11 September 2011 that in the rainy 1960s 'tuiuiús' (*J. mycteria*, which he described well) reached Acari, and one was hunted. "We ate it. The flesh is black, as tasty as that of a duck", he said. The first documented record involved one on an islet in Pau dos Ferros Dam (06°10'S, 38°11'W), Pau dos Ferros, in February 2012 (Fig. 3A), followed by two at Carnaubais (05°12'S, 36°47'W) on 28 February 2018 (Fig. 3B). Around the same time other records were made in Pernambuco (Pereira *et al.* 2012) and Ceará (WA 3194160) confirming the species' occasional occurrence in the Caatinga, during wet periods.




Figure 3A. Jabiru *Jabiru mycteria*, Pau dos Ferros, Rio Grande do Norte, Brazil, February 2012 (© J. L. C. Novaes); (B) Carnaubais, Rio Grande do Norte, Brazil, February 2018 (© C. T. Puppim-Gonçalves & A. C. C. Xavier)

### WOOD STORK *Mycteria americana*

Three in a mixed colony of several species of white herons at Umari, Caicó in the Caatinga, in 1998 (C. Medeiros) represented the first state record. An adult with heavily worn and disorganised wing feathers and a dirty body was photographed flying over Messias Targino on 13 December 2015 (WA 1945521; J. van den Bosch).

### MAGNIFICENT FRIGATEBIRD *Fregata magnificens*

There is an unpublished recovery of this frigatebird banded as a nestling (CEMAVE U07889) on Fernando de Noronha (03°52'S, 32°26'W) on 25 June 1987 by A. Filippini, and recovered at Pitangui beach (05°30'S, 35°10'W), Extremoz, sometime prior to 1997 (SNA



2019; A. Schulz-Neto pers. comm.). Regular visitor to the state's coasts, being well known to fishermen as 'garapira'. Its seasonal distribution, based on 52 records (some documented), shows individuals (including adult males) and groups of up to nine occur year-round. At Baía Formosa, flocks of 16 and 15 were seen by FSM interacting with fishing Guiana Dolphins *Sotalia guianensis* in December 2002 and January 2003, respectively.

#### **MASKED BOOBY** *Sula dactylatra*

An adult male found alive on Santa Rita beach, Extremoz, on 8 October 2008, was taken to Natal Zoo (Fig. 4), where it subsequently died. It was only recently prepared as a specimen (COUFRN 1253). The first available documented records were off Natal at the shelf break (05°43'S, 34°57'W; depth 200 m) (Lees *et al.* 2015). There are three unpublished recoveries in the state: a nestling, ringed (CEMAVE U23323) on 19 July 1994 on Santa Bárbara Island, Abrolhos National Park (17°58'S, 38°42'W), Caravelas, Bahia, caught at sea (05°06'S, 36°38'W) north of Macau in 1997; an adult, ringed on the right leg, said to be from Abrolhos, Bahia, landed on a floating platform at the Maracajaú Reef (05°40'S, 35°24'W; depth 5 m), 7 km off Maxaranguape, where it was fed, in 2000 and 2002 (Risonaldo pers. comm.; photo identified by FSM); and an adult with a ring on its right leg, photographed on Búzios beach, Nísia Floresta—not in Natal as indicated—(see WA 549593) and subsequently released by the environmental police (G. Rodrigues pers. comm.). *S. dactylatra* appears to be the commonest of the three *Sula* species that occur in the state's waters.



Figure 4. Masked Booby *Sula dactylatra*, found in Extremoz, Rio Grande do Norte, Brazil, October 2008, photographed in Natal Zoo, and now COUFRN 1253 (© D. Brandão)

#### **RED-FOOTED BOOBY** *Sula sula*

An adult in typical plumage was found on the coast in 2018 and was prepared as a specimen (COUFRN 1275) on 28 September 2018. This is the only state record.

#### **PINNATED BITTERN** *Botaurus pinnatus*

One struck the window of JBI's house close to the floodplain of the Pirangi River, Parnamirim, on 28 May 2008. Its frozen corpse was effectively destroyed by an electricity outage at COUFRN. This was the only state record for some years, but live individuals were subsequently photographed (see WikiAves 2019).

#### **LEAST BITTERN** *Ixobrychus exilis*

One observed at Povoado de Sítio in Ceará-Mirim between February 2001 and June 2002 (M. Silva) was the first state record. A female at the lagoon in Martins on 6 August 2010 was the second (FSM). The first documented record involved an adult male at the main pond in ESEC Seridó, Serra Negra do Norte, in 2012, photographed by C. A. Varela-Freire ([www.carlosvarela-agenteambiental.blogspot.com](http://www.carlosvarela-agenteambiental.blogspot.com)). Finally, an adult female was at Mossoró on 25 December 2016 (WA 2411673; RDL). The species' status in the state is unclear, but the record in December suggests possible local breeding.

**STRIPE-BACKED BITTERN** *Ixobrychus involucris*

The only state record is a photo of an adult flushed from cattails in the mobile sand dunes at Guamaré on 18 May 2012 (WA 3065361; C. Ferreira, identified by FSM).

**ROSEATE SPOONBILL** *Platalea ajaja*

The only documented state record involves a skull (SDNHM 35485) collected by F. Baker at 'Papary Lake' (06°07'S, 35°10'W), Nisia Floresta (Fig. 5). Although the date of the specimen is not given, it can be attributed to 1911, corresponding to the date of a mollusc-collecting expedition during which it was discovered (Baker 1913). Interviews with local people convinced FSM as to the presence of this species in mangroves at Macau and Guamaré until the 1960s, but suggest that it is now extinct; however, one, probably of this species, was observed flying over Macau around December 2009 (JBI). The nearest recent documented records to Rio Grande do Norte are from north-west Ceará and Alagoas, meaning the species could wander to the state.



Figure 5. Skull of Roseate Spoonbill *Platalea ajaja*, SDNHM 35485, collected in Nisia Floresta, Rio Grande do Norte, Brazil, 1911 (P. Unitt, © San Diego Natural History Museum)

**OSPREY** *Pandion haliaetus*

Widely reported in the state: Macau (Andrade 1991, Sick 1997); Seridó (Varela-Freire & Araújo 1997); the littoral around Potiguar petroleum basin: Macau, Assu, Galinhos, Guamaré, Porto do Mangue (Varela *et al.* 1997, Azevedo *et al.* 2004, Irusta & Sagot-Martin 2011, Silva *et al.* 2012); and Grossos (Girão & Albano 2011). However, there was no documented record for the state until an adult was photographed in flight at Galinhos on 21 November 2009 (WA 252946; B. França). Another record for the southern interior was at ESEC Seridó, Serra Negra do Norte (Pichorim *et al.* 2016b). A female *P. h. carolinensis* named 'Holly' banded (1088-07037) and tagged (388) as a young adult, along with its mate, in Annapolis (38°60'N, 76°29'W), Maryland, USA, was also fitted with a datalogger (Fig. 6A),



Figure 6A. Osprey *Pandion haliaetus* ('Holly'), Annapolis, Maryland, USA (© R. Bierregaard); and (B) tracking of 'Holly'.

which revealed that the bird twice migrated to Rio Grande do Norte, covering c.8,000 km on each journey. ‘Holly’ remained in the Piranhas basin (05°44’S, 36°53’W) between 16 October 2016 and 15 March 2017 and from 9 November 2017 to 14 March 2018 (Fig. 6B). On the next migration, ‘Holly’ wintered in the state of Ceará, in the Acaraú basin (04°14’S, 40°27’W) (R. Bierregaard pers. comm.). North American Ospreys had previously been recovered in the neighbouring states of Paraíba, initially on 4 February 1973, and Ceará, on 10 February 1983 (Mestre & Bierregaard 2009), but not in Rio Grande do Norte (Mestre *et al.* 2010).

#### WHITE-COLLARED KITE *Leptodon forbesi*

A presumed adult was flushed from the ground on the left bank of Jiqui River, then circled overhead with a Black Vulture *Coragyps atratus* before disappearing behind forest at Parnamirim, on 15 January 2007 (FSM). This first state record of an Endangered north-east Brazilian endemic was originally considered doubtful due to possible confusion with White-necked Hawk *Amadonastur lacernulatus*, which is now known to occur only south of the São Francisco River (Dénes *et al.* 2011). FSM subsequently observed an adult and young *L. forbesi* at the same locality, on 20 July (with JBI) and 22 July 2009, respectively. Proof of occurrence was obtained by a high-flying bird photographed at Baía Formosa on 4 September 2016 (WA 2263455; D. Gurgel).

#### SHARP-SHINNED HAWK *Accipiter striatus*

The first documented state record involved an adult banded (CEMAVE H110936) at the Universidade Federal Rural do Semi-Árido farm (05°03’S, 37°24’W), Mossoró, on 26 April 2014 (L. França & P. T. Moura; Fig. 7A). Another Caatinga record was of one captured by MP at ESEC Seridó, Serra Negra do Norte, on 12 June 2014 (Fig. 7B). The subspecies involved is *A. s. erythronemius*, with a finely barred white chest, rufous-grey belly and red flanks and thighs (Ferguson-Lees & Christie 2001), which is sometimes treated as a species (Gill & Donsker 2019).



Figure 7A. Sharp-shinned Hawk *Accipiter striatus erythronemius*, captured in Mossoró, Rio Grande do Norte, Brazil, April 2014 (© P. T. Moura); (B) Serra Negra do Norte, Rio Grande do Norte, Brazil, June 2014 (Mauro Pichorim)

#### PLUMBEOUS KITE *Ictinia plumbea*

The first state record was of four flying south-east over Serra do Feiticeiro, Lajes, on 6 March 2006 (FSM), while one perched beside the road to Jardim do Seridó (06°39’S, 36°42’W), Parelhas, on 10 April 2006 (photographed by L. Sena) may pertain to the same migratory movement. Further records suggest overwintering adults from a southern population: singles at the Jiqui and Pitimbu floodplain, Parnamirim and Nísia Floresta, on 20 July 2009, and in the same area (WA 249719; B. França) on 5 August 2009 (FSM & JBI), with two there



on 11 July 2016 (FSM). More surprisingly, two recent records of juveniles: one over São Miguel do Gostoso on 27 May 2018 (WA 3010251; T. Almeida) and one over Currais Novos on 17 February 2019 (WA 3282753; G. Medeiros), the latter date suggestive of local breeding. In Brazil, there are resident and migratory populations of *I. plumbea*, which depart the south of the country mainly in May–July, and occur sporadically in the north-east, where it is not known to breed (Somenzari *et al.* 2018).

#### **RUFOUS CRAB HAWK** *Buteogallus aequinoctialis*

In 2002, FSM became convinced that a population exists in the mangroves of Canguaretama, based on interviews with local people during a canoe trip on the Camamirim River. They spoke of the ‘gavião-piri-piri’ or ‘pixi-pixi’—onomatopoeic names for *B. aequinoctialis*—and pointed out a place where one had a nest. A calling bird was heard on the Camamirim River on 27 December 2002 and two were observed flying over the Penha River on 6 January 2003; an adult was seen in the same mangroves on 22 April 2005 (FSM). Ten years later, on the south side of this mangrove complex, at Baía Formosa, sightings were made in August and October 2015 (FSM), and it was finally documented by photos and video on the Garatuba River (06°23’S, 35°04’W) on 25 October 2015 (WA 2874182; DMH). It is estimated that the Rio Grande do Norte population of this Near Threatened species is just 5–10 pairs, with no records elsewhere in the state. In extreme north-east Brazil, it appears to have disappeared from Ceará (WikiAves 2019), where it was formerly present (Pinto & Camargo 1961). More recently, it was found in mangroves of the Mamanguape and Paraíba do Norte Rivers (Araujo *et al.* 2006), and elsewhere in nearby Paraíba (WikiAves 2019). Although recent records in mangroves in the Atlantic Forest domain are restricted to a few localities (Moreira-Lima 2013), we believe the species is recovering in parts of north-east Brazil, possibly due to the less systematic destruction of raptors than in previous decades.

#### **YELLOW-BREASTED CRAKE** *Porzana flaviventer*

An adult in wing moult was caught alive by a poacher’s hunting dog at Maiada do Fogo, in the floodplain of the Ceará-Mirim River, Ceará-Mirim, on 23 October 2005, and was given to M. Silva. Photos were taken next day by JBI (Fig. 8) before it was released at the same location. On 8 July 2007, FSM & M. Silva flushed one that briefly perched at Sítio (c.05°36’S, 35°30’W), Ceará-Mirim, in the same floodplain. One was photographed by D. Gurgel and P. Vitor in the same municipality on 23 October 2016 (WA 2336075, 2334287, 2333420). The status in Rio Grande do Norte of this Data Deficient crake (ICMBio / MMA 2018) remains to be clarified.



Figure 8. Yellow-breasted Crake *Porzana flaviventer*, captured in Ceará-Mirim, Rio Grande do Norte, Brazil, October 2005 (Jorge Bañuelos Irusta)

#### **PAINT-BILLED CRAKE** *Neocrex erythrops*

Few reports from north-east Brazil, where this crake is a seasonal visitor and a probable breeder during the rainy season (R. Otoch pers. comm.). In the state, adults have been found dead or moribund three times: a full-plumaged decaying corpse, which had been struck by a vehicle on the road to Cacimbinha, Tibau do Sul, in the Atlantic Forest, found by V.



Figure 9. Fresh corpse of Paint-billed Crane *Neocrex erythrops*, Parnamirim, Rio Grande do Norte, Brazil, March 2015, and currently COUFRN 1224 (© Z. Coringa)

Andrade on 2 May 2006, photographed by DMH and deposited in alcohol at COUFRN by FSM; one with a wound to its right eye, captured by hand and photographed immediately after release at Cajueiro stream, which divides the municipalities of Parelhas and Equador, in the Caatinga, on 5 June 2008 (WA 343447; L. Sena); and a female found dead in the early morning by M. Viana & Z. Coringa, either having struck a wall or a vehicle, at Pium (05°57'S, 35°10'W), Parnamirim, in the coastal Atlantic Forest on 23 March 2015 (COUFRN 1224) (Fig. 9). In Parelhas, an adult foraging on wet ground was photographed on 10 May 2018 (WA 2969079; L. Sena), and the *tiuk* call was sound-recorded during the same good rainy season on 2 June 2018 (WA 2992265; J. Dantas).

#### WILSON'S PLOVER *Charadrius wilsonia*

The South American resident population breeds on northern and eastern coasts of Rio Grande do Norte (Sick 1997; FSM & JBI pers. obs.); the presence of migrants from North America is yet to be proven. A specimen taken in Canguaretama sometime between September 1997 and November 1998 and deposited at UFPE (Nascimento & Larrazábal 2000) can no longer be found there (L. Naka pers. comm.). Additional studies are needed to define geographic boundaries and diagnosis of *C. w. crassirostris* and *C. w. cinnamominus*.

#### SOLITARY SANDPIPER *Tringa solitaria*

Three ringed as juveniles in North America were recaptured and released in the state: BBL (Bird Banding Laboratory, USA and Canada) 1151-15400, New Brunswick, Canada (45°43'N, 64°31'W), on 25 August 1988, recaptured at Canguaretama (06°20'S, 35°01'W) on 25 March 1995; BBL 1431-68689, New Brunswick, on 10 August 1999, recaptured at Galinhos (05°06'S, 36°10'W) on 24 October 2000; and BBL 1541-46316, New Jersey, USA (39°10'N, 74°51'W), ringed on 23 May 2000, recaptured at Galinhos on 13 November 2000 (Mestre *et al.* 2010; L. Mestre pers. comm.). Birds migrating through south-east Canada belong to the nominate form.

**WILLET** *Tringa semipalmata*

There has been one alleged recovery of a ringed Willet in the state (Lara-Resende & Leal 1982), questionable because it was mapped outside its coastal range. Forty photos taken on both the north and east coasts (WikiAves 2019) are mainly the nominate, but it is difficult to eliminate Western Willet *T. s. inornata* in some cases (O'Brien *et al.* 2006). Our voucher, an individual with an evenly heavy bill to its tip and contrastingly darker scapulars and paler coverts, photographed at Galinhos on 15 November 2014 (WA 1522100; J. van den Bosch), is of the eastern nominate. Some authors consider the races as separate species that can be syntopic on migration and in winter (Oswald *et al.* 2016).

**RUDDY TURNSTONE** *Arenaria interpres*

One, ringed in New Jersey, USA, in 2002—thus presumably *A. i. morinella*—was seen in Natal on 15 March and 6 May 2005 (FSM). Individuals were also re-trapped by S. Azevedo in Galinhos during 22 April–3 May 2003 (Araújo *et al.* 2004).

**RED KNOT** *Calidris canutus*

Three Red Knots originally ringed in New Jersey, USA (one in 2000 or 2002, and two in 2004; W. Pitts pers. comm.)—presumed to be *C. c. rufa*—were seen at the Casqueira River, Macau, on 10 January and 17 February 2005, respectively (JBI).

**SANDERLING** *Calidris alba*

There have been six sightings of four individuals, all marked in New Jersey, USA (thus presumably *C. a. rubida*), at Forte dos Reis Magos, Natal: one ringed in 2004, observed on 4, 20 and 23 November 2004 (FSM); one ringed in 2003, another in 2004, both seen on 15 December 2004 (JBI); and flag KMJ (banded FWS 1621-19137) attached at Reeds Beach on 14 May 2005, photographed on 30 September 2006 (MP).

**PECTORAL SANDPIPER** *Calidris melanotos*

The only record is of four observed by JBI in a monitored landfill (05°43'S, 35°23'W) at Ceará-Mirim on 27 September 2012 (ML 27817531).

**SOUTH POLAR SKUA** *Stercorarius maccormicki*

As distinguishing this species from some congeners can be difficult, the origin of marked individuals can be useful to guarantee identification: one, ringed prior to fledging at Palmer Station, Cormorant Islands, on 23 February 1980, was recaptured on the state's north coast on 30 October 1980 and released (Lara-Resende & Leal 1982), captured again on 26 April 1984 at the same locality and kept captive; and a nestling, ringed at Admiralty Bay, King George Island (62°S, 58°20'W) on 19 January 1986, recovered at Galinhos on 20 October 1994 (Olmos 2002). One undated specimen (COUFRN 182), donated by IBAMA / Natal, is from Rio Grande do Norte; a dark-morph immature in primary moult was caught at sea and kept captive in Macau, photographed in a cage on 12 September 2008 (JBI, identified by FSM); and one was photographed close to the coast of Natal on 28 October 2012 (WA 791473; V. Florencio). Three of these records are from the same offshore area. Although already proven to occur in the state's waters (Lara-Resende & Leal 1982, Olmos 2002), this skua was mentioned only as a potential species off extreme north-east Brazil (Lees *et al.* 2015). As with most jaegers, it is a transequatorial migrant that is expected to be fairly common in offshore waters (F. Olmos pers. comm.).



**POMARINE JAEGER** *Stercorarius pomarinus*

The first photographic record involved one around a boat, pursuing *Sula dactylatra* and *Puffinus puffinus*, 27 km east-northeast off Natal at the shelf break, where there were hundreds of terns of four species, on 25 January 2015 (Lees *et al.* 2015; WA 1593176; A. Lees). Two previous sight records by FSM: one moving south off Cotovelo beach, Parnamirim, on 15 November 2002, and another at Maracajaú Reef (05°40'S, 35°24'W), Maxaranguape, on 30 May 2005.

**PARASITIC JAEGER** *Stercorarius parasiticus*

The first documented record involved a full-plumaged dark-morph adult photographed c.20 km off Natal on 10 December 2017 (ML 77958391; F. Olmos). Previously, individuals had been seen by FSM at Maracajaú Reef (05°40'S, 35°24'W), Maxaranguape: one during 29–31 May 2005, one on 29–30 August 2005, a total of six moving south on 20–21 September 2006, and a total of five moving south on 18–19 October 2006, i.e. during both the boreal spring and autumn migrations.

**GREY-HOODED GULL** *Chroicocephalus cirrocephalus*

Hundreds of pairs of this colonial breeder occur in mixed colonies with Gull-billed Terns *Gelochelidon nilotica*, Cabot's Terns *Thalasseus acuflavidus eurygnathus* and Black-necked Stilts *Himantopus mexicanus* in or close to salt pans on the north coast (Azevedo *et al.* 2004; FSM, JBI & MP). It is recorded year-round, with fewest in December (five birds) and March (12) and largest numbers during the breeding period (April–August), with max. 1,647 nests (eggs, nestlings and a few flying juveniles) at Furado do Avelino, Macau, on 18–19 July 2009 (FSM, MP, B. França & C. Ferreira). Post-nuptial moult of adults noted August–February (FSM), and post-breeding dispersal at Galinhos, in 2012, between 25 September and 30 October (FSM & C. Ferreira). Uncommon on the state's east coast, where we are aware of just three records: at Nísia Floresta in August 2001 (JBI); at the Potengi estuary, Natal between August 2007 and March 2008 (B. França); and two adults photographed at Natal on 6 November 2013 (WA 1151661; R. Oliveira). There is an inland record of two non-breeding adults at Lagoa do Piató, 57 km from the coast, in Assu, on 9 December 2008 (FSM).

**LAUGHING GULL** *Leucophaeus atricilla*

An immature with moulting primaries photographed at Guamaré on the state's north coast on 16 May 2010 (WA 287755; B. França) was the first available documented record. Previously observed by FSM, but on the east coast at Maracajaú Reef (05°40'S, 35°24'W), Maxaranguape, where two first-years were present 12 November–6 December 2006 (Fig. 10). On the north coast, 12 (all non-adult) were at the Conchas estuary, Porto do Mangue, on 14 February 2009 (FSM & MP). Most records are from Guamaré: an adult with three first-years at Canto do Amaro (05°05'S, 36°20'W) on 9 November 2010; a first-year at the same location on 19 December 2010; a first-year between Canto do Amaro and Coroa do Moacir on 29 May



Figure 10. First-winter Laughing Gull *Leucophaeus atricilla*, Maxaranguape, Rio Grande do Norte, Brazil, December 2006 (© E. Vieira)



2011, and perhaps the same bird at the confluence of the Aratuá and Miassaba Rivers (05°06'S, 36°19'W) on 6 June 2011 (FSM); five (including a breeding-plumaged adult) on Canto do Amaro beach on 28 February 2015 (photographed by C. Ferreira, identified by FSM). Also on the north coast, a first-winter at Areia Branca on 20 January 2012 (WA 574897; C. Beier). It appears that this migrant regularly overwinters (November–February) on the state's coasts, where immatures are present as late as May or June.

### LEAST TERN *Sternula antillarum*

The alleged identification of Yellow-billed Tern *S. superciliaris* at Salina Diamante Branco (05°05'S, 36°17'W), Galinhos, was accompanied by the following comment: 'arrived March–April to breed in colony June–July 1999 and June 2000, one breeding adult collected 18 June 2000' (translated from Azevedo *et al.* 2004), and the same species was also reported by Araújo *et al.* (2004), but the specimen mentioned (UFPE 3139, in fact labelled 19 June 2000) is an adult *S. antillarum* (Fig. 11). Least Tern is a regular seasonal breeder on sand banks in some mangrove complexes on the state's north coast, as documented at Macau (courtship 29–30 April, one egg on 15 May 2005, and estimated seven pairs around Salina Soledade; FSM & JBI), Guamaré (80 individuals and 20 nests at Rolo da Costa on 29 May 2011; FSM & R. Dantas) and Galinhos (Azevedo *et al.* 2004). This species frequents coastal waters in dozens or hundreds year-round—except August, when only three recorded. Largest concentrations in January—hundreds at the shelf break off Natal (Lees *et al.* 2015)—and in April and October, with respective max. of 370 and 325 roosting on floating structures at Maracajaú Reef (05°40'S, 35°24'W), Maxaranguape (FSM). The presence of individuals from the Northern Hemisphere in these flocks is probable, but not documented. North American migrants could occur simultaneously with local breeders, at least in April.



Figure 11. Least Tern *Sternula antillarum*, UFPE 3139, collected in Galinhos, Rio Grande do Norte, Brazil, June 2000 (© V. L. Silva)

### YELLOW-BILLED TERN *Sternula superciliaris*

Well-documented records in the state are few. We consider that the following photo-documented records pertain to this species: two breeding-plumaged adults in Natal on 16 May 2011 (WA 611256; D. Barrufi); 16 adults (in various plumages) at Macau on 21 June 2012 (WA 672567, 673494, 675557; J. van den Bosch); an immature, with a brown-margined

strong bill and black wing feathers until the sixth primary, at Guamaré on 9 April 2016 (WA 2084773; J. Dantas); a subadult, with a dirty yellow bill and black until the fifth primary, at the same locality on 30 April 2016 (WA 2103818; L. Sena); a breeding-plumaged adult, with a strong bill and three black outermost primaries in Macau on the same date (WA 2104993; J. Dantas); an immature with a streaked crown, dark-tipped strong bill and black until the fifth primary, in Macau on 24 August 2016 (WA 2249902; J. Dantas); and 33, 20 with all-yellow bills, at Areia Branca on 27 February 2017 (WA 2480542; J. C. Rodrigues). Based on this, we assume that adult Yellow-billed Terns visit the state in February–June, with one record of an immature in August.

#### **GULL-BILLED TERN** *Gelochelidon nilotica*

The north coast seasonally harbours a few colonies comprising dozens of breeding pairs, mixed with breeding Grey-hooded Gulls, in or close to salt pans (FSM, JBI & MP). First proven breeding in June–July 1999 at Salina Diamante Branco, Galinhos, where common in March–August 2000 and a breeding adult was trapped on 17 April 2000 (Azevedo *et al.* 2004). Also, 50 in June, 23 in July, three in August, and post-breeding dispersal from this colony between 6 July and 13 August 2012 (FSM & C. Ferreira). Fifty nests found by FSM and JBI at Salina Soledade, Macau, on 14 May 2005, where birds absent December 2004–January 2005, returned in February, max. 224 (including four adults on nests) on 8 April, 41 nests with eggs, nestlings and a fledged juvenile on 10 June (with I. Cortez), and adults carrying food on 20–23 June and 1–3 July. At the same colony, in 2006, display in April, nests in May–July, and juveniles in June–August (JBI). On 21–22 July 2007, 22 and 15 first-years at Soledade, Macau, and Lagoa Lagamar, Carnaubais, respectively (FSM & JBI).

#### **COMMON TERN** *Sterna hirundo*

Dozens of recoveries from the state's coast have been published, but all lack detailed supporting data (Lara-Resende & Leal 1982, Cordeiro *et al.* 1996, Mestre 2007b, Mestre *et al.* 2010). In an unpublished study conducted in 2005–07 by FSM, 16 Common Terns, ringed as non-volant young in New Hampshire, Massachusetts, Connecticut, New York (USA), and the US Virgin Islands (thus assigned to the nominate form) were seen at Maracajá Reef (05°40'S, 35°24'W), Maxaranguape.

#### **ROSEATE TERN** *Sterna dougallii*

Three coastal recoveries have been published, but all lack detailed data (Lara-Resende & Leal 1982, Mestre *et al.* 2010). In an unpublished study conducted in 2005–07 by FSM, 14 Roseate Terns, ringed as non-volant young in Massachusetts and New York (USA), a nestling ringed BBL in the US Virgin Islands, and another in the Azores (all assigned to the nominate) were seen at Maracajá Reef (05°40'S, 35°24'W), Maxaranguape.

#### **CABOT'S TERN** *Thalasseus acuflavidus*

Yellow-billed *T. a. eurygnathus* is a seasonal breeder on the north coast, in a mixed colony with Grey-hooded Gull, *Gelochelidon* and *Himantopus*. First proven breeding on 19 July 2009 at Furado do Avelino (Salinor), Macau, with a juvenile and five non-volant young (FSM, MP & C. Ferreira; WA 236785; B. França). Around Salina Soledade and Ponta do Tubarão, Macau, birds were absent in December 2004–February 2005, arrived in March, max. 56 and 78 on 20–23 May and 2 July, respectively (FSM & JBI), and display was observed in March 2006 (JBI). On the same coast, 162 (mainly adults) were at the Conchas estuary, Porto do Mangue, on 14 February 2009 (FSM & MP). In addition, one ringed in Mississippi in

December 1964, thus probably of the nominate (Burger 2018), was recovered on the state's north coast in either August 1966 (Sick 1997: 337) or May 1966 (Olmos 2002).

#### **EARED DOVE** *Zenaida auriculata*

The Caatinga population is *Z. a. noronha* (Azevedo & Antas 1990). It is a nomadic mass-breeder tracking the production of seeds following the rainy season, and breeds on the ground in large colonies, where thousands are illegally hunted (J. B. de Medeiros pers. comm.) and sold (FSM pers. obs.). Three banded in Piauí and Pernambuco were recovered in Rio Grande do Norte: one, a youngster, banded (CEMAVE L16622) in Itainópolis, Piauí, on 30 March 1984, was recovered at São José do Seridó in July 1985 (Azevedo & Antas 1990).

#### **GREAT HORNED OWL** *Bubo virginianus*

Known in the state since at least the 19th century, and its onomatopoeic local name, 'jucurutu', is replicated in a municipality in Rio Grande do Norte (Cascudo 1968). Rare in the Brazilian north-east, with state records—based mainly on interviews with local people—concentrated in the south and south-west (Lima *et al.* 2018). It is included in the ESEC Seridó bird list, Serra Negra do Norte (Pichorim *et al.* 2016b), based on an undocumented record of one that flew low above FSM in a steppe-type landscape (06°35'S, 37°15'W) at dusk on 3 December 2002. Birds in the Caatinga have been considered to represent a separate subspecies (*Bubo v. deserti* Reiser, 1905), whose validity is questionable (Traylor 1958, Lima *et al.* 2018).

#### **SHORT-EARED OWL** *Asio flammeus*

The only state record was at Santa Izabel Farm (05°07'S, 36°10'W), Galinhos, on 14 August 2012, where one was flushed from bare sand dunes and was mobbed by up to seven Southern Lapwings *Vanellus chilensis* (Fig. 12; WA 3065349; C. Ferreira & FSM). Except for a handful of localities in Bahia and Piauí states, it seems there are no other records in the Brazilian north-east (König *et al.* 2009, Olsen *et al.* 2019, WikiAves 2019), thus that in Rio Grande do Norte represents a range extension of c.1,000 km north-east.



Figure 12. Short-eared Owl *Asio flammeus*, Galinhos, Rio Grande do Norte, Brazil, August 2012 (© C. Ferreira)

#### **SHORT-TAILED NIGHTHAWK** *Lurocalis semitorquatus*

Just one state record, an immature perched on a high horizontal branch of a *Tabebuia aurea* (Bignoniaceae), in Cerro Corá on 9 July 2017 by D. Oliveira *et al.* (WA 2615000; J. Dantas).

#### **BAND-WINGED NIGHTJAR** *Hydropsalis longirostris*

One documented state record of this rupicolous nightjar: a male at Cânion dos Apertados (06°20'S, 36°30'W), Currais Novos, photographed (WA 3219414) and sound-recorded (WA 3220109) on 21 December 2018 (E. Oliveira).

#### **NACUNDA NIGHTHAWK** *Podager nacunda*

Seldom recorded in extreme north-east Brazil. A female (COUFRN 1228) found dead at Natal International Airport, São Gonçalo do Amarante, on 26 July 2016, was the first documented state record. One responded to playback at ESEC Seridó, Serra Negra do Norte,

on 23 January 2013 and another was heard there on 25 May 2013 (MP), but neither record was documented. A group of 20 settled on a grassy area at Touros on 22 May 2018 (M. Silva; WA 2985934). A partial migrant in Brazil: the nominate subspecies breeds in far southern latitudes and migrates north in the austral winter, when it mixes with resident *P. n. minor*, which is not known to breed in north-east Brazil (Somenzari *et al.* 2018). Both documented state records were during the austral winter, and, together with the first published record for neighbouring Paraíba—a flock of 15 photographed on 23 June 2010 (Pereira *et al.* 2012)—induce us to believe that these were the nominate form. Should *Podager* (unexpectedly) be proven to occur during the local summer (and rainy season in the Caatinga), this would suggest that *P. n. minor* breeds in north-east Brazil.

#### LESSER NIGHTHAWK *Chordeiles acutipennis*

Of the two species of *Chordeiles* of problematic identification likely to occur, only *C. acutipennis* has been claimed: recorded but not described in São José do Seridó in July–December 1996 (Medeiros *et al.* 2000), which justified the species' inclusion in a list of birds of the Seridó region (Varela-Freire & Araújo 1997); one trapped but not described at ESEC Seridó, Serra Negra do Norte (Nascimento 2000, Pacheco 2004); seen but again not described at the same location (MMA 2004); recorded at ESEC Seridó and in Macau based on a reference in Silva *et al.* (2012), the latter corresponding to repeated observations in March–July 2005 (FSM & JBI), including a photo (WA 3089210; JBI) of an adult male in flight within a loose group of 27 birds, on 21 June 2005; and ringed but not described, and mentioned as occasional at ESEC Seridó during field work on 14 July 2012, 26 January 2013 and 1 June 2013 (C. C. O. Silva *et al.* 2017). Also at ESEC Seridó, two alleged *C. acutipennis* were captured on 21 February 2013, both with total length of 20 cm, and mass 33.3 g and 31.9 g, and two others on 9 and 12 June 2013, mass 35.0 g and 34.9 g, respectively (Hasui *et al.* 2018), but without documentation (MP pers. obs.). These weights might apply to Least Nighthawk *Nannochordeiles pusillus* (FSM pers. obs.). A photo taken in the Caatinga at Parelhas on 12 March 2006 is not identifiable to species (WA 344901). For now, we recognise *C. acutipennis* as occurring in Rio Grande do Norte based solely on WA 3089210, whereas *C. minor* is relegated to our secondary list.

#### BISCUTATE SWIFT *Streptoprocne biscutata*

The only holotype of an avian taxon collected in Rio Grande do Norte is the male specimen MNRJ 36897 from Bico da Arara (06°31'S, 36°38'W; 600 m), Acari, of *S. b. seridoensis*. Two Biscutate Swifts, ringed there during 26–27 August 1986, were recaptured on 4–5 August 1987 (Sick *et al.* 1988). The population of this non-breeding colony, first studied by Sick and others, appears to be stable or slightly diminishing, at c.90,000–110,000 birds over the last three decades (Andrade & Freitas 1987, Sick *et al.* 1988, Sick 1991; unpubl. work by FSM in 2005, 2007, 2009, 2011, 2016 and 2018).

#### HORNED SUNGEM *Heliactin bilophus*

Status uncertain, despite repeated records. One at a 'tabuleiro' / floodplain ecotone at Sagi, Baía Formosa, on 3 January 2003. A male and female fed at *Hohenbergia* (Bromeliaceae) in *restinga* at Morro Vermelho, Nísia Floresta, on 16 November 2004, and a breeding-plumaged male was between Lago Azul and Morro Vermelho on 17 April 2005. In the same municipality, an adult male fed on ornamental orange and red Asteraceae flowers at Pium on 7 January 2006 (FSM). Single males were observed in *cerrado* at Punaú (05°23'S, 35°30'W), Rio do Fogo, on 21 October and 12 November 2006, and 17 January 2007, during a study between October 2006 and November 2013 (Pichorim *et al.* 2014a). Moreover, three birds



were seen, of which a female was photographed, at Espírito Santo on 3 September 2017 (WA 2682924; J. Dantas) in a 'tabuleiro', which had burned the year before. At the same place, there are several documented records, including breeding-plumaged males and a female, in May, July and September–October. To date, *Heliactin* has not been reported during the period 18 January–16 April, and breeding in the state, although probable, is unproven.

#### **AMAZONIAN MOTMOT** *Momotus momota*

Since 2002, there have been repeated records at Mata da Estrela Private Reserve (06°23'S, 35°01'W), Baía Formosa (FSM & JFP). Of these, the first to be documented was a singing adult on 28 March 2017 (WA 2510629; P. Fernandes). The Endangered subspecies *M. m. marcgravianus*, endemic to the Pernambuco Centre of Endemism, is not accepted by some authorities, but included by others (Piacentini *et al.* 2015, ICMBio / MMA 2018, Snow & Kirwan 2019).

#### **COLLARED FOREST FALCON** *Micrastur semitorquatus*

Observed in shrubby *restinga* at Barra do Cunhaú (06°18'S, 35°02'W), Tibau do Sul, on 25 December 2002; carrying avian prey at a forest border at Baía Formosa on 28 December 2002; flying over mangroves at Canguaretama on 5 January 2003; and heard at Serra das Queimadas, Parelhas, on 11 June 2006 (FSM). One was sound-recorded at Mata da Estrela Private Reserve (06°22'S, 35°00'W; 25 m), Baía Formosa, on 30 August 2008 (M. Silva), and the first publicly available documentation is a sound-recording from the same locality on 9 September 2018 (WA 3390273; J. A. Marins).

#### **PEREGRINE FALCON** *Falco peregrinus*

Both subspecies that occur in Brazil have been recovered in the state (Silva e Silva 1996, Mestre 2007a): a young *F. p. tundrius* (BBL 987-39774) banded in Alaska, USA (65°N, 141°03'W) on 12 July 1980 and shot in Assu on 15 February 1982, c.12,370 km distant; and a female nestling *F. p. anatum* (BBL 987-15200) banded in Virginia, USA (37°83'N, 75°17'W) on 9 October 1979 and shot in Natal on 18 March 1981, 6,365 km from its nest.

#### **RED-SHOULDERED MACAW** *Diopsittaca nobilis*

Discovered in the south-east of the state: a group of eight at an Atlantic Forest / *cerrado*-like ecotone, in Mata do Pilão, Espírito Santo, on 27 May 2018 (J. Dantas), with a sound-recording from the same location on 10 June 2018 (WA 3233566; M. Silva). A pair was photographed in Natal on 24 May 2011 (WA 359163; S. Messias), but we consider these birds to be of cagebird origin.

#### **PEACH-FRONTED PARAKEET** *Eupsittula aurea*

A genetic study is needed to clarify the origin(s) of the small populations of this species in the state. As suggested by the collecting of five at Mamanguape, Paraíba, in 1957 (Pinto & Camargo 1961), it has probably been present, perhaps continuously, in Rio Grande do Norte for some decades, possibly originating from and maintained by released or escaped cagebirds, in places such as Natal. The Tibau do Sul breeding population could be feral (DMH). *E. aurea* is scarce in *cerrado* at Rio do Fogo (05°23'S, 35°30'W) (Pichorim *et al.* 2014a), arguably the best habitat for the species in the state.

#### **WHITE-FLANKED ANTWREN** *Myrmotherula axillaris*

The subspecies in the state is *M. a. luctuosa*, which is sometimes treated at species level (del Hoyo *et al.* 2019b). All records are from Baía Formosa, at Mata da Estrela Private Reserve

(06°23'S, 35°01'W), where at least 32 individuals were found during 27 December 2010–3 January 2011 (JFP), and Mata da Pituba (06°27'S, 35°03'W) (RDL).

#### **RUFOUS-WINGED ANTWREN** *Herpsilochmus rufimarginatus*

The population in the Pernambuco Centre of Endemism is generally attributed to *H. r. scapularis* (Pinto 1978, del Hoyo *et al.* 2019a), but a taxonomic review of this complex suggests that this population may be a novel taxon, whose plumage and voice is very similar to Amazonian *H. r. frater* (Silva 2013; JFP pers. obs.).

#### **WILLIS'S ANTBIRD** *Cercomacroides laeta*

The only state record was a pair seen and sound-recorded in the valley of the Pituba River, Baía Formosa, on 8 September 2009 (WA 2660533; MP, M. Silva & JBI). The Vulnerable subspecies *C. l. sabinoi* (MMA 2003) is not always accepted as valid (Bierregaard *et al.* 1997, Cavarzere 2014) and has not been evaluated by ICMBio / MMA (2018).

#### **BLACK-CHEEKED GNATEATER** *Conopophaga melanops*

The Vulnerable subspecies *C. m. nigrifrons* is endemic to the Pernambuco Centre of Endemism (ICMBio / MMA 2018). It occurs around Baía Formosa at two localities: first documented at Mata da Estrela Private Reserve (06°23'S, 35°01'W) via a sound-recording on 14 August 2016 (WA 2235397; J. Dantas); and sight records at Mata da Pituba (JBI).

#### **PLAIN XENOPS** *Xenops minutus*

The subspecies in Rio Grande do Norte is *X. m. alagoanus*, which is considered Vulnerable (ICMBio / MMA 2018) and whose voice is more similar to Amazonian *X. m. genibarbis* than to the nominate Atlantic Forest population (RDL pers. obs.). Recent molecular data showed that one collected in Pernambuco (FMNH 399212; Field Museum of Natural History, Chicago) grouped phylogenetically with birds in eastern Amazonia, and not with the Atlantic Forest clade (Burney 2009), but this author inferred that the species is paraphyletic, when in fact *alagoanus* could be ranked as a species. Widely overlooked since its description (del Hoyo *et al.* 2019c), *alagoanus* was omitted from a recent phylogeny of the species (Harvey & Brumfield 2015).

#### **RUFOUS-FRONTED THORNBIRD** *Phacellodomus rufifrons*

Reported to be expanding its range in south-east Brazil (Willis & Oniki 1993; JFP pers. obs.), this thornbird has recently reached western Rio Grande do Norte, where it was found at Serra do Mel in 2007 (G. Pereira), and Umarizal and Luís Gomes, respectively, in May and June 2010 (FSM), and the south-east, at Santo Antônio and Nova Cruz in January 2011 (JFP). It was first documented by a photo of two birds at Passa e Fica on 27 September 2015 (WA 1851438; D. Gurgel).

#### **RED-SHOULDERED SPINETAIL** *Synallaxis hellmayri*

A Brazilian endemic treated either as Data Deficient (ICMBio / MMA 2018) or Near Threatened (IUCN 2019). The first state record was one trapped at ESEC Seridó, Serra Negra do Norte, on 17 December 1995 (Nascimento 2000). A vocalising pair sound-recorded at 650 m in the Serra das Queimadas, Parelhas, on 14 November 2005 (FSM) is the second. The state's main population is in the highest parts of the Serra de Santana (FSM pers. obs.; Pichorim *et al.* 2016a). Other records are concentrated in the Seridó region, with a few on the north coast. Very local in distribution, *S. hellmayri* is probably better considered Near Threatened at state level.

**BEARDED BELLBIRD** *Procnias averano*

Was considered extinct in the state (Paiva 1983 *apud* Sick 1972, Teixeira 1992). Rediscovered at Baía Formosa on 13 June 2015 by P. Vitor, F. V. Souza, and J. Dantas, who photographed a female feeding on *Psidium* fruits (Myrtaceae) (WA 1724942), which is the only documented record.

**WHITE-THROATED SPADEBILL** *Platyrinchus mystaceus*

First record at Mata da Estrela Private Reserve, Baía Formosa, on 28 December 2002 (FSM); also known from Mata do Mucambo, Goianinha (M. Silva, FSM & JFP pers. obs.), and sound-recorded at Canguaretama (WA 2598721; F. V. Souza). The Vulnerable and declining *P. m. niveigularis* is endemic to the Pernambuco Centre of Endemism, and in need of taxonomic study (ICMBio / MMA 2018).

**WHITE-BELLIED TODY-TYRANT** *Hemitriccus griseipectus*

Uncommon in remnant patches of Atlantic Forest between Baía Formosa and Natal (Olmos 2003; FSM, JFP & RDL). The Vulnerable *H. g. naumburgae* is endemic to the Pernambuco Centre of Endemism (ICMBio / MMA 2018).

**WHITE-LORED TYRANNULET** *Ornithion inerme*

A rapid song of this tyrannulet was taped at a forest border in the valley of the Catu River (06°20'S, 35°09'W), Canguaretama, on 13 December 2005 (WA 3300921; FSM, identified by RDL & JFP). Also heard singing repeatedly in riverine forest at Mucambo (06°19'S, 35°14'W), Goianinha, on 2 January 2011 (JFP). More recently, an adult and a just-fledged juvenile at a forest border in the same locality (WA 3257427, 3300921; RDL) provided proof of breeding (RDL & FSM).

**PLAIN-CRESTED ELAENIA** *Elaenia cristata*

One of the few species documented by a historical specimen, a male (LSUMZ 10420) taken by J. H. Phares in Natal on 9 September 1945. The demography and life history of this elaenia have been studied in *restinga* woodland on dunes with 'tabuleiro' in Parnamirim (Toledo-Lima 2013). Its citation for the north coast Caatinga, in Macau (Silva *et al.* 2012), is the only record for that biome in the state and is perhaps in error (FSM & JBI).

**SWAINSON'S FLYCATCHER** *Myiarchus swainsoni*

Vocalisations exclusively attributable to this species establish its presence in the state, but its status is uncertain. Visual diagnosis is less reliable (Fig. 13). First documented via a sound-recording of its repeated call at Fazenda Malhada Vermelha (06°45'S, 36°40'W; 400 m), Parelhas, on 14 November 2005 (WA 3300894; FSM, identified by RDL), which may suggest local breeding or a late austral spring migrant. Two individuals reported as this species, with brood patches, were caught during the dry season, on 3 May and 25–26 July 1996 at ESEC Seridó, Serra Negra do Norte (Nascimento 2000). At the same locality, mist-netting work suggested the species to be a migrant (Damasceno 2015), nomadic / migratory (C. C. O. Silva *et al.* 2017) or a seasonal visitor (G. Toledo-Lima). Apart from the November documentation, records suggest the species is present in Rio Grande do Norte in March–August (RDL), i.e. the austral winter. *M. swainsoni* is a partial migrant, the nominate subspecies being the southernmost taxon, which migrates to northern South America, while sedentary populations in central Brazil may be *M. s. pelzelni* (Somenzari *et al.* 2018). Based on analysis of specimens, both subspecies have been found in nearby Ceará (Lanyon 1978).



**SHORT-CRESTED FLYCATCHER** *Myiarchus ferox*

The typical call of this *Myiarchus* was first documented at Espírito Santo on 27 May 2018 (WA 2984460; J. Dantas). *M. ferox* had been observed previously: one calling in shrubby *restinga* at Barra do Cunhaú, Tibau do Sul, on 26 December 2002; two calling in riverine forest at Mata do Mucambo, Goianinha, on 20 May 2006 (FSM); two heard at an Atlantic Forest / sugarcane plantation ecotone, Baía Formosa, on 28 December 2010; one heard at Mucambo on 30 December 2010; and one heard at Guaju mangroves, Baía Formosa, on 1 January 2011 (JFP). *M. ferox* was mentioned for the Caatinga by Silva *et al.* (2012), but specimen COUFRN 40 from this habitat in Macaíba was re-identified as *M. tyrannulus* by RDL. Although claimed to have been ringed and reportedly resident at ESEC Seridó, Serra Negra do Norte (C. C. O. Silva *et al.* 2017), we consider its presence there improbable (Pichorim *et al.* 2016b). We regard the species to be the only *Myiarchus* restricted to the Atlantic Forest in the state.

**BROWN-CRESTED FLYCATCHER** *Myiarchus tyrannulus*

One of the commonest passerines in Rio Grande do Norte, especially in almost all Caatinga areas. Less common in the coastal strip, *M. tyrannulus* was seen in mangroves at Canguaretama on 24 December 2002 (FSM), and was heard in a small patch of Atlantic Forest at Baía Formosa on 28 December 2010 (JFP & FSM). During the rainy season, *M. tyrannulus* occurs in sympatry with *M. swainsoni*. For visual identification of these two species, see Fig. 13.



Figure 13A and C. Brown-crested Flycatcher *Myiarchus tyrannulus*, Tibau, Rio Grande do Norte, Brazil, February 2018; (B and D) Swainson's Flycatcher *M. swainsoni*, Baraúna, Rio Grande do Norte, Brazil, April 2018; note the heavy bill of *M. tyrannulus*, proportionately smaller in *M. swainsoni*, in addition to its pinkish base, and the rufous-brown in the tail feathers of *M. tyrannulus*, which is absent in *M. swainsoni* (Rafael Dantas Lima)

**VELVETY BLACK TYRANT** *Knipolegus nigerrimus*

Varela-Freire & Araújo (1997) listed Blue-billed Black Tyrant *K. cyanirostris* in the dry Seridó region, but we consider this to be a misidentification of *K. nigerrimus* (see tertiary list in Appendix 1), a species collected in the state (MNRJ 2078) years ago. At Fazenda Salobro

Private Reserve (06°13'S, 37°03'W; 400 m), Jucurutu, a female was seen and two males were trapped on 31 December 2008 (FSM). *K. nigerrimus* has been documented at several other Caatinga sites in the Seridó region, e.g. Parelhas, where adults tended a nestling on 26 April 2016 (WA 2098441; L. Sena), Acari (06°31'S, 36°38'W; 600 m) and Currais Novos (WA 3099923, 3183124; T. Elis). All of the records of females involved birds with large coloured throats (e.g., WA 3194677; T. Elis; FSM), which lead us to conclude that the nominate occurs. If valid, *K. n. hoflingae*, described from Bahia based on a single female with a thinner striated throat patch, the chin and throat darker than the nominate, and a dusky wing panel, differs from females in Rio Grande do Norte. As it has been incautiously inferred that *hoflingae* is distributed throughout far north-east Brazil, including Rio Grande do Norte (Farnsworth *et al.* 2019), we offer here measurements of two adult males trapped in Jucurutu as a basis for comparative study of these taxa: both total length 159 mm; wing chord 91.5 and 92.5 mm (wingtip p4); tail 80.5 and 79.0 mm; tarsus both 21.8 mm; exposed culmen 13.7 and 14.0 mm; bill to nostrils 9.5 and 10.0 mm; gonys 3.8 and 4.0 mm; mass 21.0 and 19.75 g (CEMAVE E79861, E79864).

### SPECTACLED TYRANT *Hymenops perspicillatus*

The only state record of this austral migrant constitutes the northernmost record by c.1,400 km. An adult male was photographed foraging on the ground below a group of *Prosopis uniflora* trees (Mimosaceae) at the ecotone between dunes and mangroves at Santa Isabel Farm (05°07'S, 36°10'W), Galinhos, on 10 December 2012 (Fig. 14; WA 3065355; C. Ferreira), and was observed again by FSM two days later, foraging under the same trees. Since then, there has been a second record in north-east Brazil, on 17 May 2014, a female present for a month at Caravelas, Santa Bárbara Island, in southernmost Bahia, following strong southerly winds on 26–28 April (WA 1335827; A. Carletti). This species moves north during the austral winter, with records in the states of Paraná, São Paulo, Rio de Janeiro, Mato Grosso do Sul and Mato Grosso in April–November (Somenzari *et al.* 2018).



Figure 14. Male Spectacled Tyrant *Hymenops perspicillatus*, Galinhos, Rio Grande do Norte, Brazil, December 2012 (© C. Ferreira)

### YELLOW-BROWED TYRANT *Satrapa icterophrys*

The five available state records are all from the period May 2012–June 2013, at three Caatinga localities: Alto do Rodrigues and Macau, on 27 May 2012 and 10 June 2012 (WA 649286, 663484; J. van den Bosch), and ESEC Seridó, Serra Negra do Norte, where it was trapped not only in the austral winter (on 14 July 2012 and 1 June 2013), but also on 26 January 2013, when the region was still dry (C. C. O. Silva *et al.* 2017). The true status of this species in Rio Grande do Norte is unclear.

**GREY MONJITA** *Xolmis cinereus*

The first documented state record, extending the known range c.650 km north-east, is from Punaú, Rio do Fogo, where two were observed on 29 October 2006, and singles on 21 December 2006, 16 August 2008 and 16 November 2013, in recently burned areas (Pichorim *et al.* 2014a; WA 1689513; B. França). At the same location, one was photographed on 15 and 29 January 2017 (WA 2435098; D. Gurgel, WA 2452371; A. Rafael). Lastly, just to the west, at Pureza, an adult was photographed on 29 June 2018 (WA 3030610; M. Silva). The species frequents this *cerrado* enclave in the austral winter (June and August) and during the austral spring migration (October–December) but also in January, which might suggest local breeding. Its status in Rio Grande do Norte is still to be clarified.

**BLUE-AND-WHITE SWALLOW** *Pygochelidon cyanoleuca*

First documented in Ceará-Mirim on 17 May 2007 (WA 3459903; JBI), this swallow is uncommon in the state, where we know of just 11 records, each involving 1–3 individuals, in February–September. Ten of these, between 30 March and 9 September, during the austral autumn and winter, potentially involve southern South American *P. c. patagonica*, but the record in February may well represent the nominate form.

**BROWN-CHESTED MARTIN** *Progne tapera*

Two subspecies occur: the nominate, represented by LSUMZ 10421, taken by J. H. Phares in Natal on 9 August 1945; and *P. t. fusca*, an austral migrant (Somenzari *et al.* 2018), photographed in São Miguel do Gostoso on 18 July 2017 (WA 2634379; E. Guilherme). Groups numbering dozens to >300 individuals, including the latter form, have been recorded by JBI, mainly at the landfill of Ceará-Mirim, almost annually since 2005, between March and September (e.g., WA 3459758).

**WHITE-RUMPED SWALLOW** *Tachycineta leucorrhoa*

Two with one Blue-and-white Swallow, and a group of Southern Rough-winged Swallows *Stelgidopteryx ruficollis*, at the landfill in Ceará-Mirim on 17 May 2007, and one with a group of *Stelgidopteryx* at the same location on 27 September 2007 (Fig. 15A) (JBI), are the first photo-documented records for the state. These correspond to the wintering and pre-breeding periods for this austral migrant, which reaches the northernmost limit of its range in Rio Grande do Norte (Moreira-Lima 2013, WikiAves 2019). *T. leucorrhoa* was reportedly




Figure 15A. White-rumped Swallow *Tachycineta leucorrhoa*, Ceará-Mirim, Rio Grande do Norte, Brazil, September 2007 (Jorge Bañuelos Irusta); (B) *T. leucorrhoa*, Alto do Rodrigues, Rio Grande do Norte, Brazil, May 2012 (© J. van den Bosch)



trapped at Salina Diamante Branco, Galinhos, in April 2003 (Araújo *et al.* 2004), but was said to be resident, rather than a migrant, thereby casting doubt on the identification. The first available proof of occasional overwintering by this species is a photo of an adult at Alto do Rodrigues on 23 May 2012 (Fig. 15B; WA 646156; J. van den Bosch). At least four were photographed together, again in Ceará-Mirim, on 26 June 2013 (JBI).

#### CHILEAN SWALLOW *Tachycineta leucopyga*

This austral migrant was photographed by JBI at Ceará-Mirim on 30 May 2011 (Fig. 16), the sole documented record for the state. *T. leucopyga* had been claimed at Macau on 14 September 2008 (Silva *et al.* 2012), which would have been a new species for the Caatinga biome, but the photo was ultimately re-identified as a White-winged Swallow *T. albiventer* (WA 761685). The same authors mentioned a record at Ceará-Mirim, also in September 2008, which they considered to be the first for north-east Brazil, but the record was not documented. They also referred to Pacheco *et al.* (1996)—for a record of *T. leucopyga* in Rio de Janeiro—as its northern limit in the austral winter, but the species concerned was *T. leucorrhoa*. Nevertheless, the formal Brazilian limit for overwintering *T. leucopyga* was considered to be Rio de Janeiro (Somenzari *et al.* 2018, WikiAves 2019), consequently the documented record in Rio Grande do Norte is the northernmost by c.1,900 km.



Figure 16. Chilean Swallow *Tachycineta leucopyga* in Ceará-Mirim, Rio Grande do Norte, Brazil, May 2011 (Jorge Bañuelos Irusta)

#### BLACK-CAPPED DONACOBIUS *Donacobius atricapilla*

First noted in 2001 at Povoado de Sítio, Ceará-Mirim, and repeatedly observed there, where it was collected by M. Silva in 2007 (COUFRN 14). At least seven, one of which sang and a pair that duetted, in the same floodplain on 7–8 July 2007 (FSM & M. Silva). The photo of an adult male singing on 8 April 2010 was taken at this sole locality in the state (WA 279016; B. França).

#### LONG-BILLED GNATWREN *Ramphocaenus melanurus*

The nominate subspecies is an Atlantic Forest endemic (Moreira-Lima 2013). In Rio Grande do Norte, the species was first observed in mixed-species flocks with White-flanked Antwren *Myrmotherula axillaris*, White-fringed Antwren *Formicivora grisea*, Plain Antvireo *Dysithamnus mentalis*, Pectoral Antwren *Herpsilochmus pectoralis*, Planalto Slaty Antshrike *Thamnophilus pelzelni*, Black-cheeked Gnatcatcher *Conopophaga melanops* and Plain Xenops *Xenops minutus* at Mata da Estrela Private Reserve, Baía Formosa, on 14–15 December 2002 (FSM). *Ramphocaenus* was first documented by a sound-recording at this sole state locality on 12 March 2016 (WA 2270254; D. Gurgel).



Figure 17. Female Yellow-legged Thrushes *Turdus flavipes*, COUFRN 812 and COUFRN 844, collected in Barreira do Inferno, Parnamirim, Rio Grande do Norte, Brazil, May and July 2013, respectively (Rafael Dantas Lima)

#### YELLOW-LEGGED THRUSH *Turdus flavipes*

Not previously suspected to occur in the state, this thrush has been documented four times in the 21st century, during the austral winter: an adult male without a whitish chin (i.e. of the nominate subspecies) ringed by G. Segurado and photographed by B. França at Macaíba on 14 June 2008 (Silva *et al.* 2012), at the ecotone of Caatinga / Atlantic Forest; several of both sexes trapped, of which two females were collected in *restinga* during a study in 2012–16 (Fig. 17); and an adult female photographed at Natal on 12 April 2015 (WA 1662001; D. Gurgel). The nominate race is endemic to the Atlantic Forest (Moreira-Lima 2013), and its complex migratory status, including seasonal altitudinal movements, remain to be clarified (Somenzari *et al.* 2018).

#### CREAMY-BELLIED THRUSH *Turdus amaurochalinus*

First ringed in the state at Vila Industrial Alcanor, Macau, in May 1988 (D. Souza) at the end of the rainy season in the Caatinga, i.e. the early austral winter. A fresh-plumaged youngster with pointed tail feathers, some evidence of moult on its breast, thighs and vent, mass 60.25 g, and a very young individual (mass 61.5 g) were trapped at Serra das Queimadas, Parelhas, on 9–10 June 2006, suggesting local breeding (Fig. 18). One full-plumaged adult (mass 63 g) with a brood patch, and another full-plumaged adult (55 g) were caught on 10 and 11 March 2007 at the same location, at the start of the austral winter. Also, a heavily speckled juvenile (55 g) was trapped on 8 May 2009, and an adult female (53.5 g) with a brood patch, ringed on 9 May 2009, was recaptured with a well-developed brood patch, mass 57 g, on 19 January 2010, just over eight months later, all at Pimenteira in the Serra de Santana, Santana do Matos



Figure 18. Young Creamy-bellied Thrush *Turdus amaurochalinus*, Serra das Queimadas, Parelhas, Rio Grande do Norte, Brazil, June 2006 (© L. Sena)

(FSM). In lowland Caatinga, singing birds and adults in nuptial plumage were recorded in January–April (FSM), and five with brood patches were captured in Mossoró (Cavalcanti *et al.* 2016). In Rio Grande do Norte, this thrush forms local concentrations in the austral winter, e.g., c.100 birds in 4 ha of low *restinga* at Maxaranguape on 4 August 2015 (FSM); these were presumably either austral migrants or a seasonal gathering of residents at a rich food source. This species occurs year-round in north-east Brazil (Somenzari *et al.* 2018). The latter authors considered that records in this region do not suggest migratory behaviour, probably due to the co-existence of resident / migratory populations—which we believe to be true—but we note that, at least in Rio Grande do Norte, nesting is better established than the arrival of austral migrants.

### CHALK-BROWED MOCKINGBIRD

*Mimus saturninus*

The breeding biology of the species was studied in *restinga* on the campus of the Federal University in Natal during 2006–07 (unpubl. study by L. M. C. Dantas *et al.*). It is also a common breeder in the Caatinga, illustrated by a nest at Fazenda Salobro Private Reserve (06°13'S, 37°02'W; 400 m), Jucurutu, on 23 March 2017 (Fig. 19). Populations in north-east Brazil have traditionally been treated as *M. s. arenaceus*, but this subspecies is diagnosed only by its longer bill (Hellmayr 1934). A modern taxonomic analysis would be welcome.



Figure 19. Nestlings and egg of Chalk-browed Mockingbird *Mimus saturninus*, Jucurutu, Rio Grande do Norte, Brazil, March 2017 (© J. Sagot)

### YELLOW-RUMPED CACIQUE *Cacicus cela*

This cacique breeds around Baía Formosa, as reported by local people to FSM and revealed by photos (WikiAves 2019). It is possibly a feral population, still being reinforced by released cagebirds. A former colony in Ceará-Mirim was persecuted to extinction (M. Silva) and another colony at Nísia Floresta in the 1980s also no longer exists (DMH).

### SCREAMING COWBIRD *Molothrus rufoaxillaris*

The first mention in the state is from August 2010 (Pichorim *et al.* 2016a). Documentation of immatures in transitional plumages (WA 2164752; J. van den Bosch, WA 2641146; S. Furtado) establishes the species on the primary list. Reported to have recently invaded eastern Brazil (ICMBio / MMA 2018), and that is perhaps true for Rio Grande do Norte too but, based on WikiAves records (to 31 March 2019), it occurs only in March–September, suggesting the species is an austral winter visitor.

### CINNAMON TANAGER *Schistochlamys ruficapillus*

Two specimens from the state: UMMZ 113609, taken by E. W. van Aken in Natal on 19 February 1945; and an undated female (COUFRN 705) from the *restinga* at Parnamirim (MP). The species has been recorded almost throughout the east coast, including: an adult in shrubby *restinga* at Barra do Cunhaú (06°18'S, 35°02'W), Tibau do Sul, on 26 December 2002; an adult in low-stature Atlantic Forest at Dunas de Natal State Park, Natal, on 24 November 2004 (FSM); up to five in *cerrado* at Punaú (05°23'S, 35°30'W), Rio do Fogo (Pichorim *et al.* 2014a, where a photo was published); and in 'tabuleiro' at Jundiá (05°53'S, 35°23'W),



Macaíba (Silva *et al.* 2012). A sighting in the Caatinga at Macau (Silva *et al.* 2012) is the only record for this biome in the state, but escaped / released cagebirds may be the origin (FSM pers. obs.).

#### **SAFFRON FINCH** *Sicalis flaveola*

Interviews with local people by FSM confirmed this finch was formerly numerous in the state (Faria 2001). It is still very sought after by the cagebird trade, filling many of the environmental agency's rehabilitation centres throughout north-east Brazil. It is regularly released by the environmental police without identifying the subspecies concerned, probably leading to some genetic mixing between *S. f. brasiliensis*, the native local taxon, and *S. f. pelzelni*.

#### **BRAZILIAN TANAGER** *Ramphocelus bresilius*

Individuals frequent the *restinga* of Parnamirim and Natal (P. Vitor), but breeding is unknown (MP), and environmental agents regularly release cagebirds (FSM). Birds recorded in *restinga* and in mangroves at Baía Formosa and Atlantic Forest borders at Goianinha probably represent remnant native populations (JFP & FSM).

#### **WHITE-BELLIED SEEDEATER** *Sporophila leucoptera*

A pair was photographed in the floodplain of the Pirangi River, at Cotovelo, Parnamirim, on 1 May 2008 (JBI), at the start of the local rainy season. A male *S. l. cinereola* was photographed at Nísia Floresta on 28 August 2017 (WA 2676962; F. E. Silva, identified by FSM), at the end of the local rainy season. The so-called 'chorão' or 'patativa' is much sought after by traffickers in the state (Pichorim *et al.* 2016b, FSM pers. obs.).

#### **BLACK-THROATED SALTATOR** *Saltatricula atricollis*

Recently proven to occur in the north-west of the state: a photo of a moulting male singing at Assu on 26 June 2017 (WA 2658215; M. Silva), and a photo and sound-recording of a singing adult at Sítio do Góis, Apodi, on 11 March 2018 (WA 2908958, 2909323; J. Souza).

#### **COMMON WAXBILL** *Estrilda astrild*

This exotic invasive species, present in Natal since 1985 when it was seen building a nest (C. Medeiros), is currently common along the east coast but scarce in the dry interior of the state (FSM pers. obs.; WikiAves 2019).

#### **HOUSE SPARROW** *Passer domesticus*

Introduced from the Old World, and currently common throughout the state. The nominate is the only subspecies known to occur in Brazil (Piacentini *et al.* 2015).

## **Species on the Secondary List**

#### **RED-WINGED TINAMOU** *Rhynchotus rufescens*

The first published state record (Olmos 2003) lacked documentary evidence, and two other references to the species stemmed from interviews with local people (Silveira & Varela-Freire 1999, Pichorim *et al.* 2016a). It was heard and seen at ESEC Seridó, Serra Negra do Norte (JBI), but was not documented, and this is the sole record to support its inclusion in the Caatinga bird list for the state (Silva *et al.* 2012) along with that for the same locality (Pichorim *et al.* 2016b).



**BLUE-WINGED TEAL** *Anas discors*

Ten ringed in the USA and Canada between 1955 and 1971 were recovered in nine municipalities in the state between 1959 and 1975 (Mestre *et al.* 2010), but there is no documentary evidence to support these records (CEMAVE pers. comm., BBL pers. comm.). Subsequently, the species has not been recorded again in Rio Grande do Norte.

**AMERICAN FLAMINGO** *Phoenicopterus ruber*

A rock painting at 'Pedra Ferrada', São Rafael, has been attributed to this species (Sick 1997: 227). An adult was present at Salina Diamante Branco, Galinhos, over five consecutive days in March 2001 (Azevedo *et al.* 2004), but went undocumented.

**MAGELLANIC PENGUIN** *Spheniscus magellanicus*

Two weakened immatures, one beached on 7 August 2008 in Natal, the other captured at sea off Natal on 13 August 2008, were found during an influx of this penguin along the Brazilian coast due to a severe austral winter (Dantas *et al.* 2013). Both subsequently died, but the frozen specimens (COUFRN 28, 30) were lost due to an electricity outage.

**YELLOW-NOSED ALBATROSS** *Thalassarche chlororhynchos*

An albatross found beached on the north coast of the state, in May 2018, was rehabilitated and released on about 20 December 2018 (F. Lima pers. comm.). Two photos of the bird were used to publicise an educational project, in which it was identified as a Black-browed Albatross *T. melanophris*. In fact, the dark bill with clear yellow-green culmen identify this immature as *T. chlororhynchos* (Bugoni & Furness 2009). The species is maintained on our secondary list because the documentary evidence is not available. This is the only known state record, but one of many in north-east Brazil, e.g. one seen 55 km north-east of the Fernando de Noronha archipelago on 15 November 1996 by A. B. S. Fuchs (J. M. R. Soto, MS); 37 found beached on the north coast of Bahia in 1994–98 (Lima *et al.* 2004); an immature off Salvador, Bahia, on 25 May 2009 (WA 156301; B. Rennó); and an immature 45 km off Paracuru, Ceará, on 5 January 2018 (WA 2842425; C. E. Moura). Endangered *T. chlororhynchos* occurs along most of the Brazilian coast (ICMBio / MMA 2018), and our partial compilation of records confirms that immatures are reasonably regular off the north-east of the country.

**ZINO'S PETREL** *Pterodroma madeira*

This pelagic species, which breeds in Macaronesia, is believed to occur off the state's coast as a result of the use of dataloggers to track the tiny population (Zino *et al.* 2011, Ramos *et al.* 2016), but lacks confirmation through direct observation.

**DESERTAS PETREL** *Pterodroma deserta*

Like the previous species, occurrence off the state's coast is based on information gathered from dataloggers (Ramirez *et al.* 2013, Ramos *et al.* 2016), but lacks confirmation through direct observation.

**BULWER'S PETREL** *Bulweria bulwerii*

The occurrence of this pelagic species off Rio Grande do Norte is also based on information gathered from dataloggers (Ramos *et al.* 2015, Cruz-Flores *et al.* 2019), but lacks direct confirmation.

**BOYD'S SHEARWATER** *Puffinus boydi*

This shearwater, endemic to the Cape Verde Islands, is believed to occur off the state's coast in the non-breeding season (Zajková *et al.* 2017), but this discovery, which is based on information garnered from dataloggers, lacks direct confirmation. *P. boydi* was split from Macaronesian Shearwater *P. baroli*, itself split from Little Shearwater *P. assimilis* (Gill & Donsker 2019), but is treated as a subspecies of Audubon's Shearwater *P. lherminieri* by some authorities (Carboneras *et al.* 2019). Like *Pterodroma madeira*, *P. deserta*, and *Bulweria*, because of the relative accuracy of geolocators (+/- 200 km), it is impossible to be sure if this shearwater occurs in the state's continental waters, which reach only 27–40 km from the coast.

**BROWN BOOBY** *Sula leucogaster*

One, beached south of Natal, was kept by A. Varela-Freire between September 1986 and December 1987 (Varela-Freire 1997) but not documented. A weak individual floating in the sea off Guamaré in early March 2019 was photographed by A. Gleydson, but the photo is unavailable (RDL & FSM).

**CAPPED HERON** *Pilherodius pileatus*

One flushed by FSM from the almost-dry stony bed of Salobro stream, with still a few seasonal ponds and flanked by gallery forest, in Martins on 4 August 2010, was not documented. The species is known from nearby localities in neighbouring Ceará, Paraíba and Pernambuco (Pereira *et al.* 2012, WikiAves 2019). More records are expected in Rio Grande do Norte.

**BUFF-NECKED IBIS** *Theristicus caudatus*

This conspicuous ibis was seen once in the Seridó region around 1996, but not in 'açude Riacho do Roçado' as claimed by Medeiros *et al.* (2000; A. da Silveira pers. comm.). This record supported the species' inclusion in the first state bird list (Varela-Freire *et al.* 1999). *T. caudatus* is now considered extinct on the north coast, based on interviews with local people by FSM that indicated its presence until the 1950s at Macau and Guamaré, where it was regularly hunted. Now rare in north-east Brazil, where absent from Rio Grande do Norte to north-west Ceará and from almost all of the Caatinga (WikiAves 2019); it is no longer expected in the state.

**SWALLOW-TAILED KITE** *Elanoides forficatus*

Considered rare in the first state bird list (Varela-Freire *et al.* 1999), but there is no information concerning the origin of the claimed record. A bird perched on electric wires along the unsurfaced road to Morro Vermelho (05°59'S, 35°10'W), Nísia Floresta, on several consecutive days in 2004, was finally shot and buried there (C. Ferreira pers. comm.). The nearest records to Rio Grande do Norte are from Pernambuco; individuals on 15 October and 4 December 1999, possibly of a resident population (Roda & Pereira 2006), and on 17 October 2009 (Pereira *et al.* 2014). Records in central-north Ceará on 25 November 2007, and Alagoas on 19 October 2018 (WikiAves 2019), suggest migration as well; thus, further state records can be expected.

**SUNGREBE** *Heliornis fulica*

Around 1998, a weak Sungrebe (identified by DMH) was brought to Pipa Ecological Sanctuary, Tibau do Sul, where it died some days later but was not documented. An uncommon resident in the Atlantic Forest (Moreira-Lima 2013); because our record lacks

information as to the locality where it was found, it may not have been in Rio Grande do Norte.

#### **HUDSONIAN GODWIT** *Limosa haemastica*

An immature bird, probably a Hudsonian Godwit, was seen at Salina Soledade, Macau, on 22 June 2005 (Irusta & Sagot-Martin 2011), but not documented. Autumn migration passes inland (west) of north-east Brazil and in spring the species uses few or no stopover sites (Somenzari *et al.* 2018); thus, the species is not expected in Rio Grande do Norte.

#### **EURASIAN WHIMBREL** *Numenius phaeopus*

Seen twice in the state: one in a flock with 14 American Whimbrels *N. hudsonicus*, seen both perched and flight at Canto do Amaro mangroves, Guamaré, on 9 October 2010; and one in a flock with c.160 *N. hudsonicus* flushed from mangrove islets (05°50'S, 35°17'W) in Macaíba on 18 February 2016 (FSM). There is no documentary evidence, but *N. phaeopus* has been repeatedly documented on Fernando de Noronha (Silva e Silva 2008), and more state records can be expected.

#### **BLACK TERN** *Chlidonias niger*

One, banded in 1984 in Berlin, Germany, and therefore of the nominate subspecies, was recovered at Macau in September 1986 (Sick 1997: 335). Although *C. n. surinamensis* is the only subspecies admitted by Piacentini *et al.* (2015), these authors note, based on the Macau record, that the nominate is also expected in Brazil, and *surinamensis* should occur in Rio Grande do Norte.

#### **SCALED PIGEON** *Patagioenas speciosa*

An adult perched atop a 15 m-tall tree was seen in riverine forest along Mucambo stream (06°19'S, 35°14'W), Goianinha, on 21 January 2019 (FSM), but was not documented. Given that it occurs at neighbouring localities in Paraíba, the species was expected in the state.

#### **RUDDY QUAIL-DOVE** *Geotrygon montana*

A probable Ruddy Quail-Dove was heard singing at Mata da Estrela Private Reserve, Baía Formosa, on 29 December 2002, and two others, probably a pair of this species, were flushed at a forest border in Estivas Private Reserve, Canguaretama, on 12 December 2005 (FSM). The species is expected in Rio Grande do Norte, with the nearest documented record in neighbouring Mamanguape, Paraíba, on 12 May 2015 (WA 1694065; E. Barreto).

#### **SPECTACLED OWL** *Pulsatrix perspicillata*

The species has been observed, but not documented, in coastal dunes at Natal (Varela-Freire *et al.* 1999), but we are unable to find any record to support this assertion, and due to the lack of surface water and paucity of tall trees we believe Dunas de Natal State Park does not harbour the species (FSM pers. obs.). Nevertheless, there is a suitable habitat in the Atlantic Forest of south-east Rio Grande do Norte. It is found very nearby at Mamanguape in far north-east Paraíba (Pinto & Camargo 1961, WikiAves 2019), so it can be expected in the state.

#### **MOTTLED OWL** *Strix virgata*

The song of this owl was heard at Casqueira forest (06°24'S, 35°04'W), Baía Formosa, on 26 January 2015 during a full moon (FSM), but was not sound-recorded. The nearest

documented records are from Pernambuco (Roda & Pereira 2006) and to the north, as far as Pará.

#### **BUFF-FRONTED OWL** *Aegolius harrisii*

We know of two trustworthy records in the state: one responding to playback on two nights at ESEC Seridó, Serra Negra do Norte (Pichorim *et al.* 2016b); and one heard at Cerro Corá in July 2017 (D. Oliveira), but there is no documentary evidence.

#### **COMMON NIGHTHAWK** *Chordeiles minor*

All Brazilian records, including those photo-documented or sound-recorded, are in September–April, and *C. m. chapmani* overwinters in central Brazil and northern Argentina (Somenzari *et al.* 2018). Even photo-documented records require careful analysis due to the difficulty in separating *C. minor* from *C. acutipennis*. Two photos of birds perched on branches, at Parelhas on 12 March 2006 (WA 344901; L. Sena) and Macau on 18 March 2015 (WA 1640227; J. Dantas), could pertain to either of these species. *C. minor* appears rather common at Chapada do Araripe, in southern Ceará, where MNRJ obtained an adult female on 15 March 1989 (Teixeira *et al.* 1993). The other nearest documented record is a photo of one perched at Aquiraz, in northern Ceará, on 16 February 2006 (WA 1947161; A. Campos). Although its main migration route lies west of Rio Grande do Norte, *C. minor* is likely to be found in the state, especially in the ‘serras’ and mangroves.

#### **SICK’S SWIFT** *Chaetura meridionalis*

Seen twice at Engenho Mucambo, Goianinha—one foraging with two Fork-tailed Palm Swifts *Tachornis squamata* and several Barn Swallows *Hirundo rustica* on 5 November 2006 (FSM & MP); and one vocalising over the forest on 30 December 2010 (JFP)—which suggests it is a potential breeder, but documentary evidence to establish its occurrence in the state is lacking.

#### **WHITE-CHINNED SAPPHIRE** *Hylocharis cyanus*

An adult male was observed and its song heard at Mata do Mucambo, Goianinha, on 30 December 2010, and the species was heard again there on 2 January 2011 (JFP), but not documented.

#### **VERSCOLOURED EMERALD** *Amazilia versicolor*

Included in a bird list for Baía Formosa (Olmos 2003), without documentation. Two females, both showing a subterminal dark band on the open tail, were observed by FSM but not documented: at Dunas de Natal State Park, Natal, on 28 November 2002; and mobbing a male Ruby-topaz Hummingbird *Chrysolampis mosquitus* that was competing with it for flying insects at the border of *restinga* at Morro Vermelho, Nísia Floresta, on 17 November 2004.

#### **BARRED FOREST FALCON** *Micrastur ruficollis*

Heard by FSM at Mata da Estrela Private Reserve, Baía Formosa, on 3–4 January 2003, but not documented. Heard at the same locality on 28 February–7 March 2003, where judged uncommon and said to have been sound-recorded (Olmos 2003); however, the observer confirmed that the species was only heard (F. Olmos pers. comm.).

**STREAKED XENOPS** *Xenops rutilans*

One heard singing and two seen at Mata do Mucambo, Goianinha, on 2 January 2011 (JFP & FSM), but not documented.

**RED-HEADED MANAKIN** *Ceratopipra rubrocapilla*

Recorded at Baía Formosa during a survey on 28 February–7 March 2003 (Olmos 2003), but not documented and the species has not been encountered again in this area.

**WHISKERED FLYCATCHER** *Myiobius barbatus*

Identification of *Myiobius* can be difficult (Ridgely & Tudor 2009). The first mention of this genus in the state pertained to Black-tailed Flycatcher *M. atricaudus* seen at Mata da Estrela Private Reserve, Baía Formosa (Olmos 2003). However, records attributed to this species in the Pernambuco Centre of Endemism are likely to reflect misidentification for *M. barbatus*, which occurs—albeit not in sympatry with *atricaudus*—in the far north-east Brazilian Atlantic Forest, whereas *atricaudus* is found in ‘brejos de altitude’ in the Caatinga (RDL pers. obs.). Eight specimens of *M. barbatus* were collected in Paraíba at the border with Rio Grande do Norte in 1957 (Pinto & Camargo 1961), and one, presumably this species, was recently photographed at Mata do Pilão, Espírito Santo (WA 3002212; F. V. Souza), but is not definitely identifiable, and we therefore include the species only on the secondary list. A recent phylogeny of the genus reveals that a specimen of *barbatus* collected in the Pernambuco Centre of Endemism—presumably *M. b. mastacalis*—is sister to the nominate from the Guiana Shield (Azevedo 2017), therefore matching the pattern of several species from the same centre of endemism whose populations are more closely related to those in Amazonia than to the Atlantic Forest south of the São Francisco River (Teixeira 1987, Silveira *et al.* 2003).

**GREY ELAENIA** *Myiopagis caniceps*

One was heard at Mata do Mucambo, Goianinha on 30 December 2010, and it sang at the same place on 2 January 2011 (JFP), but was not documented.

**EULER’S FLYCATCHER** *Lathrotriccus euleri*

Three undocumented records: one *Cnemotriccus*-like tyrant-flycatcher, but with a pale-based bill and long ochre supercilium, perched on a *Sideroxylon obtusifolium* (Sapotaceae) on a sand dune, along with two or more Chilean Elaenias *E. chilensis*—presumably on austral autumn migration—at Guamaré on 27 April 2012 (FSM); one at the ‘serra’ of Venha-Ver (06°19’S, 38°26’W) in July 2012 (Mariano 2014); and one at the ‘serra’ of Martins sometime in 2016 (D. Oliveira). This species, partially migratory in Brazil (Somenzari *et al.* 2018), is expected in the state during the austral winter, and may also be resident in the ‘serras’ of the state, as it is at Serra de Baturité, Ceará (RDL pers. obs.).

**GREEN-WINGED SALTATOR** *Saltator similis*

Much sought after by traffickers and historically released by state environmental agents; one, sound-recorded (WA 3257831, 3257841) at Mata do Mucambo, Goianinha, on 20 January 2019, probably originated from a release and is therefore of unknown origin (RDL & FSM). This locality had previously been studied by ornithologists, without any records of this species until at least 2011, thereby reinforcing the hypothesis of a released or escaped individual.



## YELLOW-FACED SISKIN *Spinus yarrellii*

This Vulnerable species was included as rare in the first state bird list, being specifically mentioned for Agreste (Atlantic Forest / Caatinga ecotone) and the Caatinga (Varela-Freire *et al.* 1999), but we have found no specific records. A claimed first documented record at Florânia (Silva *et al.* 2011) was misidentified and consequently invalid (WA 295444). In the most recent Brazilian Red List (ICMBio / MMA 2018), the species is mapped for two localities in the state, one being the incorrect record just mentioned, while the source of the other is unknown to us. Although rare or disappeared today, the species is well known to those inhabitants interviewed (Pichorim *et al.* 2016a; FSM pers. obs.). It is expensive on the black market, and males are much sought after to cross with the canary, *Serinus canaria*, to produce the hybrid 'pintagol', an excellent singer and consequently the target of systematic trapping in the state. Its tendency towards extinction has been offset since at least 1999 via releases (Dr A. Leite, Cetas / IBAMA Natal, pers. comm.), and today the environmental police immediately release any birds confiscated from traffickers.

## Discussion

**Avifaunal composition.**—In biogeographical terms, the state's avifauna reflects the meeting of the wet lowland Atlantic Forest (with some taxa endemic to the Pernambuco Centre of Endemism) and dry Caatinga (four species typical of this domain) enriched by an enclave of Cerrado (two typical species). For example, East Brazilian Chachalaca *Ortalis araucuan* and *Ramphocelus bresilius* are mainly associated with the Brazilian Atlantic Forest, while typical Caatinga species include *Penelope jacucaca*, *Nyctidromus hirundinaceus*, *Anopetia gounellei* and *Synallaxis hellmayri*. Silvery-cheeked Antshrike *Sakesphorus cristatus*, Caatinga Cacholote *Pseudoseisura cristata*, Campo Troupial *Icterus jamaicai*, Pale Baywing *Agelaioides fringillarius* and White-throated Seedeater *Sporophila albogularis* are also associated with the semi-arid domain. The Cerrado enclave harbours six species mainly associated with this domain (*Heliactin bilophus*, *Xolmis cinereus*, Black-masked Finch *Coryphaspiza melanotis*, White-rumped Tanager *Cypsnagra hirundinacea*, and, especially, Blue Finch *Porphyrospiza caerulescens* and Coal-crested Finch *Charitospiza eucosma*), but *Heliactin* also occurs in coastal 'tabuleiros'. Some species like Rusty-backed Antwren *Formicivora rufa*, Rufous-winged Antshrike *Thamnophilus torquatus* and Lesser Elaenia *Elaenia chiriquensis*, occur not only in the Cerrado enclave, but also in some degraded borders of Atlantic Forest, in stable sand dunes, and in a large area of *restinga* interspersed with 'tabuleiros' that extends from Natal to the north-east corner of the state. Tawny Piculet *Picumnus fulvescens* and Ochraceous Piculet *P. limae* are usually considered typical of the Atlantic Forest and Caatinga, respectively (Moreira-Lima 2013, Araujo & Silva 2017) but, although we follow Piacentini *et al.* (2015; see Table 1), a recent taxonomic reassessment considers them to be extreme phenotypes of a clinal variation (Lima *et al.* 2020).

The state's Atlantic Forest forms part of the Pernambuco Centre of Endemism, one of the main hotspots for threatened bird taxa in South America (Teixeira 1986, Lees *et al.* 2014a). Although the state is poorer in endemic taxa than the southern states, *Penelope superciliaris alagoensis*, *Leptodon forbesi*, *Momotus momota marcgravianus*, *Herpsilochmus [rufimarginatus]* sp., *Xenops minutus alagoanus*, *Conopophaga melanops nigrifrons*, *Platyrrinchus mystaceus niveigularis* and *Hemitriccus griseipectus naumburgae* have already been identified there, and some disjunct populations of Amazonian species occur too, such as *Cercomacroides laeta* and *Myiobius [barbatus]* sp. Note that some other taxa may prove to be more related to Amazonian taxa than to those in the Atlantic Forest (Teixeira 1987, Silveira *et al.* 2003, Ledo & Colli 2017).



Many common species in the coastal Atlantic Forest are absent from the semi-arid Caatinga. However, others (Caatinga Antwren *Herpsilochmus sellowi*, Black-capped Antwren *H. atricapillus*, Ochre-cheeked Spinetail *Synallaxis scutata*, Pectoral Sparrow *Arremon taciturnus* and Flavescent Warbler *Myiothlypis flaveola*) occur in 'brejos de altitude' in the Caatinga. Some of these represent remnants of the ancient corridors that existed between the Amazon and Atlantic forests (Batalha-Filho *et al.* 2013, Ledo & Colli 2017). In Rio Grande do Norte, the 'serras', where the 'brejos de altitude' could occur, are lower (<850 m) than in Ceará (<1,150 m) and Paraíba (<1,190 m), and therefore lack the climatic conditions to support ombrophilous forests, there being only small patches of semi-deciduous forests.

Mangroves in the state harbour some species typical of this environment such as Yellow-crowned Night Heron *Nyctanassa violacea*, Buteogallus *aequinoctialis*, Mangrove Rail *Rallus longirostris*, *Charadrius wilsonia* and Bicoloured Conebill *Conirostrum bicolor*. Others, such as Little Blue Heron *Egretta caerulea*, Little Wood Rail *Aramides mangle*, *Limnodromus griseus*, *Tringa semipalmata* and Plain-bellied Emerald *Amazilia leucogaster*, clearly prefer mangroves and other coastal habitats.

The state of Rio Grande do Norte lies at a subequatorial latitude, which, in conjunction with the coastal effect plus prevailing trade winds, favours the arrival or passage of many species. This is partially reflected by the documentation of two taxa of the same species, e.g., *Thalasseus acuflavidus* (nominate, *eurygnathus*), *Falco peregrinus* (*tundrius*, *anatum*), *Progne tapera* (nominate, *fusca*), with others liable to occur, such as *Charadrius wilsonia* (*crassirostris*, nominate expected), *Tringa semipalmata* (nominate, *inornata* expected), *Chlidonias niger* (presumably nominate, *surinamensis* expected) and *Myiarchus swainsoni* (nominate, *pelzelni* expected).

Except beached birds and the recent use of dataloggers, few data were available on seabirds until the first pelagic birding trips off the state (Lees *et al.* 2015, WikiAves 2019). Numerous other species are expected, and seabirds are likely to account for many future additions to our list (see Potential species).

**Potential species.**—As indicated earlier, the avifauna of Rio Grande do Norte has only recently been studied, consequently we have prepared a list of species likely to occur or to have occurred in the state (see Appendix 2). References concerning these 'potential species', along with those relating to our tertiary list, are presented in Supplementary Material.

Our speculation as to potential species is based on distributional patterns indicated by documented records in conjunction with the availability of suitable habitats in the state. Migrant species, due to the state's geographical position and to the trade winds, can originate from boreal and austral latitudes, and the Old World. This group of species has already produced some ornithological surprises in the state, such as *Hymenops perspicillatus* from the south, and transatlantic *Numenius phaeopus*, thus migrant species deflected by drift are simultaneously exceptional and expected; see the lists for Atol das Rocas Biological Reserve (Schulz Neto 2004b), Fernando de Noronha (Silva e Silva 2008), and some Ceará records, e.g., Northern Gannet *Morus bassanus* (Teixeira *et al.* 2016), Curlew Sandpiper *Calidris ferruginea* (Musher *et al.* 2016), Collared Pratincole *Glareola pratincola* (WA 1672238) and Lesser Black-backed Gull *Larus fuscus* (Girão *et al.* 2006). For such reasons, we have compiled a reasonably extensive list of potential species for which we indicate their estimated probability of occurrence.

The lack of historical data precludes any assessment of the state's avifauna through time. The number of bird species could have been much larger during the early post-Columbian period; for example, toucans probably existed several centuries ago (Silva 2003). Massive clear-cutting of Atlantic Forest and its replacement by sugarcane plantations had a significant negative impact on avian diversity (Teixeira 1986), and it appears very possible

that some species recorded during a visit to Mamanguape in neighbouring Paraíba in 1957 (Pinto & Camargo 1961) also once occurred in nearby Rio Grande do Norte. These include Golden-tailed Parrotlet *Touit surdus* and Swallow-winged Puffbird *Chelidoptera tenebrosa*. Yellow-olive Flycatcher *Tolmomyias sulphurescens*, Planalto Tyrannulet *Phyllomyias fasciatus* and several other forest species are currently known from Guaribas Biological Reserve, at Mamanguape (Almeida & Teixeira 2010). Northern Lesser Woodcreeper *Xiphorhynchus atlanticus* at Mataraca (G. Pereira pers. comm.), and other species at Rio Tinto (Araujo & Nishida 2007), two localities in extreme north-east Paraíba, could occur in south-east Rio Grande do Norte. Groups of Boat-billed Heron *Cochlearius cochlearius* have been observed in nearby Paraíba (Araujo *et al.* 2006; E. Souza pers. comm.), which suggests that this heron also will be found in the state. In the interior, where the Caatinga predominates, semi-deciduous forest enclaves are most promising for discovering new species for the state, e.g., Grey-headed Spinetail *Cranioleuca semicinerea*, Planalto Woodcreeper *Dendrocolaptes platyrostris* and Moustached Woodcreeper *Xiphocolaptes falcirostris*.

The potential list encompasses 92 species (55 non-passerines and 37 passerines) with 16 of high, 36 medium, and 40 low (roughly estimated) probability of being found in the state. Along with the 425 species on the consolidated list, those species with high or medium chance of occurrence in the state produce a reasonable potential total of c.480 species in Rio Grande do Norte and its continental waters. But it should be borne in mind that, for now, just 391 have been recorded on the primary list.

**Threats to the state's avifauna.**—Our consolidated list includes a noteworthy number of threatened taxa: ten globally and 21 nationally threatened taxa, and 11 globally and seven Near Threatened, all of which depend on conservation of forests (Atlantic Forest, *restinga* and mangroves, 'serras', and arboreal Caatinga) and the Cerrado. In addition, the lack of respect for federal anti-hunting legislation threatens *Crypturellus noctivagus zabele*, *Penelope superciliaris alagoensis*, *P. jacucaca*, *Spinus yarrellii*, seven species of waders, and eight pelagic species. Other species of conservation concern are *Leptodon forbesi* (Critically Endangered), *Herpsilochmus pectoralis* and *Coryphaspiza melanotis* (both Vulnerable).

We trust this paper will become a basic reference for the construction of a state Red List of birds to be protected by new environmental laws. Simultaneously, restoration initiatives linked to compliance with environmental laws and conservation of water resources should and could be linked to conservation projects for these species, aiming to create new habitat corridors.

As noted by Lees *et al.* (2015), the area at the shelf break is rich in pelagic birds that may suffer some persecution (*cf.* Lima *et al.* 2004). For example, we have encountered a *Puffinus gravis* with its left wing ensnared by a fish hook and nylon line; a *P. puffinus* lacking one wing, probably cut by fishermen after it became caught in their nets; a *Fregetta tropica* with one leg; an *Oceanites oceanicus* lacking half of the primaries in one wing, perhaps caused by a shotgun (or by a jaeger); two *Sula dactylatra* caught on fish hooks; a *Stercorarius maccormicki* captured alive and released, trapped again and kept captive, and another caught on a fish hook and also held captive.

The state's mangroves swamps were partially destroyed centuries ago by the salt industry and since the 1970s by shrimp producers. Today, egg collecting persists in and around some saltpans. Although federal law does not permit hunting, poaching is common in the state and this activity has provided recoveries of ringed birds, e.g., *Falco peregrinus*, waders, ducks (*Anas discors*) and other waterbirds. There is intense pressure on nesting colonies of *Zenaida auriculata*, and we have regularly witnessed the hunting and sale of Tataupa Tinamou *Crypturellus tataupa*, *Penelope jacucaca*, *P. superciliaris*, *Nyctanassa violacea*, Great Egret *Ardea alba*, *Elanoides forficatus*, Black-chested Buzzard-Eagle *Geranoaetus*

*melanoleucus*, Roadside Hawk *Rupornis magnirostris*, Common Gallinule *Gallinula galeata*, Blue Ground Dove *Claravis pretiosa*, all of the *Columbina* species, many White-tipped Doves *Leptotila verreauxi*, and *Conirostrum bicolor* (Pichorim *et al.* 2016a; FSM pers. obs.).

The caging and trafficking of wild birds creates an endless drain on groups such as doves, psittacids and passerines (Pichorim *et al.* 2014b, 2016a; FSM pers. obs.). All eight species of psittacids on the primary list have been observed in cages, and many end up in the IBAMA / Natal rehabilitation centre or Natal Zoo. Among passerines, the most impacted families are Icteridae, Thraupidae, Turdidae, Mimidae, Cardinalidae and Fringillidae. Of the latter, *Spinus yarrellii* is on the verge of extinction, as it is over most of its range. White-naped Jay *Cyanocorax cyanopogon* is a common cagebird too. These pressures lead to local decreases and the disruption of the natural distribution of species, e.g., some psittacids, *Cyanocorax*, *Ramphocelus bresilius*, *Cacicus cela* and *Saltator similis*, due to frequent release by local people and environmental agencies of thousands of birds in inappropriate habitats. Also, some non-native species have been erroneously released, such as Seven-coloured Tanager *Tangara fastuosa*.

Windfarms kill raptors, migrants such as *Limnodromus griseus*, *Arenaria interpres* and Semipalmated Plover *Charadrius semipalmatus*, gulls, doves, cuckoos, hummingbirds, etc. (FSM & JBI pers. obs.). There are now thousands of wind turbines along almost the entire north coast and the northern half of the east coast, as well as the tops of some 'serras'. Their full impacts remain to be assessed, as existing monitoring schemes are inadequate.

The most threatened and already impoverished avifauna is surely that of the Atlantic Forest, which is subject to constant loss of its small remnants to sugarcane plantations and, more recently, by touristic developments. Moreover, some of these species are sought after as cagebirds and edible gamebirds, such as the threatened *Penelope supercilialis alagoensis*.

In this article we present the first state bird list based on strict criteria for compilation and documentation (Carlos *et al.* 2010, Lees *et al.* 2014b). It also represents the first critical review of all published records, from which obvious errors and doubtful cases have been removed, following careful consideration, and completed with unpublished records. Our consolidated list of the birds of Rio Grande do Norte encompasses 425 species, with 391 species on the primary list and 34 species on the secondary list.

Species lists are ongoing, never-ending exercises due to the dynamic nature of avian communities and the factors influencing them such as changing weather patterns, human population growth and the destruction of ecosystems. This is the first step in consolidating a state list and we hope that it will spark interest in further studies of the Rio Grande do Norte avifauna.

#### Acknowledgements

This work is dedicated to the memory of Antonio Adalberto Varella-Freire, one of the pioneers of field zoology in Rio Grande do Norte. We thank Severino Mendes de Azevedo, Luciano N. Naka and Victor Leandro Silva (UFPE), Helder F. P. de Araujo and Nayla Fábila F. do Nascimento (CAHZ), Luís Fábio Silveira (MZUSP), Philip Unitt (SDNHM), James V. Remsen and Steve Cardiff (LSUMZ), Christopher Milensky (USNM) and Brett W. Benz (UMMZ), who sent information and photos of specimens held in their collections; Simone Dena for information from the FNJV archive; MCZ, MPEG and MNRJ for providing online data from their collections; William Pitts / BBL for information on resighted waders, Jeff Spindel (North America), Judy Pierce (US Virgin Islands) and Norman Ratcliffe (Europe) for information about Roseate Terns, SNA (ICMBio / CEMAVE) for data on banded bird recoveries, Luiz Mestre for information on banded Solitary Sandpipers, and all contributors and administrators of online databases, especially Bruno França and WikiAves, who provided numerous new records for the avifauna of Rio Grande do Norte and neighbouring states. RDL thanks Ms Dione Seripierri for many references used in this work. We are grateful to Fábio Olmos for identification of some seabirds, Roberto Otoch, Weber Girão, Ciro Albano, Robson Silva e Silva, Luciano Moreira Lima and Glauco Pereira for correspondence and references, Deodato Souza, Albano Schulz-Neto, Marcelo da Silva and Damião Oliveira for some data, and C. E. Agne for identification of *Tachycineta leucopyga*. We thank the botanist Jomar Jardim (identification of *Psidium*), the always reliable

Valdenir Andrade for information on birds brought to Pipa Ecological Sanctuary, and Flávio Lima, director of 'Project Cetaceans of the Costa Branca' at UERN, Mossoró, for authorising use of unpublished data. Maurizélia de Brito, director of Atol das Rocas Biological Reserve, provided information about seabirds. Many thanks to Rob Bierregaard for permission to mention the results and a photo from his project on Ospreys. We are grateful to the manager of Salinor (Macau) who sanctioned an important bird census there. Special thanks to biologist Francisco de Assis of Dunas de Natal State Park, Adson Borges Machado, director of ESEC Seridó in 2002, and Damião Dantas de Souza, director of the Açú National Forest, and José Fernando Neto for accommodation and sharing relevant data. Rosemeire Dantas accompanied FSM on many visits to Guamaré and the Potengi estuary. Claudio Ferreira, another faithful companion during many field trips, along with Luiz Sena, Zelito Coringa, Jules Sagot, José Luís C. Novaes, Carolina T. Puppim-Gonçalves, Ana Carolina C. Xavier, Douglas Brandão, Pedro T. Moura, Estevão Vieira and Jan van den Bosch permitted us to publish their photos. Iracema Miranda and family proffered their constant support of our field work on the state's north coast. We have been informed, guided in the field, transported and permitted to research birds by dozens of kind and receptive people of all ages, who often accommodated us in their homes; the number far exceeds the limits of this publication, but we are pleased to remember Antonio Francisco and Nira in Mossoró, Seu Berino, 'Lula' and 'Baia' in Canguaretama, Ivo Cortez in Macau, 'Toninho' in Guamaré, Aluisio in Galinhos, Luiz Sena and family in Parelhas and Equador, Chico de Otávio in Martins, Genaro Moreira de Carvalho in Coronel João Pessoa, Oscar and his father, and Luís Belo and his father Pedro in Luís Gomes, Lydia Brasileira, Francisco Reis de Araújo and Lucas Gaehwiler in Jucurutu, Estevão Vieira and Sandra Magalhães in Maracaju, Chico do Fole in Tenente Laurentino Cruz, Antônio Amaro Sobrinho, 'Pelé' Amaro, Chico Amaro de Oliveira and family, and Del de Alice in Santana do Matos, the Pereira family in Cerro Corá, José Zacarias dos Santos, 'Sapo' and the Rabelo family in Baía Formosa, Ricardo 'Cobra' Tersuliano's parents in the *restinga* of Maxaranguape, 'Caboco' in Poço Branco, Juan Pablo B. in Serra Caiada and Sítio Novo, 'Canindé' at Papagaio Farm and Sr. Diapino and his son Francisco de Assis at Barra da Onça Farm / Santana do Matos, Juan Carlos Vargas companion on field trips in Lajes and Sr. Lorival, his son João and grandson João Pedro in Boa Vista / Lajes, Dr Frederico Lima in Mucambo / Goianinha, Maurício Galvão Meira e Sá and Sr. Raimundo in Bico da Arara / Acari. In France, FSM received great support from Alain Sagault, Geneviève Boussus, Pascale, Philippe and André Sagot. Debbie Robertson offered her very kind help with the English version of the manuscript. We warmly thank Fernando Straube, Fábio Olmos and especially Guy Kirwan, for their extensive reviews, comments and suggestions.

#### References:

- Almeida, A. C. C. & Teixeira, D. M. 2010. Aves da Reserva Biológica Guaribas, Mamanguape, Paraíba, Brasil. *Rev. Nordestina Biol.* 19(2): 3–14.
- Alves, G. 1977. Provincia Fluminis Grandis. *Rev. Inst. Ceará* 91: 57–89.
- Amarante, O. A. C., Silva, F. J. L., Rios Filho, L. G. & Vidal, M. J. B. 2003. *Potencial eólico do Estado do Rio Grande do Norte*. Companhia Energética do Rio Grande do Norte, Natal.
- Andrade, M. A. 1991. Águia-pescadora ameaçada. *Ciência Hoje* 13(75): 60–61.
- Andrade, M. A. & Freitas, M. V. 1987. O enigma dos andorinhões. *Ciência Hoje* 37: 8–9.
- Andrade-Lima, D. 1982. Present-day forest refuges in northeastern Brazil. Pp. 123–135 in Prance, G. T. (ed.) *Biological diversification in the tropics*. Columbia Univ. Press, New York.
- Antas, P. T. Z. 1987a. A nidificação no chão da rolinha-roxa-mirim, *Columbina minuta*, em Pedro Avelino, Rio Grande do Norte. *Rev. Bras. Zool.* 3: 465–466.
- Antas, P. T. Z. 1987b. A nidificação da avoante no Nordeste do Brasil, relacionada com o substrato fornecido pela vegetação. *Rev. Bras. Zool.* 3: 467–470.
- Antas, P. T. Z. 1994. Migration and other movements among the lower Paraná River valley wetlands, Argentina, and the south Brazil/Pantanal wetlands. *Bird Conserv. Intern.* 4: 181–190.
- Araújo, F. A. A., Vianna, R. S. T., Vada, M. Y., Silva, E. V., Doretto, L., Cavalcante, G. C., Azevedo, S. M. J., Magalhães, V. S., Gomes, J. L., Queiroz, P. V. S., Larrazábal, M. E., Martins, L. C., Rodrigues, S. G. & Vasconcelos, P. F. C. 2004. Inquérito sorológico em aves migratórias e residentes de Galinhos, RN, para detecção do vírus da Febre do Nilo Ocidental e outros vírus. *Bol. Eletrônico Epidemiológico* 4(2): 1–12.
- Araujo, H. F. P. & Nishida, A. K. 2007. Considerações sobre colônias de garças (Ciconiiformes, Ardeidae) no Estado da Paraíba, Brasil. *Ornithologia* 2: 34–40.
- Araujo, H. F. P. & Silva, J. M. C. 2017. The avifauna of the Caatinga: biogeography, ecology, and conservation. Pp. 181–210 in Silva, J. M. C., Leal, I. R. & Tabarelli, M. (eds.) *Caatinga: the largest tropical dry forest region in South America*. Springer, Berlin.
- Araujo, H. F. P., Rodrigues, R. C. & Nishida, A. K. 2006. Composição da avifauna em complexos estuarinos no Estado da Paraíba, Brasil. *Rev. Bras. Orn.* 14: 249–259.
- Azevedo, S. B. 2017. Revisão sistemática do gênero *Myiobius* Gray, 1839 (Aves: Onychorhynchidae). M.Sc. thesis. Museu Paraense Emílio Goeldi & Universidade Federal do Pará, Belém.
- Azevedo, S. M. J. & Antas, P. T. Z. 1990. Técnicas de captura para o anilhamento de *Zenaida auriculata noronha*. Pp. 78–84 in Azevedo, S. M. J. (ed.) *An. IV Encontro Nac. Anilhadores Aves-ENAV (1988)*. Univ. Federal de Pernambuco, Recife.



- Azevedo, S. M. J., Dias-Filho, M. M., Larrazábal, M. E., Lyra-Neves, R. M. & Telino-Jr, W. R. 2000. Movimentação de *Progne chalybea* (Aves: Hirundinidae) em Pernambuco, Brasil. Pp. 149–159 in Alves, M. A. S., Silva, J. M. C., Sluys, M. V., Bergallo, H. G. & Rocha, C. F. D. (ed.) *A ornitologia no Brasil, pesquisa atual e perspectivas*. Ed. Univ. Estadual do Rio de Janeiro, Rio de Janeiro.
- Azevedo, S. M. J., Larrazábal, M. E. & Pena, O. 2003. First record of *Charadrius melodus* Ord (Aves Charadriiformes) in Brazil. *Rev. Bras. Zool.* 20: 559–560.
- Azevedo, S. M. J., Larrazábal, M. E. & Pena, O. 2004. Aves aquáticas de ambientes antrópicos (salinas) do Rio Grande do Norte, Brasil. Pp. 255–266 in Branco, J. O. (ed.) *Aves marinhas e insulares brasileiras: bioecologia e conservação*. Ed. Univ. do Vale do Itajaí, Santa Catarina.
- Baker, F. 1913. The land and fresh-water mollusks of the Stanford Expedition to Brazil. *Proc. Acad. Nat. Sci. Phil.* 65: 618–672.
- Batalha-Filho, H., Fjeldså, J., Fabre, P. & Miyaki, C. Y. 2013. Connections between the Atlantic and the Amazonian forest avifaunas represent distinct historical events. *J. Orn.* 154: 41–50.
- Bierregaard, R. O., Cohn-Haft, M., Stotz, D. F. & Teixeira, D. M. 1997. Cryptic biodiversity: an overlooked species and new subspecies of antbird (Formicariidae) with a revision of *Cercomacra tyrannina* in northwestern South America. Pp. 111–128 in Remsen, J. V. (ed.) *Studies in Neotropical ornithology honoring Ted Parker*. Orn. Monogr. 48.
- Brown, K. S. 1982. Historical and ecological factors in the biogeography of aposematic Neotropical butterflies. *Amer. Natur.* 22: 453–471.
- Bugoni, L. & Furness, R. W. 2009. Ageing immature Atlantic Yellow-nosed *Thalassarche chlororhynchos* and Black-browed *T. melanophris* albatrosses in wintering grounds using bill colour and moult. *Mar. Orn.* 37: 249–252.
- Burger, J. 2018. *Birdlife of the Gulf of Mexico*. Texas A&M University Press, College Station.
- Burney, C. W. 2009. Comparative phylogeography of Neotropical birds. Ph.D. thesis. Louisiana State Univ., Baton Rouge.
- Carboneras, C., Jutglar, F. & Kirwan, G. M. 2019. Audubon's Shearwater (*Puffinus lherminieri*). In del Hoyo, J., Elliott, A., Sargatal, J., Christie, D. A. & de Juana, E. (eds.) *Handbook of the birds of the world Alive*. Lynx Edicions, Barcelona (retrieved from <https://www.hbw.com/node/52580> in July 2019).
- Carlos, C. J., Trimble, J. & Fedrizzi, C. E. 2004. A specimen of Cape Petrel *Daption capense* from north-east Brazil. *Bull. Brit. Orn. Cl.* 124: 287–290.
- Carlos, C. J., Straube, F. C. & Pacheco, J. F. 2010. Conceitos e definições sobre documentação de registros ornitológicos e critérios para a elaboração de listas de aves para os estados brasileiros. *Rev. Bras. Orn.* 18: 355–361.
- Cascardo, L. C. 1968. *Nomes da terra: geografia, história e toponímia do Rio Grande do Norte*. Sebo Vermelho, Natal.
- Cavalcanti, L. M. P., Paiva, L. V. & França, L. F. 2016. Effects of rainfall on bird reproduction in a semi-arid Neotropical region. *Zoologia* 33(6): 1–66.
- Cavarzere, V. A. 2014. Taxonomy, phylogeny and distribution of *Cercomacra* Sclater, 1858 (Aves: Thamnophilidae). Ph.D. thesis. Univ. de São Paulo, São Paulo.
- Comitê Brasileiro de Registros Ornitológicos (CBRO). 2009. Lista das aves do Brasil (v. 09 August 2009). <http://www.cbro.org.br>.
- Comitê Brasileiro de Registros Ornitológicos (CBRO). 2010. Lista das aves do Brasil, 9ª edn. (v. 18 October 2010). <http://www.cbro.org.br>.
- Comitê Brasileiro de Registros Ornitológicos (CBRO). 2011. Lista das aves do Brasil, 10ª edn. (v. 25 January 2011). <http://www.cbro.org.br>.
- Comitê Brasileiro de Registros Ornitológicos (CBRO). 2014. Lista das aves do Brasil, 11ª edn. <http://www.cbro.org.br>.
- Coelho, A. G. M. 1978. Lista de algumas espécies de aves do Nordeste do Brasil. *Notul. Biol.* 1: 1–7.
- Cordeiro, P. H. C., Flores, J. M. & Nascimento, J. L. X. 1996. Análise das recuperações de *Sterna hirundo* no Brasil entre 1980 e 1994. *Ararajuba* 4: 3–7.
- Cruz-Flores, M., Ramos, R., Sardà-Serra, M., López-Souto, S., Militão, T. & González-Solís, J. 2019. *Migración y ecología espacial de la población española de petrel de Bulwer*. SEO / BirdLife, Madrid.
- Dantas, G. P. M., Almeida, V. S., Maracini, P., Serra, S. D., Chame, M., Labarthe, N., Kolesnikovas, C., Siciliano, S., Matias, C. A. R., Moura, J. F., Campos, S. D. E., Mader, A. & Serafini, P. P. 2013. Evidence for northward extension of the winter range of Magellanic Penguins along the Brazilian coast. *Mar. Orn.* 41: 195–197.
- Dénes, F. V., Silveira, L. F., Seipke, S. H., Thorstrom, R., Clark, W. S. & Thiollay, J.-M. 2011. The White-collared Kite (*Leptodon forbesi* Swann, 1922) and a review of the taxonomy of the Gray-headed Kite (*Leptodon cayanensis* Latham, 1790). *Wilson J. Orn.* 123: 326–331.
- Diniz, M. T. M. & Pereira, V. H. 2015. Climatologia do Estado do Rio Grande do Norte, Brasil: sistemas atmosféricos ativos atuantes e mapeamento de tipos de clima. *Bol. Goiano Geogr.* 35: 488–506.
- Espírito Santo, A. R. S. & Silva, C. M. S. 2016. Características climáticas da Cidade de Natal. *Parque Cidade Rev.* 6(1): 1–15.
- Faria, O. L. 1961. *A caça nos sertões do Seridó*. Serviço de Informação Agrícola, Rio de Janeiro.

- Faria, O. L. 2001. *Notas de carregação*. Coleção Nação Potiguar, Fundação Hélio Galvão, Rio de Janeiro.
- Farnsworth, A., Langham, G. & Boesman, P. 2019. Velvet Black-tyrant (*Knipolegus nigerrimus*). In del Hoyo, J., Elliott, A., Sargatal, J., Christie, D. A. & de Juana, E. (eds.) *Handbook of the birds of the world Alive*. Lynx Edicions, Barcelona (retrieved from <https://www.hbw.com/node/57396> in June 2019).
- Ferguson-Lees, J. & Christie, D. A. 2001. *Raptors of the world*. Christopher Helm, London.
- Fitzpatrick, J. 2019. Planalto Tyrannulet (*Phyllomyias fasciatus*). In del Hoyo, J., Elliott, A., Sargatal, J., Christie, D. A. & de Juana, E. (eds.) *Handbook of the birds of the world Alive*. Lynx Edicions, Barcelona (retrieved from <https://www.hbw.com/node/57120> in June 2019).
- Gill, F. & Donsker, D. (eds.) 2019. IOC world bird list (v 9.2). <https://doi.org/10.14344/IOC.ML.9.2>.
- Geiger, R. 1961. *Überarbeitete Neuauflage von Geiger, R.: Köppen-Geiger / Klima der Erde*. (Wandkarte 1:16 Mill.) Klett-Perthes, Gotha.
- Girão, W. & Albano, C. 2011. Região do Banco dos Cajuais. Pp. 137–140 in Valente, R. M., Silva, J. M. C., Straube, F. C. & Nascimento, J. L. X. (eds.) *Conservação de aves migratórias Neárticas no Brasil*. Conservação Internacional, Belém.
- Girão, W., Albano, C., Pinto, T., Campos, A., Meirelles, A. C. & Silva, C. P. N. 2006. First record of the Lesser Black-backed Gull *Larus fuscus* Linnaeus, 1758 for Brazil. *Rev. Bras. Orn.* 14: 463–464.
- Grantsau, R. K. H. 2010. *Guia completo para a identificação das aves do Brasil*. Ed. Vento Verde, São Paulo.
- Harvey, M. G. & Brumfield, R. T. 2015. Genomic variation in a widespread Neotropical bird (*Xenops minutus*) reveals divergence, population expansion, and gene flow. *Mol. Phyl. & Evol.* 83: 305–316.
- Hasui, É., Metzger, J. P., Pimentel, R. G., Silveira, L. F., Bovo, A. A., Martensen, A. C., Uezu, A., Regolin, A. L., Bispo de Oliveira, A. Á., Gatto, C. A., Duca, C., Andretti, C. B., Banks-Leite, C., Luz, D., Mariz, D., Alexandrino, E. R., Barros, F. M., Martello, F., Pereira, I. M., Silva, J. N., Ferraz, K. M., Naka, L. N., Anjos, L., Efe, M. A., Pizo, M. A., Pichorim, M., Gonçalves, M. S., Cordeiro, P. H., Dias, R. A., Muylaert, R. D., Rodrigues, R. C., Costa, T. V., Cavarzere, V., Tonetti, V. R., Silva, W. R., Jenkins, C. N., Galetti, M. & Ribeiro, M. C. 2018. Atlantic birds: a data set of bird species from the Brazilian Atlantic Forest. *Ecology* 99: 497–497.
- Hellmayr, C. E. 1934. Catalogue of birds of the Americas and the adjacent islands, pt. 7. *Publ. Field Mus. Nat. Hist. Zool. Ser.* 13(330).
- Howell, S. N. G. 2010. Identification and taxonomy of White-bellied Storm Petrels, with comments on WP report in August 1986. *Dutch Birding* 32: 36–42.
- del Hoyo, J., Collar, N. & Kirwan, G. M. 2019a. Northern Rufous-winged Antwren (*Herpsilochmus scapularis*). In del Hoyo, J., Elliott, A., Sargatal, J., Christie, D. A. & de Juana, E. (eds.) *Handbook of the birds of the world Alive*. Lynx Edicions, Barcelona (retrieved from <https://www.hbw.com/node/1343587> in June 2019).
- del Hoyo, J., Collar, N. & Kirwan, G. M. 2019b. Silvery-flanked Antwren (*Myrmotherula luctuosa*). In del Hoyo, J., Elliott, A., Sargatal, J., Christie, D. A. & de Juana, E. (eds.) *Handbook of the birds of the world Alive*. Lynx Edicions, Barcelona (retrieved from <https://www.hbw.com/node/1343585> in June 2019).
- del Hoyo, J., Collar, N. & Kirwan, G. M. 2019c. Plain Xenops (*Xenops genibarbis*). In del Hoyo, J., Elliott, A., Sargatal, J., Christie, D. A. & de Juana, E. (eds.) *Handbook of the birds of the world Alive*. Lynx Edicions, Barcelona (retrieved from <https://www.hbw.com/node/1343639> in June 2019).
- Instituto Chico Mendes de Biodiversidade/Ministério do Meio Ambiente. 2018. Vol. 3 – Aves. In ICMBio / MMA (ed.) *Livro Vermelho da fauna brasileira ameaçada de extinção*, 1ª ed. ICMBio / MMA, Brasília.
- Instituto Nacional de Meteorologia (INMET). 2019. [www.inmet.gov.br](http://www.inmet.gov.br) (accessed 28 April 2019).
- IUCN. 2019. The IUCN Red List of threatened species 2019-1. <https://www.iucnredlist.org>.
- Irusta, J. B. & Sagot-Martin, F. 2011. Complexo litorâneo da Bacia Potiguar. Pp. 141–145 in Valente, R. M., Silva, J. M. C., Straube, F. C. & Nascimento, J. L. X. (eds.) *Conservação de aves migratórias Neárticas no Brasil*. Conservação Internacional, Belém.
- König, C., Weick, F. & Becking, J.-H. 2009. *Owls of the world*. Second edn. Christopher Helm, London.
- Lamm, D. W. 1948. Notes on the birds of the states of Pernambuco and Paraíba, Brazil. *Auk* 65: 261–283.
- Lanyon, W. E. 1978. Revision of the *Myiarchus* flycatchers of South America. *Bull. Amer. Mus. Nat. Hist.* 161: 427–628.
- Lara-Resende, S. M. & Leal, R. P. 1982. Recuperação de anilhas estrangeiras no Brasil. *Brasil Florestal* 12(52): 27–53.
- Larrazábal, M. E., Azevedo, S. M. J. & Pena, O. 2002. Monitoramento de aves limícolas na Salina Diamante Branco, Galinhos, Rio Grande do Norte, Brasil. *Rev. Bras. Zool.* 19: 1081–1089.
- Ledo, R. M. D. & Colli, G. R. 2017. The historical connections between the Amazon and the Atlantic Forest revisited. *J. Biogeogr.* 44: 2551–2563.
- Lees, A. C., Albano, C., Kirwan, G. M., Pacheco, J. F. & Whittaker, A. 2014a. The end of hope for Alagoas Foliage-gleaner *Philydor novaesi*? *Neotrop. Birding* 14: 20–28.
- Lees, A. C., Naka, L. N., Aleixo, A., Cohn-Haft, M., Piacentini, V. Q., Santos, M. P. D. & Silveira, L. F. 2014b. Conducting rigorous avian inventories: Amazonian case studies and a roadmap for improvement. *Rev. Bras. Orn.* 22: 107–120.
- Lees, A. C., Olmos, F. & Campos, A. 2015. Here be gadflies: pelagic birding off north-east Brazil. *Neotrop. Birding* 17: 11–18.



- Lima, P. C., Grantsau, R., Lima, R. C. F. R. & Santos, S. S. 2004. Occurrence and mortality of seabirds along the northern coast of Bahia, and the identification key of the Procellariiformes order and the Stercorariidae family. *Atualidades Orn.* 121: 1–62.
- Lima, R. D., Dantas, H. R. & Oliveira, D. V. 2018. Ocorrência de *Bubo virginianus* (Strigiformes: Strigidae) no extremo nordeste do Brasil: primeiro registro documentado para o Estado do Rio Grande do Norte e revisão dos registros no Nordeste brasileiro. *Atualidades Orn.* 206: 11–15.
- Lima, R. D., Tomotani, B. M. & Silveira, L. F. 2020. Colour variation and taxonomy of *Picumnus limae* Sneathlage, 1924 and *P. fulvescens* Stager, 1961 (Piciformes: Picidae). *J. Orn.* 161: 491–501.
- Macedo, H. A. M. 2003. Expedições arqueológicas em Carnaúba dos Dantas, Rio Grande do Norte, Brasil: resultados das prospecções realizadas entre 1996 e 1997. *Cadernos Centro Mem. Oeste Santa Catarina* 17(18): 81–216.
- Marcgrave, G. 1648. *Historiae rerum naturalium Brasiliae*. Lundunum Batavorum, Amsterdam.
- Mariano, E. F. 2014. Relações biogeográficas entre a avifauna de florestas de altitude no Nordeste do Brasil. Ph.D. thesis. Univ. Federal da Paraíba, João Pessoa.
- Medeiros, J. A., Silveira, A. G. & Varela-Freire, A. A. 2000. Zoogeografia do Rio Grande do Norte VI: listagem provisória das aves do açude Riacho do Roçado, São José do Seridó, Rio Grande do Norte. *Sociedade e Território* 14: 29–34.
- Mestre, L. A. M. 2007a. Recuperações no Brasil de falcões-peregrinos (*Falco peregrinus*) anilhados na América do Norte entre 1967 e 2001. *Ornithologia* 2: 72–80.
- Mestre, L. A. M. 2007b. Registros das migrações de trinta-réis-boreal *Sterna hirundo*: análise das recuperações de indivíduos marcados na América do Norte (1928–2005) e Brasil (1983–2005). *Ornithologia* 2: 81–87.
- Mestre, L. A. M. & Bierregaard, R. O. 2009. The role of Amazonian rivers for wintering ospreys (*Pandion haliaetus*): clues from North American band recoveries in Brazil between 1937 and 2006. *Stud. Neotrop. Fauna & Environ.* 44: 141–147.
- Mestre, L. A. M., Roos, A. L. & Nunes, M. F. 2010. Análise das recuperações no Brasil de aves anilhadas no exterior entre 1927 e 2006. *Ornithologia* 4: 15–35.
- Ministério do Meio Ambiente (MMA). 2003. Instrução normativa n° 3, de 27/05/2003. Lista das espécies terrestres da fauna brasileira ameaçadas de extinção segundo o workshop da Fundação Biodiversitas de dezembro de 2002 com categorias IUCN. Diário Oficial da União, Seção 1 101: 88–97, Brasília, Distrito Federal.
- Ministério do Meio Ambiente (MMA). 2004. Aves. Pp. 52–77 in *Plano de Manejo da Estação Ecológica do Seridó*. MMA, Brasília.
- Minns, J., Buzzetti, D., Albano, C., Whittaker, A., Grosset, A. & Parrini, R. 2010. *Birds of Brazil / Aves do Brasil*. DVD-ROM. Ed. Avis Brasilis, Vinhedo.
- Moreira-Lima, L. 2013. Aves da Mata Atlântica: riqueza, composição, status, endemismos e conservação. M.Sc. thesis. Univ. de São Paulo, São Paulo.
- Murphy, R. C. 1936. *Oceanic birds of South America*. Amer. Mus. Nat. Hist., New York.
- Musher, L. J., Almeida, B. J. M., Fedrizzi, C. E., Holderbaum, J. M., Lees, A. C., Mizrahi, D. & Rodrigues, R. C. 2016. Curlew Sandpipers *Calidris ferruginea* in the western Atlantic: the first, second and third Brazilian records from Ceará and Maranhão. *Rev. Bras. Orn.* 24: 62–67.
- Nacinovic, J. B. & Teixeira, D. M. 1989. As aves de Fernando de Noronha - uma lista sistemática anotada. *Rev. Bras. Biol.* 49: 709–729.
- Nascimento, J. L. X. 2000. Estudo comparativo da avifauna em duas estações ecológicas da Caatinga: Aiuaba e Seridó. *Melospittacus* 3: 12–35.
- Nascimento, J. L. X. & Larrazábal, M. E. L. 2000. Alimentação de aves limícolas em Barra de Cunhaú, Canguaretama, Rio Grande do Norte. *Melospittacus* 3: 91–109.
- O'Brien, M., Crossley, R. & Karlson, K. 2006. *The shorebird guide*. Houghton Mifflin, New York.
- Oliveira, A. C. P., Penha, A. S., Souza, R. F. & Loiola, M. I. B. 2012. Floristic composition of a savanna community in Rio Grande do Norte, northeastern Brazil. *Acta Bot. Bras.* 26: 559–569.
- Olmos, F. 2000. Revisão dos registros de *Fregetta tropica* para o Brasil (Procellariiformes: Hydrobatidae). *Nattereria* 1: 27–28.
- Olmos, F. 2002. Non-breeding seabirds in Brazil: a review of band recoveries. *Ararajuba* 10: 31–42.
- Olmos, F. 2003. Birds of Mata Estrela private reserve, Rio Grande do Norte, Brazil. *Cotinga* 20: 23–30.
- Olsen, P. D., Kirwan, G. M., Christie, D. A. & Marks, J. S. 2019. Short-eared Owl (*Asio flammeus*). In del Hoyo, J., Elliott, A., Sargatal, J., Christie, D. A. & de Juana, E. (eds.) *Handbook of the birds of the world Alive*. Lynx Edicions, Barcelona (retrieved from <https://www.hbw.com/node/55128> in June 2019).
- Oswald, J. A., Harvey, M. G., Remsen, R. C., Foxworth, D. U., Cardiff, S. W., Dittmann, D. L., Megna, L. C., Carling, M. D. & Brumfield, R. T. 2016. Willet be one species or two? A genomic view of the evolutionary history of *Tringa semipalmata*. *Auk* 133: 593–614.
- Pacheco, J. F. 2004. As aves da Caatinga – uma análise histórica do conhecimento. Pp. 189–240 in Silva, J. M. C., Tabarelli, M., Fonseca, M. T. & Lins, L. V. (eds.) *Biodiversidade da Caatinga: áreas e ações prioritárias para a conservação*. Ministério do Meio Ambiente, Brasília.

- Pacheco, J. F. & Whitney, B. M. 1995. Range extensions for some birds in northeastern Brazil. *Bull. Brit. Orn. Cl.* 115: 157–163.
- Paiva, M. P. 1983. Algumas considerações sobre a fauna da região semi-árida do Nordeste brasileiro. *Coleção Mossoroense, Sér. B* 404: 1–31.
- Pereira, G. A., Lobo-Araújo, L. W., Leal, S., Medcraft, J. P., Marantz, C. A., Toledo, M. T. F., Araujo, H. F. P., Albano, C., Pinto, T., Santos, C. H. A., Serapião, L. C. H., Malacco da Silva, G. B. & Pioli, D. 2012. Important bird records from Alagoas, Pernambuco and Paraíba, north-east Brazil. *Cotinga* 34: 17–22.
- Piacentini, V. Q. 2011. Taxonomia e distribuição geográfica dos representantes do gênero *Phaethornis* Swainson, 1827 (Aves: Trochilidae). Ph.D. thesis. Univ. de São Paulo, São Paulo.
- Piacentini, V. Q., Aleixo, A., Agne, C. E., Maurício, G. N., Pacheco, J. F., Bravo, G. A., Brito, G. R. R., Naka, L. N., Olmos, F., Posso, S., Silveira, L. F., Betini, G. S., Carrano, E., Franz, I., Lees, A. C., Lima, L. M., Pioli, D., Schunck, F., Amaral, F. S. R., Bencke, G. A., Cohn-Haft, M., Figueiredo, L. F. A., Straube, F. C. & Cesari, E. 2015. Annotated checklist of the birds of Brazil by the Brazilian Ornithological Records Committee / Lista comentada das aves do Brasil pelo Comitê Brasileiro de Registros Ornitológicos. *Rev. Bras. Orn.* 23: 91–298.
- Pichorim, M., Silva, M., França, B. R. A., Oliveira-Jr, T. M. & Rodrigues, M. C. 2014a. A Cerrado bird community in the northernmost portion of northeastern Brazil - recommendations for conservation. *Rev. Bras. Orn.* 22: 347–362.
- Pichorim, M., Câmara, T. P. F., Oliveira-Jr, T. M., Oliveira, D. V., Nascimento, E. P. G. & Mobley, J. A. 2014b. A population of Blue-winged Macaw *Primolius maracana* in northeastern Brazil: recommendations for a local Conservation Action Plan. *Trop. Conserv. Sci.* 7: 488–507.
- Pichorim, M., Oliveira, D. V., Oliveira-Jr, T. M., Câmara, T. P. F. & Nascimento, É. P. G. 2016a. Pristine semi-arid areas in northeastern Brazil remain mainly on slopes of mountain ranges: a case study based on bird community of Serra de Santana. *Trop. Zool.* 29: 189–204.
- Pichorim, M., Damasceno, J. P. T. & Toledo-Lima, G. 2016b. *Guia de aves da Estação Ecológica do Seridó*. Ed. Caule de Papiro, Natal.
- Pinto, O. M. O. 1938. Catálogo das aves do Brasil e lista dos exemplares que as representam no Museu Paulista, 1ª parte. *Rev. Mus. Paulista* 22: 1–566.
- Pinto, O. M. O. 1944. *Catálogo das aves do Brasil e lista dos exemplares existentes na coleção do departamento de zoologia*, 2ª parte. Secretária da Agricultura, São Paulo.
- Pinto, O. M. O. 1978. *Novo catálogo das aves do Brasil*, 1ª parte. Empresa Gráfica Revista dos Tribunais, São Paulo.
- Pinto, O. M. O. & Camargo, E. A. 1961. Resultados ornitológicos de quatro recentes expedições do Departamento de Zoologia ao nordeste do Brasil, com a descrição de seis novas subespécies. *Arq. Zool. São Paulo* 11: 193–284.
- Praxedes, G. C., Araújo, M. F. F. & Varella-Freire, A. A. 1997. Zoogeografia do Rio Grande do Norte II: levantamento preliminar da ornitofauna da micro-região da Serra Verde (Taipu, RN). *Cadernos Northerio-grandenses Temas Geogr.* 11: 82–89.
- Queiroz, L. P., Cardoso, D., Fernandes, M. F. & Moro, M. F. 2017. Diversity and evolution of flowering plants of the Caatinga domain. Pp. 23–63 in Silva, J. M. C., Leal, I. R. & Tabarelli, M. (eds.) *Caatinga: the largest tropical dry forest region in South America*. Springer, Berlin.
- Ramírez, I., Paiva, V. H., Menezes, D., Silva, I., Phillips, R. A., Ramos, J. Á. & Garthe, S. 2013. Year-round distribution and habitat preferences of the Bugio petrel. *Mar. Ecol. Progr. Ser.* 476: 269–284.
- Ramos, R., Sanz, V., Militão, T., Bried, J., Neves, V. C., Biscoito, M., Phillips, R. A., Zino, F. & González-Solís, J. 2015. Leapfrog migration and habitat preferences of a small oceanic seabird, Bulwer's petrel (*Bulweria bulwerii*). *J. Biogeogr.* 42: 1651–1664.
- Ramos, R., Ramírez, I., Paiva, V. H., Militão, T., Biscoito, M., Menezes, D., Phillips, R. A., Zino F. & González-Solís, J. 2016. Global spatial ecology of three closely-related gadfly petrels. *Sci. Rep.* 6: 23447. <https://dx.doi.org/10.1038/srep23447>.
- Ridgely, R. S. & Tudor, G. 2009. *Field guide to the songbirds of South America: the passerines*. Univ. of Texas Press, Austin.
- Robertson, B. C., Stephenson, B. M., Ronconi, R. A., Goldstien, S. J., Shepherd, L., Tennyson, A., Carlile, N. & Ryan, P. G. 2016. Phylogenetic affinities of the *Fregetta* storm-petrels are not black and white. *Mol. Phyl. & Evol.* 97: 170–176.
- Rocha-Neto, M., Santos, M. P., Batista, M. G. & Duarte, M. A. G. 1998. Lista preliminar da avifauna da Escola das Dunas de Pitangui, município de Extremoz, Rio Grande do Norte. *Paradigmas* 11: 107–110.
- Roda, S. A. & Pereira, G. A. 2006. Distribuição recente e conservação das aves de rapina florestais do Centro Pernambuco. *Rev. Bras. Orn.* 14: 331–344.
- Santos, K. B. 2017. Efeitos da pluviosidade e disponibilidade de alimento na reprodução de aves em uma floresta seca neotropical. M.Sc. thesis. Univ. Federal Rural do Semi-árido, Mossoró.
- Schulz-Neto, A. 1998. Aspectos biológicos da avifauna marinha na Reserva Biológica do Atol das Rocas, Rio Grande do Norte, Brasil. *Hornero* 15: 17–28.

- Schulz-Neto, A. 2004a. Aves insulares do arquipélago de Fernando de Noronha. Pp. 147–168 in Branco, J. O. (ed.) *Aves marinhas e insulares brasileiras: bioecologia e conservação*. Ed. UNIVALI, Itajaí.
- Schulz-Neto, A. 2004b. Aves marinhas do Atol das Rocas. Pp. 169–192 in Branco, J. O. (ed.) *Aves marinhas e insulares brasileiras: bioecologia e conservação*. Ed. UNIVALI, Itajaí.
- Sick, H. 1972. A ameaça da avifauna brasileira. Pp. 99–153 in Academia Brasileira de Ciências (ed.) *Espécies da fauna brasileira ameaçadas de extinção*. Academia Brasileira de Ciências, Rio de Janeiro.
- Sick, H. 1985. *Ornitologia brasileira: uma introdução*. Ed. Univ. de Brasília, Brasília.
- Sick, H. 1991. Distribution and subspeciation of the Biscutate Swift *Streptoprocne biscutata*. *Bull. Brit. Orn. Cl.* 111: 38–40.
- Sick, H. 1993. *Birds in Brazil*. Princeton Univ. Press.
- Sick, H. 1997. *Ornitologia brasileira*. Ed. Nova Fronteira, Rio de Janeiro.
- Sick, H., Andrade, M. A., Mattos, G. T. & Freitas, M. V. 1988. Anilhamento de *Streptoprocne biscutata* no Rio Grande do Norte. Pp. 65–72 in Sander, M. (ed.) *An. III Encontro Nac. Anilhadores Aves*. Univ. do Vale do Rio dos Sinos, São Leopoldo.
- Silva, A. C. 2003. As representações zoomórficas na subtradição Seridó. M.Sc. thesis. Univ. Federal de Pernambuco, Recife.
- Silva, C. C. O., Pichorim, M., Moura, P. T. S. & França, L. F. 2017a. Seasonality, abundance and detection bias of birds in a tropical dry forest in north-eastern South America. *J. Trop. Ecol.* 33: 365–378.
- Silva, J. M. C., Barbosa, L. C. F., Leal, I. R. & Tabarelli, M. 2017b. The Caatinga: understanding the challenges. Pp. 3–19 in Silva, J. M. C., Leal, I. R. & Tabarelli, M. (eds.) *Caatinga: the largest tropical dry forest region in South America*. Springer, Berlin.
- Silva, J. M. C. & Castelletti, C. H. M. 2003. Status of the biodiversity of the Atlantic Forest of Brazil. Pp. 43–59 in Galindo-Leal, C. & Câmara, I. G. (eds.) *The Atlantic Forest of South America: biodiversity status, threats, and outlook*. Center for Applied Biodiversity Science and Island Press, Washington DC.
- Silva, M. 2013. Taxonomia e biogeografia da espécie politépica *Herpsilochmus rufimarginatus* (Temminck, 1822) (Aves: Thamnophilidae). M.Sc. thesis. Univ. Federal do Rio Grande do Norte, Natal.
- Silva, M., França, B. R. A., Hagi, L. Y. G. L., Rocha Neto, M., Oliveira, D. V. & Pichorim, M. 2011. New sites and range extensions for endemic and endangered birds in extreme north-east Brazil. *Bull. Brit. Orn. Cl.* 131: 234–240.
- Silva, M., França, B. R. A., Irusta, J. B., Souto, G. H. B. O., Oliveira-Júnior, T. M., Rodrigues, M. C. & Pichorim, M. 2012. Aves de treze áreas de caatinga no Rio Grande do Norte, Brasil. *Rev. Bras. Orn.* 20: 312–328.
- Silva e Silva, R. 1996. Records and geographical distribution of the Peregrine Falcon *Falco peregrinus* Tunstall, 1771 (Aves, Falconidae) in Brazil. *Pap. Avuls. Zool., São Paulo* 39: 249–270.
- Silva e Silva, R. 2008. *Aves de Fernando de Noronha / Birds of Fernando de Noronha*. Ed. Avis Brasilis, Vinhedo.
- Silveira, A. G. & Varela-Freire, A. A. 1999. Zoogeografia do Rio Grande do Norte V: resultado preliminar da composição da avifauna do município de Major Sales, Rio Grande do Norte. *Sociedade e Território* 13: 22–30.
- Silveira, A. G., Santos, M. P. D., Batista, M. A. G., Duarte, M. A. G. & Rocha-Neto, M. 2001. As aves e sua diversidade de habitats na Escola das Dunas de Pitangui. Pp. 59–69 in Rocha-Neto M (ed.) *Guia ilustrado: fauna da escola das Dunas de Pitangui, ecossistema terrestre*. Moura Ramos Gráfica e Editora, Natal.
- Silveira, L. F. 2008. *Tangara fastuosa* (Lesson, 1831). Pp. 554–555 in Machado, A. B. M., Drummond, G. M. & Paglia, A. (eds.) *Livro vermelho da fauna brasileira ameaçada de extinção*. Ministério do Meio Ambiente e Fundação Biodiversitas, Brasília.
- Silveira, L. F., Olmos, F. & Long, A. J. 2003. Birds in Atlantic Forest fragments in northeast Brazil. *Cotinga* 20: 32–46.
- Sistema Nacional de Anilhamento de Aves Silvestres (SNA). 2019. Banco de dados do Centro Nacional de Pesquisa e Conservação de Aves Silvestres (CEMAVE) / Instituto Chico Mendes da Biodiversidade (ICMBio). <http://www.icmbio.gov.br/cemave/sna> (accessed July 2019).
- Snow, D. & Kirwan, G. M. 2019. Amazonian Motmot (*Momotus momota*). In del Hoyo, J., Elliott, A., Sargatal, J., Christie, D. A. & de Juana, E. (eds.) *Handbook of the birds of the world Alive*. Lynx Edicions, Barcelona (retrieved from <https://www.hbw.com/node/55826> in June 2019).
- Somenzari, M., Amara, P. P., Cueto, V. R., Guaraldo, A. C., Jahn, A. E., Lima, D. M., Lima, P. C., Lugarini, C., Machado, C. G., Martinez, J., Nascimento, J. L. X., Pacheco, J. F., Paludo, D., Prestes, N. P., Serafini, P. P., Silveira, L. F., Sousa, A. E. B. A., Sousa, N. A., Souza, M. A., Telino-Júnior, W. R. & Whitney, B. M. 2018. An overview of migratory birds in Brazil. *Pap. Avuls. Zool., São Paulo* 58: 1–66.
- Souza, M. S. & Medeiros, O. 1982. *Inscrições rupestres no Rio Grande do Norte*. Coleção Textos Acadêmicos da Univ. Federal do Rio Grande do Norte, Natal.
- Tabarelli, M. & Santos, A. M. M. 2004. Uma breve descrição sobre a história natural dos brejos nordestinos. Pp. 17–24 in Porto, K. C., Cabral, J. J. P. & Tabarelli, M. (eds.) *Brejos de altitude em Pernambuco e Paraíba: história natural, ecologia e conservação*. Ministério do Meio Ambiente, Brasília.
- Teixeira, D. M. 1986. The avifauna of the north-eastern Brazilian Atlantic Forest: a case of mass extinction? *Ibis* 128: 167–168.
- Teixeira, D. M. 1987. A new tyrannulet (*Phylloscartes*) from northeastern Brazil. *Bull. Brit. Orn. Cl.* 107: 37–41.

- Teixeira, D. M. 1992. As fontes do paraíso – um ensaio sobre a ornitologia no Brasil Holandês (1624-1654). *Rev. Nordestina Biol.* 7(1/2): 1–149.
- Teixeira, D. M., Otoch, R., Luigi, G., Raposo, M. A. & Almeida, A. C. C. 1993. Notes on some birds of northeastern Brazil (5). *Bull. Brit. Orn. Cl.* 113: 48–52.
- Teixeira, R. S. C., Otoch, R. & Raposo, M. A. 2016. First record of Northern Gannet *Morus bassanus* in the Southern Hemisphere. *Bull. Brit. Orn. Cl.* 136: 151–152.
- Toledo-Lima, G. S. T. 2013. Sobrevivência de adultos e estimativa populacional da guaracava-de-topete-uniforme *Elaenia cristata* (Aves: Tyrannidae) em fragmentos de mata atlântica no nordeste do Brasil. M.Sc. thesis. Univ. Federal do Rio Grande do Norte, Natal.
- Tomotani, B. M. & Silveira, L. F. 2016. A reassessment of the taxonomy of *Crypturellus noctivagus* (Wied, 1820). *Rev. Bras. Orn.* 24: 34–45.
- Traylor, M. A. 1958. Variation in South American Great Horned Owls. *Auk* 75: 143–149.
- Varela-Freire, A. A. (ed.). 1997. *Fauna Potiguar V.1: a fauna das dunas costeiras da cidade do Natal e do litoral oriental do Rio Grande do Norte*. Ed. Univ. Federal do Rio Grande do Norte, Natal.
- Varela-Freire, A. A. & Araújo, S. A. 1997. Zoogeografia do Rio Grande do Norte I: a ornitofauna do Seridó Potiguar. *Cadernos Norte-rio-grandenses Temas Geogr.* 11: 52–81.
- Varela-Freire, A. A., Andrade, H. T. A., Mannarino, R. P. & Farias, J. G. B. 1997. Caracterização da fauna terrestre e rizófila de *Eichornia crassipes* (Pontederaceae) nas áreas de influência da Petrobras, Bacia Potiguar, Norte do Rio Grande do Norte, Brasil. Relatório interno de maio 1997. Projeto Petrobras/FUNPEC/Zoologia.
- Varela-Freire, A. A., Silveira, A. G., Souza, D. L., Irusta, J. B., Silva, M., Lima, M. L., Assis, M. O. C. & Maciel, R. A. M. 1999. *Lista atualizada das aves do Estado do Rio Grande do Norte*. Instituto de Desenvolvimento e Meio Ambiente do Rio Grande do Norte, Natal.
- Vargas, O. D. E. 2017. Taxonomia e distribuição geográfica do complexo *Penelope superciliaris* Temminck, 1815 (Aves: Galliformes: Cracidae). M.Sc. thesis. Museu de Zoologia da Universidade de São Paulo, São Paulo.
- Veloso, H. P., Rangel Filho, A. L. R. & Lima, J. C. A. 1991. *Classificação da vegetação brasileira, adaptada a um sistema universal*. Instituto Brasileiro de Geografia e Estatísticas, Rio de Janeiro.
- Werneck, F. P. 2011. The diversification of eastern South American open vegetation biomes: historical biogeography and perspectives. *Quaternary Sci. Rev.* 30: 1630–1648.
- Whitney, B. M., Pacheco, J. F., Buzzetti, D. R. C. & Parrini, R. 2000. Systematic revision and biogeography of the *Herpsilochmus pileatus* complex, with description of a new species from northeastern Brazil. *Auk* 117: 869–891.
- WikiAves. 2019. A enciclopédia das aves do Brasil. <http://www.wikiaves.com> (accessed May 2019).
- Willis, E. O. & Oniki, Y. 1993. New and reconfirmed birds from the state of São Paulo, Brazil, with notes on disappearing species. *Bull. Brit. Orn. Cl.* 113: 23–34.
- Zino, F., Phillips, R. & Bischoff, M. 2011. Zino's Petrel movements at sea: a preliminary analysis of datalogger results. *Birding World* 24: 216–219.

*Addresses:* François Sagot-Martin, Grupo Ornitológico Potiguar, Rua Coronel Costa Pinheiro 1263, Tirol, Natal, RN, 59015-050, Brazil, e-mail: gopforpus@yahoo.com.br. Rafael Dantas Lima, Universidade do Estado do Rio Grande do Norte, Departamento de Ciências Biológicas, Av. Prof. Antônio Campos, s/n, Presidente Costa e Silva, Mossoró, RN, 59625-620, Brazil, e-mail: rafael-dlima@hotmail.com. José Fernando Pacheco, Comitê Brasileiro de Registros Ornitológicos, e-mail: pacheco@oikos.com.br. Jorge Bañuelos Irusta, Irusta Consultoria, Rua Marabá 350, Parnamirim, RN, 59161-230, Brazil, e-mail: jb.irusta@gmail.com. Mauro Pichorim, Universidade Federal do Rio Grande do Norte, Departamento de Botânica e Zoologia, Av. Senador Salgado Filho 3000, Lagoa Nova, Natal, RN, 59078-900, Brazil, e-mail: pichorimmauro@gmail.com. David Maurice Hassett, Santuário Ecológico de Pipa, Tibau do Sul, RN, Brazil, e-mail: davidmhassett@yahoo.com.br

## Appendix 1

Tertiary list: species with published records for Rio Grande do Norte, but with questionable or invalid physical evidence, and whose occurrence in the state appears improbable based on current knowledge. References: R1—Varela-Freire & Araújo (1997); R2—Praxedes *et al.* (1997); R3—Rocha-Neto *et al.* (1998); R4—Silveira & Varela-Freire (1999); R5—Varela-Freire *et al.* (1999); R6—Medeiros *et al.* (2000); R7—Silveira *et al.* (2001); R8—Azevedo *et al.* (2003); R9—MMA (2003); R10—Olmos (2003); R11—Azevedo *et al.* (2004); R12—MMA (2004); R13—Silveira (2008); R14—Grantsau (2010); R15—Silva *et al.* (2012); R16—Moreira-Lima (2013); R17—Cavalcanti *et al.* (2016); R18—Pichorim *et al.* (2016b); R19—Santos (2017); R20—ICMBio / MMA (2018); R21—Somenzari *et al.* (2018); R22—Fitzpatrick (2019). Notes below (N1–27).

| Taxon   | English name               | Reference  | Notes |
|---|----------------------------|------------|-------|
| <b>TINAMIFORMES</b>   |                            |            |       |
| <b>Tinamidae</b>  |                            |            |       |
| <i>Crypturellus erythropus</i> (Pelzeln, 1863)              | Red-legged Tinamou         | R4         | N1    |
| <b>ANSERIFORMES</b>   |                            |            |       |
| <b>Anatidae</b>   |                            |            |       |
| <i>Anas platalea</i> Vieillot, 1816                         | Red Shoveler               | R4         | N1    |
| <b>PELECANIFORMES</b>                                       |                            |            |       |
| <b>Threskiornithidae</b>                                    |                            |            |       |
| <i>Mesembrinibis cayennensis</i> (J. F. Gmelin, 1789)       | Green Ibis                 | R12, R15   | N2    |
| <b>ACCIPITRIFORMES</b>                                      |                            |            |       |
| <b>Accipitridae</b>   |                            |            |       |
| <i>Harpagus bidentatus</i> (Latham, 1790)                   | Double-toothed Kite        | R6         | N3    |
| <b>CHARADRIIFORMES</b>                                      |                            |            |       |
| <b>Charadriidae</b>   |                            |            |       |
| <i>Charadrius melodus</i> Ord, 1824                         | Piping Plover              | R8, R11    | N4    |
| <b>Sternidae</b>  |                            |            |       |
| <i>Sterna hirundinacea</i> Lesson, 1831                     | South American Tern        | R5         | N5    |
| <i>Thalasseus maximus</i> (Boddaert, 1783)                  | Royal Tern                 | R9         | N6    |
| <b>APODIFORMES</b>  |                            |            |       |
| <b>Trochilidae</b>  |                            |            |       |
| <i>Discosura longicaudus</i> (J. F. Gmelin, 1788)           | Racket-tailed Coquette     | R16        | N7    |
| <i>Amazilia lactea</i> (Lesson, 1832)                       | Sapphire-spangled Emerald  | R3, R7     | N8    |
| <b>GALBULIFORMES</b>  |                            |            |       |
| <b>Galbulidae</b>   |                            |            |       |
| <i>Nystalus chacuru</i> (Vieillot, 1816)                    | White-eared Puffbird       | R4, R5     | N9    |
| <b>PICIFORMES</b>   |                            |            |       |
| <b>Picidae</b>  |                            |            |       |
| <i>Picus flavigula</i> (Boddaert, 1783)                     | Yellow-throated Woodpecker | R2, R5, R6 | N10   |
| <i>Celeus flavus</i> (Statius Muller, 1776)                 | Cream-coloured Woodpecker  | R5         | N5    |
| <b>PASSERIFORMES</b>  |                            |            |       |
| <b>Thamnophilidae</b>                                       |                            |            |       |
| <i>Thamnophilus palliatus</i> (M. H. C. Lichtenstein, 1823) | Chestnut-backed Antshrike  | R5         | N5    |
| <i>Thamnophilus caeruleus</i> Vieillot, 1816                | Variable Antshrike         | R3, R7     | N11   |
| <i>Batara cinerea</i> (Vieillot, 1819)                      | Giant Antshrike            | R5         | N5    |
| <i>Myrmoderus loricatus</i> (M. H. C. Lichtenstein, 1823)   | White-bibbed Antbird       | R3, R7     | N12   |
| <b>Dendrocolaptidae</b>                                     |                            |            |       |



| Taxon  | English name                | Reference          | Notes    |
|--|-----------------------------|--------------------|----------|
| <i>Xiphocolaptes falcirostris</i> (Spix, 1824)         | Moustached Woodcreeper      | R20                | N13      |
| <b>Furnariidae</b>                                     |                             |                    |          |
| <i>Furnarius rufus</i> (J. F. Gmelin, 1788)            | Rufous Hornero              | R2, R4, R5, R6, R7 | N14      |
| <i>Phacellodomus ruber</i> (Vieillot, 1817)            | Greater Thornbird           | R5                 | N5       |
| <i>Synallaxis spixi</i> P. L. Sclater, 1856            | Spix's Spinetail            | R6                 | N15      |
| <i>Cranioleuca pallida</i> (Wied, 1831)                | Pallid Spinetail            | R5                 | N5       |
| <b>Onychorhynchidae</b>                                |                             |                    |          |
| <i>Myiobius atricaudus</i> Lawrence, 1863              | Black-tailed Flycatcher     | R10                | N16      |
| <b>Tyrannidae</b>                                      |                             |                    |          |
| <i>Elaenia parvirostris</i> Pelzel, 1868               | Small-billed Elaenia        | R17                | N17      |
| <i>Phyllomyias fasciatus</i> (Thunberg, 1822)          | Planalto Tyrannulet         | R22                | N18      |
| <i>Legatus leucophaeus</i> (Vieillot, 1818)            | Piratic Flycatcher          | R12, R18           | N20      |
| <i>Casiornis rufus</i> (Vieillot, 1816)                | Rufous Casiornis            | R19                | N19      |
| <i>Philohydor lictor</i> (M. H. C. Lichtenstein, 1823) | Lesser Kiskadee             | R4                 | N1       |
| <i>Myiozetetes cayanensis</i> (Linnaeus, 1766)         | Rusty-margined Flycatcher   | R5, R7             | N5, N21  |
| <i>Tyrannus albogularis</i> Burmeister, 1856           | White-throated Kingbird     | R21                | N22      |
| <i>Knipolegus cyanirostris</i> (Vieillot, 1818)        | Blue-billed Black Tyrant    | R1, R5             | N23      |
| <b>Turdidae</b>  |                             |                    |          |
| <i>Turdus fumigatus</i> M. H. C. Lichtenstein, 1823    | Cocoa Thrush                | R4, R5             | N1, N5   |
| <b>Icteridae</b>                                       |                             |                    |          |
| <i>Anumara forbesi</i> (P. L. Sclater, 1886)           | Forbes's Blackbird          | R1, R2, R5, R20    | N13, N24 |
| <b>Thraupidae</b>                                      |                             |                    |          |
| <i>Paroaria coronata</i> (J. F. Miller, 1776)          | Red-crested Cardinal        | R2                 | N25      |
| <i>Tangara fastuosa</i> (Lesson, 1831)                 | Seven-coloured Tanager      | R13, R14, R20      | N26      |
| <i>Tangara episcopus</i> (Linnaeus, 1766)              | Blue-grey Tanager           | R2                 | N25      |
| <i>Sicalis columbiana</i> Cabanis, 1851                | Orange-fronted Yellow Finch | R7                 | N27      |
| <i>Sporophila caerulea</i> (Vieillot, 1823)            | Double-collared Seedeater   | R2                 | N25      |

N1—Undocumented records in Major Sales (Silveira & Varela-Freire 1999), based on information from a local person, using Frisch (1981), are not consistent with the known distribution based on documentary evidence.

N2—The undocumented record of *Mesembrinibis cayennensis*, based on a sight record near the headquarters of the ESEC Seridó, Serra Negra do Norte (MMA 2004), is not consistent with the known distribution based on available documentation. This record supported the species' inclusion in Silva *et al.* (2012). Following Pichorim *et al.* (2016b), we consider it to be a misidentification, and believe the species involved was more likely to be Limpkin *Aramus guarauna* (FSM pers. obs.).

N3—An undocumented record of *Harpagus bidentatus*, referred to as 'ripina', from Riacho do Roçado dam, São José do Seridó, sometime in July–December 1996 (Medeiros *et al.* 2000), is probably a misidentification. The known distribution indicates that the species is not expected in the Caatinga, especially during the long dry season, and, in Rio Grande do Norte, 'ripina' is the vernacular name for Roadside Hawk *Rupornis magnirostris*.

N4—Claimed records of *Charadrius melodus* at Salina Diamante Branco, Galinhos, in October and December 2000 (Azevedo *et al.* 2003, 2004) supported this species' inclusion in the Brazilian primary list (CBRO 2009,

2010, 2011, 2014), but they are misidentifications (Piacentini *et al.* 2015). The voucher UFPE 3656 is, in fact, a *C. semipalmatus* (L. Naka pers. obs.).

N5—Inclusion of this species in the first state bird list (Varela-Freire *et al.* 1999) is inconsistent with the known distribution.

N6—Rio Grande do Norte was included in a list of Brazilian states where *Thalasseus maximus* occurs (MMA 2003), but we are unable to find any evidence for this, and the species is neither cited nor mapped in the state in the most recent Brazilian Red List (ICMBio / MMA 2018). See Appendix 2.

N7—Indication of *Discosura longicaudus* as 'Resident in the Atlantic Forest between Rio Grande do Norte and Rio de Janeiro' (Moreira-Lima 2013) appears speculative, and no evidence for its presence in the state is known.

N8—Alleged documentation for *Amazilia lactea* in the state (a photo in Silveira *et al.* 2001: 66) is a misidentification for Glittering-throated Emerald *A. fimbriata* (FSM pers. obs.). *A. lactea* had already been reported, but not documented, at the same locality, Dunas de Pitangui School, Extremoz, between April and December 1996 (Rocha-Neto *et al.* 1998).

N9—Undocumented records of *Nystalus chacuru* in the Agreste, the Caatinga, and coastal dunes (Varela-Freire *et al.* 1999), are inconsistent with the known distribution. *N. chacuru* had been included in the bird list of Major Sales (Silveira & Varela-Freire 1999), but the first author subsequently confirmed that the species involved was Spot-backed Puffbird *N. maculatus* (A. G. da Silveira pers. comm.).

N10—Undocumented records of *Piculus flavigula* in the Caatinga (Praxedes *et al.* 1997, Varela-Freire *et al.* 1999, Medeiros *et al.* 2000) are inconsistent with the species' known distribution, and probably reflect confusion with Golden-green Woodpecker *P. chryschloros*.

N11—Alleged documentation for *Thamnophilus caerulescens* in the state (a photo in Silveira *et al.* 2001: 69) is a misidentification of female White-fringed Antwren *Formicivora grisea* (FSM pers. obs.). *T. caerulescens* had already been reported, but not documented, at the same locality, Dunas de Pitangui School, Extremoz, between April and December 1996 (Rocha-Neto *et al.* 1998).

N12—An undocumented record of *Myrmoderus loricatus* at Dunas de Pitangui School, Extremoz, sometime during April–December 1996 (Rocha-Neto *et al.* 1998), is inconsistent with the known distribution. It was omitted from a subsequent bird list (Silveira *et al.* 2001).

N13—Indication of occurrence in the state (ICMBio / MMA 2018) is not supported by any available evidence.

N14—Undocumented records of *Furnarius rufus* in the state (Praxedes *et al.* 1997, Silveira & Varela-Freire 1999, Varela-Freire *et al.* 1999, Medeiros *et al.* 2000, Silveira *et al.* 2001), some based on observations of nests (which in fact resemble those of Pale-legged Hornero *F. leucopus*), are inconsistent with the species' known distribution.

N15—An undocumented record of *Synallaxis spixi* at Riacho do Roçado Dam, São José do Seridó, during July–December 1996 (Medeiros *et al.* 2000), is inconsistent with the species' known distribution.

N16—The record of *Myiobius atricaudus* at Mata da Estrela Private Reserve, Baía Formosa (Olmos 2003), is treated here as a misidentification of *M. barbatus* (see Species accounts). See Appendix 2.

N17—*Elaenia parvirostris* was mentioned in a bird list from Mossoró (Cavalcanti *et al.* 2016) without details; the identification must be considered questionable.

N18—Indication for the presence of *Phyllomyias fasciatus* in the state (Fitzpatrick 2019) cannot be confirmed; see Appendix 2.

N19—*Casiornis rufus* was mentioned in a bird list from Mossoró (Santos 2017) without details; the identification must be considered questionable.

N20—An undocumented record of *Legatus leucophaeus* at ESEC Seridó, Serra Negra do Norte (MMA 2004), supported its inclusion in the locality's bird list (Pichorim *et al.* 2016b), but to us it seems a probable misidentification for Variegated Flycatcher *Empidonomus varius*. Nevertheless, *Legatus* is likely to occur in Atlantic Forest remnants in the south-east of the state, but not the Caatinga (see Appendix 2).

N21—A photo (Fig. 9.8) in Silveira *et al.* (2001), taken at Dunas de Pitangui School, Extremoz, and misidentified as *Myiozetetes cayanensis*, is, in fact, of Social Flycatcher *M. similis* (FSM pers. obs.).

N22—Visual identification of *Tyrannus albogularis*, which in many respects resembles some Tropical Kingbirds *T. melancholicus*, can be difficult. Somenzari *et al.* (2018) noted that, based on WikiAves records, this species breeds in Brazil during October–April (including Rio Grande do Norte), but we believe that all

photos from the state are misidentified *T. melancholicus*, and that *T. albogularis* does not breed in far north-east Brazil (see Appendix 2).

N23—A record of *Knipolegus cyanirostris* in the Seridó (Varela-Freire & Araújo 1997) supported its inclusion in the state bird list (Varela-Freire *et al.* 1999), but is inconsistent with the species’ known distribution. More likely the observation involved *K. nigerrimus*.

N24—The Vulnerable Brazilian endemic *Anumara forbesi* (ICMBio / MMA 2018), much sought after in the cagebird trade, has been reported in the state without details (Varela-Freire & Araújo 1997, Praxedes *et al.* 1997, Varela-Freire *et al.* 1999), but all reports are of questionable identification and origin.

N25—An undocumented record in the Agreste of Taipu (Praxedes *et al.* 1997) is inconsistent with the known distribution. It is also of questionable identification, and it may have involved a cagebird.

N26—*Tangara fastuosa* has been mapped for the state (ICMBio / MMA 2018) based on Silveira (2008) and Grantsau (2010). A previous Brazilian Red List (MMA 2003) also included Rio Grande do Norte within this tanager’s range, probably based on Silveira & Varela-Freire (1999), who reported it to be rare and threatened, and that it had been found in ‘coastal sand dunes, and tabuleiro of Capim Macio.’ In these locations within Dunas de Natal State Park, confiscated cagebirds were released (Dr A. Leite, Cetas / IBAMA Natal, pers. comm.). Olmos (2003) did not detect the species in Mata da Estrela Private Reserve and environs, Baía Formosa, during 28 February to 7 March 2003. Neither did JFP, accompanied by FSM, find the species in south-east Rio Grande do Norte between 26 December 2010 and 3 January 2011. Over the last ten years, many hundreds of birds have been trapped by MP’s students in the *restinga* of Natal and Parnamirim, and the semi-deciduous forest of Mata da Estrela, but no *T. fastuosa*. Two sight records, both of singles in flight beside the Jiqui River, Parnamirim, on 28 December 2006 and on 15 January 2007 (FSM), pertain to releases by environmental agents. If the Vulnerable *T. fastuosa* ever occurred in the state, its population became extinct long ago, and no feral population apparently exists today. Following our criteria, this species is included in the tertiary list.

N27—A record of *Sicalis columbiana* from Dunas de Pitangui School, Extremoz (Silveira *et al.* 2001), is inconsistent with the known distribution. It is also of questionable identification, and may have involved a cagebird.

Appendix 2

Potential List: species with some probability of occurrence in Rio Grande do Norte, including its continental waters, based on known patterns of distribution. Notes: (1) The species has been recorded nearby, in a neighbouring state, and we believe that suitable habitat exists in Rio Grande do Norte; (2) the species has been recorded nearby, in a neighbouring state, but we consider that there is little or no suitable habitat in Rio Grande do Norte; (3) the species has been recorded in a neighbouring state and we consider that suitable habitat exists for the species in Rio Grande do Norte; (4) the known distribution and / or migration of the species is consistent with its possible / probable occurrence in the state; and (5) the species has probably already occurred in the state, but is currently possibly extinct there. Full notes, including references to nearby records, are presented in Supplementary Material.

| Taxon  | English Name           | Probability of occurrence | Notes |
|--|------------------------|---------------------------|-------|
| <i>Crypturellus soui</i> (Hermann, 1783)         | Little Tinamou         | Medium                    | 1     |
| <i>Anas acuta</i> Linnaeus, 1758                 | Northern Pintail       | Low                       | 4     |
| <i>Anas georgica</i> J. F. Gmelin, 1789          | Yellow-billed Pintail  | Medium                    | 3, 4  |
| <i>Odontophorus capueira</i> (Spix, 1825)        | Spot-winged Wood Quail | Low                       | 1, 5  |
| <i>Pterodroma feae</i> (Salvadori, 1899)         | Cape Verde Petrel      | Medium                    | 4     |
| <i>Pterodroma incerta</i> (Schlegel, 1863)       | Atlantic Petrel        | Medium                    | 4     |
| <i>Pachyptila desolata</i> (J. F. Gmelin, 1789)  | Antarctic Prion        | Medium                    | 4     |
| <i>Pachyptila belcheri</i> (Mathews, 1912)       | Slender-billed Prion   | Medium                    | 4     |
| <i>Procellaria aequinoctialis</i> Linnaeus, 1758 | White-chinned Petrel   | Medium                    | 4     |
| <i>Calonectris edwardsii</i> (Oustalet, 1883)    | Cape Verde Shearwater  | Medium                    | 4     |
| <i>Puffinus lherminieri</i> Lesson, 1839         | Audubon’s Shearwater   | Medium                    | 4     |

| <b>Taxon</b>                                    | <b>English Name</b>        | <b>Probability of occurrence</b> | <b>Notes</b> |
|---|----------------------------|----------------------------------|--------------|
| <i>Fregetta grallaria</i> (Vieillot, 1818)      | White-bellied Storm Petrel | Low                              | 4            |
| <i>Pelagodroma marina</i> (Latham, 1790)        | White-faced Storm Petrel   | Low                              | 4            |
| <i>Oceanodroma castro</i> (Harcourt, 1851)      | Band-rumped Storm Petrel   | Medium                           | 4            |
| <i>Hydrobates pelagicus</i> (Linnaeus, 1758)    | European Storm Petrel      | Medium                           | 4            |
| <i>Phaethon aethereus</i> Linnaeus, 1758        | Red-billed Tropicbird      | Medium                           | 3, 4         |
| <i>Phaethon rubricauda</i> Boddaert, 1783       | Red-tailed Tropicbird      | Low                              | 3, 4         |
| <i>Phaethon lepturus</i> Daudin, 1802           | White-tailed Tropicbird    | High                             | 3, 4         |
| <i>Fregata ariel</i> (G. R. Gray, 1845)         | Lesser Frigatebird         | Low                              | 4            |
| <i>Morus bassanus</i> (Linnaeus, 1758)          | Northern Gannet            | Low                              | 3, 4         |
| <i>Cochlearius cochlearius</i> (Linnaeus, 1766) | Boat-billed Heron          | High                             | 1            |
| <i>Ardeola ralloides</i> (Scopoli, 1769)        | Squacco Heron              | Medium                           | 3, 4         |
| <i>Ardea cinerea</i> Linnaeus, 1758             | Grey Heron                 | Medium                           | 4            |
| <i>Ardea purpurea</i> Linnaeus, 1766            | Purple Heron               | Low                              | 4            |
| <i>Egretta tricolor</i> (Statius Muller, 1776)  | Tricoloured Heron          | Low                              | 3, 4         |
| <i>Egretta gularis</i> (Bosc, 1792)             | Western Reef Heron         | Medium                           | 4            |
| <i>Egretta garzetta</i> (Linnaeus, 1766)        | Little Egret               | Low                              | 4            |
| <i>Platalea leucorodia</i> Linnaeus, 1758       | Eurasian Spoonbill         | Low                              | 4            |
| <i>Harpagus diodon</i> (Temminck, 1823)         | Rufous-thighed Kite        | High                             | 1, 4         |
| <i>Milvus migrans</i> (Boddaert, 1783)          | Black Kite                 | Low                              | 4            |
| <i>Buteo platypterus</i> (Vieillot, 1823)       | Broad-winged Hawk          | Medium                           | 3, 4         |
| <i>Spizaetus tyrannus</i> (Wied, 1820)          | Black Hawk-Eagle           | High                             | 1            |
| <i>Crex crex</i> (Linnaeus, 1758)               | Corn Crake                 | Low                              | 4            |
| <i>Pardirallus nigricans</i> (Vieillot, 1819)   | Blackish Rail              | Medium                           | 1            |
| <i>Gallinula angulata</i> Sundevall, 1850       | Lesser Moorhen             | Low                              | 4            |
| <i>Bartramia longicauda</i> (Bechstein, 1812)   | Upland Sandpiper           | Low                              | 3, 4         |
| <i>Tringa totanus</i> (Linnaeus, 1758)          | Common Redshank            | Low                              | 4            |
| <i>Calidris ferruginea</i> (Pontoppidan, 1763)  | Curlew Sandpiper           | Low                              | 3, 4         |
| <i>Calidris pugnax</i> (Linnaeus, 1758)         | Ruff                       | Medium                           | 3, 4         |
| <i>Glareola pratincola</i> (Linnaeus, 1766)     | Collared Pratincole        | Low                              | 3, 4         |
| <i>Stercorarius skua</i> (Brünnich, 1764)       | Great Skua                 | Medium                           | 4            |
| <i>Stercorarius antarcticus</i> (Lesson, 1831)  | Brown Skua                 | Low                              | 4            |
| <i>Stercorarius longicaudus</i> Vieillot, 1819  | Long-tailed Jaeger         | Medium                           | 4            |
| <i>Leucophaeus pipixcan</i> (Wagler, 1831)      | Franklin's Gull            | Medium                           | 4            |
| <i>Larus fuscus</i> Linnaeus, 1758              | Lesser Black-backed Gull   | Medium                           | 1, 4         |
| <i>Anous minutus</i> Boie, 1844                 | Black Noddy                | Low                              | 4            |
| <i>Gygis alba</i> (Sparrman, 1786)              | Common White Tern          | Low                              | 4            |

| <b>Taxon</b>   | <b>English Name</b>          | <b>Probability of occurrence</b> | <b>Notes</b> |
|--|------------------------------|----------------------------------|--------------|
| <i>Thalasseus maximus</i> (Boddaert, 1783)                                 | Royal Tern                   | Low                              | 3, 4         |
| <i>Coccyzus minor</i> (J. F. Gmelin, 1788)                                 | Mangrove Cuckoo              | Low                              | 3, 4         |
| <i>Florisuga fusca</i> (Vieillot, 1817)                                    | Black Jacobin                | Medium                           | 1, 4         |
| <i>Chelidoptera tenebrosa</i> (Pallas, 1782)                               | Swallow-winged Puffbird      | Low                              | 1, 5         |
| <i>Picummus pernambucensis</i> J. T. Zimmer, 1947                          | Pernambuco Piculet           | High                             | 1            |
| <i>Falco tinnunculus</i> Linnaeus, 1758                                    | Eurasian Kestrel             | Low                              | 4            |
| <i>Pyrrhura griseipectus</i> Salvadori, 1900                               | Grey-breasted Parakeet       | Low                              | 3, 5         |
| <i>Touit surdus</i> (Kuhl, 1820)   | Golden-tailed Parrotlet      | High                             | 1            |
| <i>Thamnophilus palliatus</i> (M. H. C. Lichtenstein, 1823)                | Chestnut-backed Antshrike    | Medium                           | 1            |
| <i>Myrmoderus ruficauda</i> (Wied, 1831)                                   | Scalloped Antbird            | Low                              | 1, 5         |
| <i>Pyriglena pernambucensis</i> J. T. Zimmer, 1931                         | Pernambuco Fire-eye          | Medium                           | 1, 5         |
| <i>Hyllopezus ochroleucus</i> (Wied, 1831)                                 | White-browed Antpitta        | Low                              | 2            |
| <i>Sclerurus cearensis</i> E. Snethlage, 1924                              | Ceara Leaf-tosser            | Low                              | 2            |
| <i>Xiphorhynchus atlanticus</i> (Cory, 1916)                               | Northern Lesser Woodcreeper  | High                             | 1            |
| <i>Dendrocolaptes platyrostris</i> Spix, 1825                              | Planalto Woodcreeper         | High                             | 1            |
| <i>Xiphocolaptes falcirostris</i> (Spix, 1824)                             | Moustached Woodcreeper       | High                             | 1            |
| <i>Automolus lammi</i> J. T. Zimmer, 1947                                  | Pernambuco Foliage-gleaner   | Medium                           | 1, 5         |
| <i>Megaxenops parnaguae</i> Reiser, 1905                                   | Great Xenops                 | Low                              | 2            |
| <i>Synallaxis hypospodia</i> P. L. Sclater, 1874                           | Cinereous-breasted Spinetail | Low                              | 4            |
| <i>Cranioleuca semicinerea</i> (Reichenbach, 1853)                         | Grey-headed Spinetail        | High                             | 1            |
| <i>Myiobius atricaudus</i> Lawrence, 1863                                  | Black-tailed Flycatcher      | Medium                           | 3            |
| <i>Schiffornis turdina</i> (Wied, 1831)                                    | Thrush-like Schiffornis      | Medium                           | 1, 5         |
| <i>Iodopleura pipra</i> (Lesson, 1831)                                     | Buff-throated Purple-tuft    | High                             | 1            |
| <i>Xipholena atropurpurea</i> (Wied, 1820)                                 | White-winged Cotinga         | Medium                           | 1            |
| <i>Mionectes oleagineus</i> (M. H. C. Lichtenstein, 1823)                  | Ochre-bellied Flycatcher     | Medium                           | 1            |
| <i>Tolmomyias sulphureus</i> (Spix, 1825)                                  | Yellow-olive Flycatcher      | Medium                           | 1            |
| <i>Tolmomyias poliocephalus</i> (Taczanowski, 1884)                        | Grey-crowned Flycatcher      | Medium                           | 1            |
| <i>Hemitriccus mirandae</i> (E. Snethlage, 1925)                           | Buff-breasted Tody-Tyrant    | Low                              | 2            |
| <i>Zimmerius acer</i> (Salvin & Godman, 1883)                              | Guianan Tyrannulet           | High                             | 1            |
| <i>Phyllomyias fasciatus</i> (Thunberg, 1822)                              | Planalto Tyrannulet          | High                             | 1            |
| <i>Legatus leucophaeus</i> (Vieillot, 1818)                                | Piratic Flycatcher           | High                             | 1            |
| <i>Rhytipterna simplex</i> (M. H. C. Lichtenstein, 1823)                   | Greyish Mourner              | Medium                           | 1            |
| <i>Philohydor lictor</i> (M. H. C. Lichtenstein, 1823)                     | Lesser Kiskadee              | Low                              | 1, 5         |
| <i>Tyrannus albogularis</i> Burmeister, 1856                               | White-throated Kingbird      | Low                              | 4            |
| <i>Griseotyrannus aurantioatrocristatus</i> (d'Orbigny & Lafresnaye, 1837) | Crowned Slaty Flycatcher     | Low                              | 3, 4         |
| <i>Contopus virens</i> (Linnaeus, 1766)                                    | Eastern Wood Pewee           | Low                              | 3, 4         |



| Taxon   | English Name                | Probability of occurrence | Notes |
|---|-----------------------------|---------------------------|-------|
| <i>Contopus cinereus</i> (Spix, 1825)                 | Tropical Pewee              | Low                       | 3, 4  |
| <i>Progne subis</i> (Linnaeus, 1758)                  | Purple Martin               | High                      | 1, 4  |
| <i>Riparia riparia</i> (Linnaeus, 1758)               | Bank Swallow                | Medium                    | 4     |
| <i>Catharus fuscescens</i> (Stephens, 1817)           | Veery                       | Low                       | 4     |
| <i>Setophaga fusca</i> (Statius Muller, 1776)         | Blackburnian Warbler        | Low                       | 3, 4  |
| <i>Geothlypis aequinoctialis</i> (J. F. Gmelin, 1789) | Masked Yellowthroat         | Medium                    | 1     |
| <i>Hemithraupis guira</i> (Linnaeus, 1766)            | Guira Tanager               | High                      | 1     |
| <i>Tiaris fuliginosus</i> (Wied, 1830)                | Sooty Grassquit             | Medium                    | 1     |
| <i>Sporophila angolensis</i> (Linnaeus, 1766)         | Chestnut-bellied Seed Finch | Low                       | 1     |

Supplementary Material

The database of the Rio Grande do Norte avifauna, an Excel spreadsheet containing all the lists (primary, secondary, tertiary, and potential) with full details and Brazilian Portuguese names, is available on BioOne.

