

Effects of simulated moose *Alces alces* browsing on the morphology of rowan *Sorbus aucuparia*

Authors: Jager, Nathan R. De, and Pastor, John

Source: Wildlife Biology, 16(3) : 301-307

Published By: Nordic Board for Wildlife Research

URL: <https://doi.org/10.2981/09-105>

BioOne Complete (complete.BioOne.org) is a full-text database of 200 subscribed and open-access titles in the biological, ecological, and environmental sciences published by nonprofit societies, associations, museums, institutions, and presses.

Your use of this PDF, the BioOne Complete website, and all posted and associated content indicates your acceptance of BioOne's Terms of Use, available at www.bioone.org/terms-of-use.

Usage of BioOne Complete content is strictly limited to personal, educational, and non - commercial use. Commercial inquiries or rights and permissions requests should be directed to the individual publisher as copyright holder.

BioOne sees sustainable scholarly publishing as an inherently collaborative enterprise connecting authors, nonprofit publishers, academic institutions, research libraries, and research funders in the common goal of maximizing access to critical research.

Effects of simulated moose *Alces alces* browsing on the morphology of rowan *Sorbus aucuparia*

Nathan R. De Jager & John Pastor

In much of northern Sweden moose *Alces alces* browse rowan *Sorbus aucuparia* heavily and commonly revisit previously browsed plants. Repeated browsing of rowan by moose has created some concern for its long-term survival in heavily browsed areas. We therefore measured how four years of simulated moose browsing at four population densities (0, 10, 30 and 50 moose/1,000 ha) changed plant height, crown width, available bite mass, the number of bites per plant and per plant forage biomass of rowan saplings. Increased biomass removal led to a significant decline in plant height ($P < 0.001$), but a significant increase in the number of bites per plant ($P = 0.012$). Increases in the number of bites per plant more than compensated for weak decreases in bite mass, leading to a weak increase in per plant forage biomass ($P = 0.072$). With the decline in plant height and increase in the number of stems per plant, a greater number of bites remain within the height reach of moose relative to unbrowsed controls. Moose therefore stand to benefit from revisiting previously browsed plants, which may result in feeding loops between moose and previously browsed rowan saplings.

Key words: *Alces alces*, bite mass, forage biomass, herbivory, moose, plant morphology, rowan, *Sorbus aucuparia*

Nathan R. De Jager*, Department of Ecology, Evolution, and Behavior, University of Minnesota, St. Paul, Minnesota 55108, USA - e-mail: ndejager@usgs.gov

John Pastor, Department of Biology, University of Minnesota Duluth, Minnesota 55811, USA - e-mail: jpastor@umn.edu

*Present address: US Geological Survey, Upper Midwest Environmental Sciences Center, La Crosse, Wisconsin 54603, USA

Corresponding author: Nathan R. De Jager

Received 30 November 2009, accepted 11 May 2010

Associate Editor: Christophe Bonenfant

Different woody plant species demonstrate a variety of growth responses to increasing browsing intensity by large herbivores (McInnes et al. 1992, Edenius et al. 1993, Danell et al. 1994, 1997, Persson et al. 2005, De Jager & Pastor 2008, De Jager et al. 2009). Some species may compensate for lost tissue due to browsing, whereas others may not. Plants that do compensate for lost tissue often do so at the cost of height growth, which keeps them within easy reach of herbivores and leads to even more browsing in the future (Pastor & Danell 2003). Over time, highly palatable plants that are repeatedly browsed are often overtopped and subsequently replaced with less palatable plants, thereby changing patterns of plant community composition (Krefting

1974, Risenhoover & Maass 1987, McInnes et al. 1992, Thompson et al. 1992, De Jager & Pastor 2009), soil fertility (McInnes et al. 1992, Pastor et al. 1993, 1998, De Jager & Pastor 2009) and in some cases decreasing biodiversity (van Wieren & Bakker 2008).

Among the characteristics on which moose *Alces alces* select plants are available bite size, bite density and forage biomass (Belovsky 1978, Renecker & Hudson 1986, Vivås & Sæther 1987, Shipley & Spalinger 1995, Shipley et al. 1998, 1999, Shipley 2007). Tree height and crown width can influence rates of browsing by moose and also control plant survival, forest succession and biodiversity. However, if moose browsing influences tree height, then

it can determine whether plants remain within reach and therefore are available for browsing in the future. Thus, the changes in the mass and density of bites that regrow in response to browsing and changes in tree height and crown width ought to subsequently affect the future forage supply to moose and their preference for previously browsed plants.

Although there has been much work on the effects of moose on these properties for many tree species within the boreal biome (see the above references), we are not aware of any study that has systematically examined the response of these morphological features of rowan *Sorbus aucuparia* to a year-round gradient of known moose browsing. Ungulate browsing is suggested to be the most important biotic factor in limiting the growth of rowan in northern Europe (Raspé et al. 2000). Moose find rowan highly palatable, presumably due to its high nutrient content (Heroldova et al. 2003) and because rowan offers moose relatively large and nutritious bites within easy reach (Shipley et al. 1999). Although rowan can be the most abundant species in the seedling cohort in productive forests of northern Sweden, it is often almost entirely absent from the tree layer in highly browsed areas (Linder et al. 1997, Motta 2003). This has created some concern for the long-term survival of rowan in heavily browsed areas (Moen 2004, Mysterud 2006).

In our study, we measured the responses of available bite mass, number of bites per plant, forage biomass, plant height and the width of plant crowns of rowan following four years of four different simulated intensities of moose browsing (0, 10, 30 and 50 moose/1,000 ha) and at two sites of high productivity in northern Sweden. Our previous studies of the responses of birch *Betula pubescens* and *B. pendula* and Scots pine *Pinus sylvestris* in this same experimental setting suggest that the deciduous species have a greater ability to replace lost tissue than conifers (Persson et al. 2005, De Jager & Pastor 2008). Despite declines in plant height and crown width, forage production of birch saplings were either unaffected at the '10 moose' level of tissue removal or responded unimodally with initial increases in forage production at the '10 and 30 moose' levels, followed by declines in forage production at the '50 moose' levels. We therefore hypothesize that increasing moose population density will cause monotonic declines in sapling height and crown width of rowan, but that forage

biomass will compensate for lost tissue due to simulated moose browsing. Such a growth response would suggest that rowan has a high capacity to replace lost tissue due to moose browsing and that previously browsed saplings may become even more palatable to moose relative to unbrowsed controls. On the other hand, if increased population density results in monotonic declines in forage biomass, we will conclude that rowan has a low tolerance for browsing and that previously browsed plants are less palatable to moose than unbrowsed control trees.

Material and methods

We simulated the interactive effects of moose population density and site productivity on plant and ecosystem responses in eight exclosures measuring 70 × 70 m, and were situated 50–90 km north and northwest of Umeå, Sweden (63°50'N, 20°18'E; Persson et al. 2005). The exclosures were constructed in young forest stands of Scots pine planted following clear-cutting approximately 11 years previously and interspersed with naturally reproducing deciduous trees, mainly birch, but also rowan, aspen *Populus tremula* and willows *Salix* spp. Sites were selected along a forest productivity gradient spanning the range of productivities for the region (Hägglund & Lundmark 1987, Fridman et al. 2001). In our study, we examined data from two of the most productive sites, Mörtsjöstavaren and Åtmyrberget, because these were the only sites where rowan was common. These sites had comparable estimates of site productivity and litterfall (Persson et al. 2005), and are typical of the types of plant communities and fertile soil types where rowan is commonly found (Linder et al. 1997).

Within each exclosure, browsing at four moose population densities (0, 10, 30 and 50 moose/1,000 ha) was simulated in 25 × 25-m plots beginning in early spring 1999. A full description of the methods used to simulate moose foraging is given in Persson et al. (2005) and is briefly summarized here. We calculated the consumption per month in the treatment plots for the different moose densities based on an estimated daily food intake of moose of 5 kg dry mass in winter and 10 kg in summer (Persson et al. 2000). The species composition of food plants in each monthly harvest was based on the seasonal composition of a moose's diet (Cederlund et al. 1980, Bergström & Hjelt 1987, Shipley

et al. 1998). During winter (November–April) living twigs > 50 cm above ground were clipped at the mean diameter observed to be browsed by moose in the area, which was 4 mm for rowan and corresponding diameters for other species encountered such as birch, pine, aspen and willow. The mass of all removed plant material was measured each time to 0.1 g precision and then deposited outside the study area. Dung (natural) and urine (artificially made by dissolving urea in water) corresponding to the amounts expected from the simulated moose densities were also added to the treatment plots (Persson et al. 2000).

During early June 2003, after four years of simulated browsing, we measured the number of bites that would be available to moose per tree, the available bite mass, forage biomass per tree, tree height and crown width on the nearest rowan trees > 50 cm in height at 5 m intervals along transects 5 m apart until a total of nine trees were measured in each treatment plot. Tree height was estimated from the ground to the tallest shoot on the plant and crown width was the distance measured across the widest portion of the tree canopy. Available bite mass (g/bite) was measured by selecting all of the twigs on the tree and measuring twig length from the tip to the mean bite diameter observed to be browsed in the areas, which is 4 mm for rowan (Persson et al. 2005). If the diameter of a twig was < 4 mm, the length of the twig was measured from the tip to the point where it branched into a second twig large enough to be considered a distinct bite (> 10 cm in length or about the amount one can fit in the hand when mimicking the act of a moose biting a twig) and both the diameter and length were recorded.

We then developed regression equations to determine the dry weight of rowan bites from basal diameters and lengths ($\ln \text{bite mass} = -3.379 + 0.537 (\ln \text{length}) + 1.714 (\ln \text{diameter})$; $R^2 = 0.97$). Estimates made using twig basal diameter and length yielded the average bite mass available to moose during winter. The total potential number of bites per tree was the total number of these shoots on a tree, and forage biomass was estimated by adding together the dry mass of each shoot on the tree.

We pooled the data from both sites because they had comparable estimates of litterfall and site productivity (Persson et al. 2005) and because there were only two of them. We then tested the effects of the increasing moose browsing treatments on plant height, crown width, bite mass, number of bites per

plant and forage biomass using ANOVA and post-hoc Tukey's *t*-tests. For ANOVAs that were not significant ($P > 0.05$), we also tested for differences between the trees in the control plots (without moose) and trees from all treatments combined (with moose).


Results

After four years of simulated browsing at both sites, the height of rowan trees decreased as biomass removal increased ($P < 0.001$; Fig. 1). Decreases in rowan height were greatest in the '50 moose' plot where mean plant height declined by ~ 75 cm relative to unbrowsed control trees. We did not detect a significant change in tree crown width with increasing population density ($P = 0.213$), but there was a difference of ~ 10 cm in width between the trees found in the control plots and those found in all treatment plots combined and this difference was significant ($t = 2.04$, $P = 0.037$). The number of bites per plant increased with increasing population density ($P = 0.012$) and was greatest in the '50 moose' plot where there was an increase of about two bites per plant relative to the control trees. Available bite mass declined with increasing population density ($P = 0.045$) but the only pairwise difference was between the '10' and '50 moose' plots (see Fig. 1). Finally, we observed a weak and statistically insignificant ($P = 0.072$) increase in per plant forage biomass with increasing moose browsing. However, we observed a significant increase in forage biomass of approximately 1.5 g, or the size of one bite, when comparing trees from all treatments with the unbrowsed control trees ($t = 2.69$, $P = 0.01$).

Discussion

Simulated moose browsing increased forage biomass of rowan by approximately 1.5 g/plant over unbrowsed control trees. This plant response to browsing resulted from an increase of about one or two bites per tree, a response that compensated for a slight decrease in bite mass. Therefore, the small change in forage biomass resulted from two almost compensating processes. This weak effect did not happen because of 'no response' on the part of the plant; indeed, the increased number of bites which compensated for the decreased bite mass at least prevented forage biomass from declining.

Figure 1. Mean and standard error of crown width (A), tree height (B), bites per plant (C), bite mass (D) and forage biomass (E) of rowan trees measured in 2003 in relation to browsing treatments (0, 10, 30 and 50 moose/1,000 ha) administered during 1999-2002. N = 18 within each treatment, and differing letters above bars indicate significant differences at $\alpha = 0.05$.



In contrast, plant height of rowan declined strongly and significantly to these same simulated browsing rates. Furthermore, similar tissue removal rates strongly decreased forage biomass of other plant species such as birch and Scots pine (Persson et al. 2005, De Jager & Pastor 2008). Thus, biomass production by rowan appears to at least compensate for lost tissue and may even increase production at moose population densities (equivalent to 50 moose/1,000 ha) which would otherwise decrease the productivity of Scots pine and birch, which are its main competitors.

In contrast to the increased number of twigs produced, simulated browsing decreased the height and width of plants, causing browsed plants to remain within the height reach of moose for a longer time period and the biomass produced to be concentrated within a smaller volume. Reductions in plant height resulted from replacement of the terminal shoot with several lateral shoots that only differed slightly in size. As a result, plant height

declined monotonically with increased moose density because the terminal shoots were removed with increasingly high probability with increasing moose density. However, because the lateral shoots that continually re-grew were only slightly smaller than those of unbrowsed trees, crown width remained relatively constant across all levels of the moose density treatment. This all-or-nothing response in crown width therefore results from the replacement of terminal shoots (reduction in plant height) with compensatory growth of multiple side shoots that maintain a relatively large size.

Declines in plant height appear to be a common effect of moose browsing (Pastor & Danell 2003, Persson et al. 2005, De Jager & Pastor 2008). The decreased plant height coupled with the increased number of side shoots and only slightly decreased bite mass suggests that even plants receiving extremely high levels of browsing may compensate for lost tissue and hence provide more food to moose relative to unbrowsed control plants. Bio-

mass production of the two species of birch, aspen and willow all increase under some conditions following browsing at low to moderate levels (Danell et al. 1985, Stolter 2008, De Jager & Pastor 2008, De Jager et al. 2009), and in many cases moose revisit these plants in preference to unbrowsed plants of the same species (Bergström 1984, Danell et al. 1985, Stolter 2008). Similar feedbacks most likely exist between moose and rowan but are probably sustained at even greater levels of moose browsing intensity given the monotonic growth responses presented here.

More than 90% of the winter food base of moose in northern Sweden is composed of three commonly occurring species (commercially planted Scots pine and the two naturally regenerating birch species; Persson et al. 2005). Rowan appears to have a much greater capacity to respond to browsing than either birch or pine. This may explain why rowan is often heavily and repeatedly browsed by moose even though it is far less common in the environment than these other species. In fact, moose browse as much as 80% of the rowan in some areas (Solbraa 2002). Rowan is thus highly palatable but low in abundance. The decline in plant height reported here and the ability of rowan to replace lost tissue at extremely high moose densities likely make previously browsed plants especially susceptible to browsing, when they occur alongside more abundant but less palatable plants. Under such conditions, generalist herbivores, such as moose, can reduce the abundance of the highly preferred but uncommon plant species (Solberg et al. 2003). Because the uncommon but highly preferred plant species constitutes a quantitatively small part of the total food intake of the herbivore, they may have little return impact on herbivore population density. Consequently, rare but palatable species may be continually browsed and not released from herbivory because the more abundant forage species support the herbivore population and thereby maintain high foraging rates (Coomes et al. 2003, Mysterud 2006). Thus, feeding loops between moose and rowan could eventually lead to local extinction of the seed source and long-term impacts on plant abundance and diversity despite maintenance of forage biomass.

Taking the results presented here and those presented in Persson et al. (2005) and De Jager & Pastor (2008) into account, a moose population density of 10 moose/1,000 ha will have very little influence on the growth of birch, Scots pine or

rowan, while divergent responses of these species occur at population densities of > 10 moose/1,000 ha. In many areas of Sweden moose densities are only 5-10/1,000 ha (Ball et al. 2000), but in some young forest stands during winter, population densities can be even higher than the densities we simulated. Thus, managers interested in the long-term effects of different sustained moose densities on plant community composition, soil fertility and biodiversity ought to be aware of the different species-specific growth responses to short-term browsing and the subsequent palatability of previously browsed plants, especially where moose populations exceed 10 moose/1,000 ha.

Acknowledgements - our research was supported by a grant from the National Science Foundation's Long-term Research in Environmental Biology Program, and their support is greatly appreciated. We thank everyone who helped to maintain the exclosures and apply the moose population density treatments over the years. We also thank Kjell Danell and Inga-Lill Persson for their logistical support, and for their hospitality while we visited Sweden.

References

- Ball, J.P., Danell, K. & Sunesson, P. 2000: Response of a herbivore community to increased food quality and quantity: an experiment with nitrogen fertilizer in a boreal forest. - *Journal of Applied Ecology* 37: 247-255.
- Belovsky, G.E. 1978: Diet optimization in a generalist herbivore: the moose. - *Theoretical Population Biology* 14: 105-134.
- Bergström, R. 1984: Rebrowsing on birch *Betula pendula* and *B. pubescens* stems by moose. - *Alces* 12: 870-896.
- Bergström, R. & Hjeljord, O. 1987: Moose and vegetation interactions in northwestern Europe and Poland. - *Swedish Wildlife Research Supplement* 1: 213-227.
- Cederlund, G., Ljungqvist, H., Markgren, G. & Stålfeldt, F. 1980: Foods of moose and roe deer at Grimsö in central Sweden. Results from rumen content analysis. - *Swedish Wildlife Research* 11: 169-247.
- Coomes, D.A., Allen, R.B., Forsyth, D.M. & Lee, W.G. 2003: Factors preventing the recovery of New Zealand forests following control of invasive deer. - *Conservation Biology* 17: 450-459.
- Danell, K., Bergström, R. & Edenius, L. 1994: Effects of large mammalian browsers on architecture, biomass and nutrients of woody plants. - *Journal of Mammalogy* 75: 833-844.
- Danell, K., Haukioja, E. & Huss-Danell, K. 1997: Morphological and chemical responses of mountain birch

- leaves and shoots to winter browsing along a gradient of plant productivity. - *Ecoscience* 4: 296-303.
- Danell, K., Huss-Danell, K. & Bergström, R. 1985: Interactions between browsing moose and two species of birch in Sweden. - *Ecology* 66: 1867-1878.
- De Jager, N.R. & Pastor, J. 2008: Effects of moose *Alces alces* population density and site productivity on the canopy geometries of birch *Betula pubescens* and *B. pendula* and Scots pine *Pinus sylvestris*. - *Wildlife Biology* 14(2): 251-262.
- De Jager, N.R. & Pastor, J. 2009: Declines in moose population density at Isle Royale National Park, MI, USA and accompanied changes in landscape patterns. - *Landscape Ecology* 24: 1389-1403.
- De Jager, N.R., Pastor, J. & Hodgson, A.L. 2009: Scaling the effects of moose browsing on forage distribution from the geometry of plant canopies to landscapes. - *Ecological Monographs* 79: 281-297.
- Edenius, L., Danell, K. & Bergström, R. 1993: Impact of herbivory and competition on compensatory growth of woody plants: winter browsing by moose on Scots pine. - *Oikos* 66: 286-292.
- Fridman, J., Kempe, G. & Nilsson, P. 2001: Forestry statistics 2001. Official statistics of Sweden. - Swedish University of Agricultural Sciences, Umeå, Sweden, 20 pp. (In Swedish with an English summary).
- Hägglund, B. & Lundmark, J-E. 1987: Hanledning I bonitering med skogshögskolans boniteringssystem Del 2, diagram och tabeller. - Skogsstyrelsen, Jönköping, Sweden, 70 pp. (In Swedish).
- Heroldova, M., Homolka, M. & Kamler, J.V. 2003: Breakage of rowan caused by red deer - an important factor for *Sorbetto-Piceetum* stand regeneration? - *Forest Ecology and Management* 181: 131-138.
- Krefting, L.W. 1974: The ecology of the Isle Royale moose with special reference to the habitat. - Technical Bulletin 297-1974, Forest Series 15. Agricultural Experiment Station, University of Minnesota, Minnesota, USA, 297 pp.
- Linder, P., Elfving, B. & Zackrisson, O. 1997: Stand structure and successional trends in virgin boreal forest reserves in Sweden. - *Forest Ecology and Management* 98: 17-33.
- McInnes, P., Naiman, R.J., Pastor, J. & Cohen, Y. 1992: Effects of moose browsing on vegetation and litter of the boreal forest, Isle Royale, Michigan, USA. - *Ecology* 73: 2059-2975.
- Moen, T.O. 2004: Ikke mat nok til alle elgene: Må bestandene halveres? - *Villmarksliv* 32: 18-22. (In Norwegian).
- Motta, R. 2003: Ungulate impact on rowan (*Sorbus aucuparia* L.) and Norway Spruce (*Picea abies* L. Karst) height structure in mountain forests in the eastern Italian Alps. - *Forest Ecology and Management* 181: 139-150.
- Mysterud, A. 2006: The concept of overgrazing and its role in management of large herbivores. - *Wildlife Biology* 12(2): 129-141.
- Pastor, J. & Danell, K. 2003: Moose-vegetation-soil interactions: a dynamic system. - *Alces* 39: 177-192.
- Pastor, J., Dewey, B., Naiman, R.J., McInnes, P. & Cohen, Y. 1993: Moose browsing and soil fertility in the boreal forests of Isle Royale National Park. - *Ecology* 74: 467-480.
- Pastor, J., Dewey, B., Moen, R., Maladenoff, D.J., White, M. & Cohen, Y. 1998: Spatial patterns in the moose-forest-soil ecosystem on Isle Royale, Michigan, USA. - *Ecological Applications* 8: 411-424.
- Persson, I-L., Danell, K. & Bergström, R. 2000: Disturbance by large herbivores in boreal forests with special reference to moose. - *Annales Zoologici Fennici* 37: 251-263.
- Persson, I-L., Danell, K. & Bergström, R. 2005: Different moose densities and accompanied changes in tree morphology and browse production. - *Ecological Applications* 15: 1296-1305.
- Raspé, O., Findlay, C. & Jacquemart, A-L. 2000: *Sorbus aucuparia* L. - *Journal of Ecology* 88: 910-930.
- Renecker, L.A. & Hudson, R.J. 1986: Seasonal foraging rates of free-ranging moose. - *Journal of Wildlife Management* 50: 143-147.
- Risenhoover, K.L. & Maass, S.A. 1987: The influence of moose on the composition and structure of Isle Royale forests. - *Canadian Journal of Forest Research* 17: 357-364.
- Shipley, L.A. 2007: The influence of bite size on foraging at larger spatial and temporal scales by mammalian herbivores. - *Oikos* 116: 1964-1974.
- Shipley, L.A., Blomquist, S. & Danell, K. 1998: Diet choices made by free-ranging moose in northern Sweden in relation to plant distribution, chemistry, and morphology. - *Canadian Journal of Zoology* 76: 1722-1733.
- Shipley, L.A., Illius, A.W., Danell, K., Hobbs, N.T. & Spalinger, D.E. 1999: Predicting bite size selection of mammalian herbivores: a test of a general model of diet optimization. - *Oikos* 84: 55-68.
- Shipley, L.A. & Spalinger, D.E. 1995: Influence of size and density of browse patches on intake rates and foraging decisions of young moose and white-tailed deer. - *Oecologia* 104: 112-121.
- Solberg, E.J., Sand, H., Linnell, J.D.C., Strand, O. & Wabakken, P. 2003: Store rovdyrs innverkning på hjorteviltet i Norge: Økologiske prosesser og konsekvenser for jaktuttak og jaktutøvelse. - NINA Fagrapport 63, 71 pp. (In Norwegian).
- Solbraa, K. 2002: Veiledning i elgbeitetaksering. - Skogbrukets Kursinstitutt, Honne, Biri, Norway, 28 pp. (In Norwegian).
- Stolter, C. 2008: Intra-individual plant response to moose browsing: feedback loops and impacts on multiple consumers. - *Ecological Monographs* 78: 167-183.

- Thompson, I.D., Curran, W.J., Hancock, J.A. & Butler, C.E. 1992: Influence of moose browsing on successional forest growth on black spruce sites in Newfoundland. - *Forest Ecology and Management* 47: 29-37.
- van Wieren, S.E. & Bakker, J.P. 2008: The impact of browsing and grazing herbivores on biodiversity. - In: Gordon, I.J. & Prins, H.H.T. (Eds.); *The ecology of browsing and grazing*. Springer Verlag, Berlin, Germany, pp. 263-292.
- Vivås, H.J. & Sæther, B-E. 1987: Interactions between a generalist herbivore, the moose *Alces alces* and its food resources: an experimental study of winter foraging behaviour in relation to browse availability. - *Journal of Animal Ecology* 56: 509-520.