

Vegetation types with *Cladium mariscus* (Cyperaceae) in Greece

Authors: Theocharopoulos, Michael, Georgiadis, Theodoros, Dimitrellos, Georgios, Chochliouros, Stergios, and Tiniakou, Argyro

Source: *Willdenowia*, 36(1) : 247-256

Published By: Botanic Garden and Botanical Museum Berlin (BGBM)

URL: <https://doi.org/10.3372/wi.36.36120>

BioOne Complete (complete.BioOne.org) is a full-text database of 200 subscribed and open-access titles in the biological, ecological, and environmental sciences published by nonprofit societies, associations, museums, institutions, and presses.

Your use of this PDF, the BioOne Complete website, and all posted and associated content indicates your acceptance of BioOne's Terms of Use, available at www.bioone.org/terms-of-use.

Usage of BioOne Complete content is strictly limited to personal, educational, and non - commercial use. Commercial inquiries or rights and permissions requests should be directed to the individual publisher as copyright holder.

BioOne sees sustainable scholarly publishing as an inherently collaborative enterprise connecting authors, nonprofit publishers, academic institutions, research libraries, and research funders in the common goal of maximizing access to critical research.

MICHAEL THEOCHAROPOULOS, THEODOROS GEORGIADIS, GEORGIOS DIMITRELLOS, STERGIOS CHOCHLIOUROS & ARGYRO TINIAKOU

Vegetation types with *Cladium mariscus* (*Cyperaceae*) in Greece

Abstract

Theocharopoulos, M., Georgiadis, T., Dimitrellos, G., Chochliouros, S. & Tiniakou, A.: Vegetation types with *Cladium mariscus* (*Cyperaceae*) in Greece. – Willdenowia 36 (Special Issue): 247-256. – ISSN 0511-9618; © 2006 BGBM Berlin-Dahlem.

doi:10.3372/wi.36.36120 (available via <http://dx.doi.org/>)

Populations of *Cladium mariscus* have been observed in twenty-nine localities, mainly in western Greece, fourteen of which are new records. The species grows in a broad spectrum of habitats (lakes, lagoons, shallow marshes, humid grasslands), depending on the presence of fresh water. Correspondence factor analysis of phytosociological relevés identified seven different vegetation types with *C. mariscus* in Greece, which are discussed in short.

Key words: autecology, phytosociology, wetlands, fresh water, distribution.

Introduction

Cladium mariscus L. is a robust perennial of the *Cyperaceae* that can reach up to 2.5 m in height. It is a cosmopolitan taxon with its main distribution area in Europe and the Mediterranean, but is also found in N Africa and W Asia (Meusel & al. 1965, Tutin & al. 1980). It grows in wetland environments in shallow ponds, on the shores of lakes, lagoons and water channels, and in the humid grasslands surrounding them.

For Greece, contrary to W and Central Europe (Balátová-Tuláková 1963, De Sloover 1970, Rivas Martinez & al. 1980, Pautou & Girel 1981, Devvilez & Iserentant 1981), no complete data exist on the distribution, ecology and syntaxonomic position of *Cladium mariscus* communities. Information regarding its geographical distribution is limited mainly to Haláscy (1904). Published phytosociological data for the species were available only from W Crete, including description of the *Dorycnio recti-Cladietum marisci* (Gradstein & Smittenberg 1977).

This paper deals with the present geographical distribution and ecology of *Cladium mariscus* in Greece. It includes the preliminary results of Theocharopoulos & al. (2000).

Material and methods

Flora Hellenica (Strid & Tan 1997, 2002), Med-Checklist (Greuter & al. 1984-1989), Flora Europaea (Tutin & al. 1968-80, 1993) and Flora of Turkey and the East Aegean Islands (Davis 1965-85) were used for taxa identification and nomenclature.

For the phytosociological characterisation of the *Cladium mariscus* stands, 55 relevés were analysed, mainly during May-July, in the years 2001-02, following the Braun-Blanquet (1964) approach. These relevés, ranging in size from 40 to 100 m², were taken from the most representative stands of *C. mariscus* in thirteen areas of Greece (Table 1). All relevés were interpreted using the numerical (ordination) method of the correspondence factor analysis (Benzecri & al. 1976). The groups (Fig. 2) obtained by ordination, using presence or absence of taxa (Roux 1997), are presented in a phytosociological table (Table 2), produced with the software SORT 4.0.

Names of syntaxa are used according to Braun-Blanquet & al. (1952), Horvat & al. (1974), Gradstein & Smittenberg (1977), Meriaux (1981), Biondi (1989) and Rivas-Martínez & al. (2001).

Twenty-six soil samples (two per site) were taken from depths of 30 cm (the root limit of *Cladium mariscus*) to determine the grain size (Bouyoucos 1951) and the percentage of organic matter (Nelson & Sommer 1982). In addition, pH, conductivity and total water hardness (Rand & al. 1975) were measured from all sites. At sites without surface water (humid grasslands, shores of lakes, etc.), samplings were made from nearby water channels or by digging shallow holes beside the plants.

Results

1. Distribution of *Cladium mariscus* in Greece

The geographical distribution of *Cladium mariscus* follows the wetlands of W and NW Greece. Further southwards it extends to W Crete (Fig. 1). Stands were found on 29 sites, 14 of which (Lakes Voulkaria and Kaiafa, Lagoons Korission, Keri and Saltini, Alyki Aigiou, Agios Nikolaos and Vonitsa bays, Roumani/Nafplio, Kryoneri, Agia, Kalodiki, Kalogria marshes, and Gorge Prassiano) are new records for Greece.

2. Characteristics of the *Cladium mariscus* sites

Stands were found mainly near the coast, but two sites (Lakes Agras and Cheimaditida) are mainland basins of intermediate altitude (c. 500 m). These sites are supplied with fresh water from karstic springs.

Bioclimatically (Mavromatis 1980) all sites (Table 1) belong to the humid or semi-humid bioclimatic type, except Nafplio and Artaki, which belong to the semi-dry type. The average lowest temperature of the coldest months ranges from -4.2 C (Lake Cheimaditida) to +7.6 C (Lagoon Keri).

Bedrock of the study areas is mainly limestone, while neogene formations (marls, siltstones, sandstones) and flysch are outcropping in some cases. Schists and gneisses are found in Lake Cheimaditida (Bornovas & al. 1983). The soils develop mostly on alluvial deposits and on local peat formations (Christanis 1999). Analysis of the particle size distribution shows mainly sandy-loam and loamy-sand types. Organic matter content in these soil samples ranged from 3 to 56.7 % (Table 1).

Analysis of the physicochemical water parameters (pH, conductivity and CaCO₃) showed pH values ranging from 6.1 to 7.3, conductivity from 580 to 2530 ms/cm, and CaCO₃ from 215 to 2895 mg/lit (Table 1). The above pH values are normal for wetlands surrounded by limestone, since their chemical parameters are determined to a large extent by the surrounding bedrock. The high conductivity values indicate a large inflow of dissolved solids probably due to weathering of the surrounding limestone rocks. The relief of these rocks also facilitates the transport of inorganic matter into the wetlands. The high CaCO₃ values appear to confirm the karstic origin of the waters.




Fig. 1. Distribution of *Cladium mariscus* in Greece (▲ : new records, ● : old records) – 1: Ag. Germanos; 2: Lake Vegoritida-Lake Petron; 3: Lake Cheimaditida; 4: Lake Agras; 5: Litochoro; 6: Episkepsi; 7: Korission lagoon; 8: Kalodiki marsh; 9: Saltini lagoon; 10: Ag. Nikolaos bay; 11: Lake Voulkaria; 12: Vonitsa; 13: Lake Trichonida; 14: Artaki; 15: Phaliro; 16: Messolongi; 17: Krioneri marsh; 18: Agias marsh; 19: Psathopirgos; 20: Alyki Egiou; 21: Livadi; 22: Kalogria; 23: Keri lagoon; 24: Lake Kaiafa; 25: Roumani/Nafplio marsh; 26: Elos; 27: Lake Agias; 28: Georgioupoli; 29: Prassiano gorge.

3. Vegetation types with *Cladium mariscus* in Greece

For the ordination of the 55 relevés with *Cladium mariscus* we used presence-absence data. The factor correspondence analysis (Fig. 2) allowed us to distinguish seven vegetation types where *C. mariscus* plays a constitutive or differentiating role: (1) *Elymus farctus*-*Cladium mariscus* community, (2) *Saccharum ravennae*-*Cladium mariscus* community, (3) *Dorycnio recti*-*Cladietum marisci*, (4) *Eupatorium cannabinum*-*Cladium mariscus* community, (5) *Carex riparia*-*Cladium mariscus* community, (6) *Cladietum marisci*, (7) *Schoenoplectus lacustris*-*Cladium mariscus* community.

The distribution of the vegetation groups on Axis 1 (from left to right) of Fig. 2 shows a transition from “terrestrial” to “aquatic” ecosystems, while Axis 2 (from bottom to top) seems to be related to both altitude and latitude.

3.1. *Elymus farctus*-*Cladium mariscus* community (Fig. 2: group I; Table 2)

This community is found mostly in the southern area of the coastal marsh Livadi (Artaki-Euboea area), in small stands on sandy-loam soils with fresh water layers, usually close to shallow soil channels. The differential taxa are *Juncus maritimus*, *Dittrichia viscosa*, *Cirsium creticum*, *Agrostis stolonifera*, *Lolium perenne* and *Rumex conglomeratus*, while *Elymus farctus* shows high

Table 1. Bioclimatic, geological, pedological and physicochemical data of the sampling sites.

Relevé Nr	Locality / altitude	Bioclimatic type	Geological substrate	Organic matter %	Particle size	Physicochemical parameters of the water		
						pH	Conduct. $\mu\text{S}/\text{cm}$	CaCO_3 mg/l
1-6	Marsh Artaki / 3 m	semi-dry / temperate	alluvial deposits & limestone	4-15	sandy-loam /silty-loam	7.2	630	290
7, 9, 10	Lake Kaiafa / 40 m	sub-humid / temperate	deposits & limestone	3-56.7	sandy-loam /loamy-sand	6.8	2530	660
8, 11, 12, 15	Bay Agios Nikolaos / 10 m	humid / temperate	alluvial deposits & limestone	6-13.5	sandy-loam	7.2	680	276
13, 14	Lagoon Saltini / 2m	humid / temperate	alluvial deposits & limestone	9-27.2	sandy-loam /loamy-sand	6.9	1170	258
16-18	Trichonida Lake / 18 m	sub-humid / temperate	alluvial deposits, flysch & limestone	6-15	sandy-loam	6.1	710	215
19-25	Lagoon Korission / 4 m	humid / temperate	deposits & limestone	12.8-17	sandy-loam /loamy-sand	7.1	610	287
26-29	Lagoon Keri / 3 m	humid / hot	alluvial deposits & limestone	5-19	sandy-loam	7.3	580	295
30-32	Marsh Roumani Nafplio / 5 m	semi-dry / temperate	alluvial deposits & limestone	12-23	sandy-loam	6.1	1758	2895
33-39	Lake Agras / 500 m	sub-humid / severe	terrestrial deposits & limestone	8-20	sandy-loam	7.3	640	385
40-45	Cheimaditida Lake / 800 m	sub-humid / severe	alluvial deposits, schists & gneisses	16-25	sandy-loam	6.9	1290	270
46	Marsh Kryoneri / 1m	sub-humi / temperate	alluvial deposits & limestone	7-38	loamy sandy-loam	6.8	2370	460
49	Marsh Agia / 1 m	sub-humid / temperate	alluvial deposits	11-35	sandy-loam /loamy-sand	6.8	985	385
47, 48, 50-55	Lake Voulkaria / 0.5 m	humid / temperate	alluvial deposits & limestone	12-48	sandy-loam	6.6	1530	447

cover values. The location of this community close to the seashore together with sandy-loam/silty-loam soils favours the presence of the above ammophytic species.

Characteristic in this community is the presence of *Carex hispida*, which grows in small groups, and (with the exception of *Phragmites australis*) the absence of Phragmitetea species. The presence of the subnitrophilous species *Hordeum murinum*, *Stellaria media*, *Galium aparine* and *Bromus sterilis* is due to human impact. Most of the marsh has been drained and transformed to intensively cultivated agricultural land, meadow and pasture.

3.2. *Saccharum ravennae-Cladium mariscus* community (Fig. 2: group II; Table 2)

This community grows in damp hollows among the sand dunes with shallow freshwater layers. It is dominated by *Saccharum ravennae* and *Schoenus nigricans* and shows a variant with *Juncus subulatus* and *Carex extensa*. It is found in the coastal marsh of Lake Kaiafa, Lake Korission, and the coastal area between Agios Nikolaos bay and Saltini lagoon. At these sites *Cladium mariscus* participates with various cover values. It can reach such a dominance that it hinders the growth of the above taxa, especially the small *S. nigricans* with high demand of light, and forms dense homogeneous stands with only a few *Phragmites australis* individuals. These are humid grasslands with few or many damp places and plentiful organic matter, forming peat in places.

By the presence of *Rumex conglomeratus*, *Lythrum junceum*, *Pulicaria dysenterica* and *Dittrichia viscosa* this community can be assigned to the alliance Molinio-Holoschoenion. A related association with a similar floristic composition but without *Cladium mariscus* (Eriantho




Fig. 2. Ordination diagram (axis 1, 2) of the fifty-five relevés with *Cladium mariscus* with the seven vegetation types (here in Roman numerals) distinguished.

ravennae-Schoenetum nigricantis (Pign. 1953) Gehu 1984) has been previously recorded from Corfu Island (Biondi 1989).

3.3. *Dorycnio recti-Cladietum marisci* Gradstein & Smittenberg 1977 (Fig. 2: group III; Table 2) Stands of this vegetation type appear on wet, muddy soils on alluvial deposits on shores of fresh-water lakes or lagoon environments fed with fresh water. *Cladium mariscus* populations with 50-80 % cover co-exist with the tall legume *Dorycnium rectum*. The following taxa of the Molinio-Holoschoenion give the association its physiognomy: *Dittrichia viscosa*, *Rumex conglomeratus*, *Juncus inflexus*, *Carex distans* and *Pulicaria dysenterica*.

The *Dorycnio recti-Cladietum marisci* was found in four areas, each with its own local variant: (a) Korission lagoon, along the southwest shore, variant with *Typha latifolia*; (b) Keri lagoon, along the southwest shore, variant with *Carex divisa*; (c) Lake Trichonida, on the south shore (Ag. Triada-Dafnias), in abandoned cultivations and irrigation channels, variant with *Equisetum telmateia*; (d) Roumani marsh close to Nafplio, mainly in scattered places with permanent humidity and along small non-paved irrigation channels between cultivations, variant with *Equisetum arvense*.

3.4. *Eupatorium cannabinum-Cladium mariscus* community (Fig. 2: group IV; Table 2)

This community is found in the SW of the artificial Lake Agras, sometimes in very dense stands. The community grows on rich organic soils often flooded in winter. The presence of *Juncus inflexus*, *Pulicaria dysenterica*, *Cirsium creticum*, *Carex distans*, *Lythrum junceum* and *Agrostis stolonifera* allows the *Eupatorium cannabinum-Cladium mariscus* community to be placed in the Molinio-Holoschoenion alliance. The community forms a variant with *Molinia coerulea*.

3.5. *Carex riparia-Cladium mariscus* community (Fig. 2: group V; Table 2)

This community is found as shore vegetation in two small areas NW of Lake Cheimaditida. It is characterized by *Carex riparia* which, together with *Cladium mariscus*, forms fairly dense

Table 2. Communities mit *Cladium mariscus* – 1: *Elymus farctus-Cladium mariscus* community; 2: *Saccharum ravennae-Cladium mariscus* community; 3: *Dorycnio recti-Cladietum marisci*; 4: *Eupatorium cannabinum-Cladium mariscus* community; 5: *Carex riparia-Cladium mariscus* community; 6: *Cladietum marisci*; 7: *Schoenoplectus lacustris-Cladium mariscus* community.

Additional, occasional species: *Carex pendula* (21, 52: 1, 25: +), *Cirsium arvense* (33, 35, 39: +), *Nymphaea alba* (51: +, 54, 55: 1), *Paspalum paspalodes* (53: +, 54: 2, 55: 1), *Silybum marianum* (1, 2: +), *Bromus sterilis* (1: 2, 3: 1), *Centaureum erythraea* (7, 12: +), *Holcus lanatus* (8:1, 14: +), *Brachypodium sylvaticum* (7, 31: +), *Plantago major* (26:1, 32: r), *Urtica dioica* (1: r), *Sisymbrium officinale* (2: +), *Centaurea solstitialis* (4: +), *Tamarix parviflora* (8: +), *Ulmus minor* (12: +), *Mentha longifolia* (12: 1), *Gladiolus* sp. (12: 1), *Oenanthe silaifolia* (12: 1), *Smilax aspera* (14: +), *Pistacia lentiscus* (15: +), *Inula crithmoides* (15: +), *Arundo donax* (16: +), *Fraxinus ornus* (16: r), *Hydrocotyle vulgaris* (20: 1), *Vitex agnus-castus* (21: +), *Carex otrubae* (26: 1), *Scirpoides holoschoenus* (30: 1), *Melilotus officinalis* (31: 1), *Sinapis arvensis* (31: +), *Prunella vulgaris* (32: +), *Melissa officinalis* (32: 1), *Trifolium fragiferum* (32: +), *Salix cinerea* (32: +), *Hypericum perforatum* (35: +), *Galium verum* (39: +), *Equisetum ramosissimum* (49: +), *Solanum dulcamara* (53: +), *Veronica beccabunga* (54: 2).

Relevé number	111111	11112222222222333	3333333	444444	444455	5555
	123456	789012345	67890123456789012	3456789	012345	678901 2345
Number of species	111111	1111111111	11111121111111 223	1111 11	112111	11111 1111
	965225	380158345	52126310179149640	4760925	832632	960111 6421
Plot size (x 10 m2)	155485	168451418	55455565555115855	1111111	555555	511511 5555
	0	0 0 0	00	0000000		00 00
Total vegetation cover (x 100%)	111111	1111111111	111111111111111111	1111111	111111	111111 1111
Date (Year/month/day)	000000	000000000	00000000000000000	0000000	000000	000000 0000
	222222	1111111111	111111111111111111	1111111	111111	222222 2222
	000000	000000000	00000000000000000	0000000	000000	000000 0000
	555555	767676766	44455555555555888	7777777	777777	666666 6666
	222222	222101011	22211111112222000	2222222	222222	211211 1111
	888888	646565655	7776666666777222	1111111	000000	166366 6666
Water depth (cm)						256156 1200
						000000 0555
Community	1	2	3	4	5	6 7
Characteristic and differential taxa of the vegetation types:						
<i>Cladium mariscus</i>	444324	422525554	33355555544324454	4345445	454534	345455 3222
<i>Elymus farctus-Cladium mariscus</i> community						
<i>Elymus farctus</i>	232552	-----	-----	-----	-----	-----
<i>Juncus maritimus</i>	+++++	-----+	-----13	-----	-----	-----
<i>Carex hispida</i>	2-21-1	-----	-----11	-----	-----	-----
<i>Lolium perenne</i>	-1+-1-	-----	-----	---+---	-----	-----
<i>Saccharum ravennae-Cladium mariscus</i> community						
<i>Saccharum ravennae</i>	-----	222142131	++-----+-----11+	-----	-----	-----
<i>Schoenus nigricans</i>	-----	114112111	--+-----1-----	-----	-----	-----
<i>Juncus subulatus</i>	-----	1211+---	-----+-----	-----	-----	-----
<i>Carex extensa</i>	-----	1121	-----	-----	-----	1121
<i>Cynanchum acutum</i>	-----	++1---1--	-----	-----	-----	-----
<i>Dorycnio recti-Cladietum marisci</i>						
<i>Dorycnium rectum</i>	-----	-1-----	11--1+--+1222211	-----	-----	--+-+-----
<i>Equisetum telmateia</i>						
<i>Equisetum telmateia</i>	-----	-----	232	-----	-----	-----
<i>Periploca graeca</i>	-----	----11---	111-----+---	-----	-----	-----
<i>Typha latifolia</i>						
<i>Typha latifolia</i>	-----	-----	---+++111+-----	-----	-----	-----
<i>Stachys palustris</i>	-----	-----	----111-----	-----	1-----	-----

Relevé number	111111	1111222222222333	3333333	444444	444455	5555	
	123456	789012345	67890123456789012	3456789	012345	678901	2345
<i>Carex divisa</i>	-----	-----	-----1--11111--	-----	-----	-----	----
<i>Aster tripolium</i>	-----	-1-----1--	-----111----	-----	-----	-----	----
<i>Orchis laxiflora</i>	-----	-----	-----+1----	-----	-----	-----	----
<i>Equisetum arvense</i>	-----	-----	-----111	-----	-----	-----	----
<i>Picris echioides</i>	-----	-----	-----111	-----	-----	-----	----
<i>Aster squamatus</i>	-----	-----	-----1-1	-----	-----	-----	----
<i>Verbena officinalis</i>	-----	-----	-----+1	-----	-----	-----	----
<i>Potentilla reptans</i>	-----	-----	-----+1	-----	-----	-----	----
<i>Eupatorium cannabinum-Cladium mariscus</i> community							
<i>Eupatorium cannabinum</i>	-----	-----	111-----	1122212	1111--	-+-----	----
<i>Satureja vulgaris</i>	-----	-----	-----	+++--1-	-----	-----	----
<i>Juncus articulatus</i>	-----	-----	-----	++111+	+-----	-----	----
<i>Molinia coerulea</i>	-----	-----	-----	121----	-----	-----	----
<i>Carex riparia-Cladium mariscus</i> community							
<i>Carex riparia</i>	-----	-----	-----	-----	222222	-----	----
<i>Cyperus flavescens</i>	-----	-----	-----	-----	322221	-----	----
<i>Lycium europaeum</i>	-----	-----	-----	-----1	--1-+-	-----	----
<i>Ononis spinosa</i>	-----	-----	-----	-----	--rr+	-----	----
<i>Calamagrostis epigejos</i>	-----	-----	-----	-----	++++	-----	----
<i>Cladietum marisci</i>							
<i>Phragmites australis</i>	321222	++11+1+11	3-3+1-11223334+1	-----	++11--	522322	2322
<i>Schoenoplectus lacustris-Cladium mariscus</i> community							
<i>Schoenoplectus lacustris</i>	-----	-----	-----	-----	+--11-1	-----+	3333
<i>Molinio-Holoschoenion vulgaris</i>							
<i>Dittrichia viscosa</i>	++---+	12111+211	--+1111+111221+-	-----	-----	-----	----
<i>Rumex conglomeratus</i>	1-1-+-	+---+---	++-+1-++11-1-+-	-r-----	-----	-----	----
<i>Pulicaria dysenterica</i>	-----	---+1-21	---1-----1+1	111--+1	++---	-1-1--	----
<i>Juncus inflexus</i>	-----	--1-1-1--	-----2-1	--+++-	-----	-----	----
<i>Holoschoenetalia and Molinio-Arrhenatheretea</i>							
<i>Cirsium creticum</i>	1+11++	+1+++++++	1212+11--1111222	1111111	++-+11	++-----	----
<i>Agrostis stolonifera</i>	122222	-----	-----+++	333----	++-+11	-----	+---
<i>Lippia nodiflora</i>	-----	1-1--1--	---1-----	1+-----	11-1--	-----	----
<i>Lythrum junceum</i>	-----	-1--1-1-	-21-----	1-1++	11111-1	-----	----
<i>Carex distans</i>	-----	-----	1+++--1+1-1-	++1++++	---1-	-----	----
<i>Cynodon dactylon</i>	-----	-----	-----+	-----	+-11	-----	----
<i>Juncus effusus</i>	-----	-----	-----1----	-----	-----	-----	+---
<i>Magnocaricion elatae and Magnocaricetalia</i>							
<i>Samolus valerandi</i>	-----	-----1-	---1-1-11----	-----	11111-	--+---	----
<i>Lycopus europaeus</i>	-----	-----	-----1-----	111	-1+-1-	11+1+	1--+-
<i>Athaea officinalis</i>	-----	-----	-----+	-----	1+---	-----	----
<i>Galium palustre</i>	-----	-----	-----	+++--+	1+-11	+-+--	----
<i>Phragmition communis and Phragmitetalia</i>							
<i>Typha angustifolia</i>	-----	-----	-----1	-----	-----	122-21	2334
<i>Sparganium erectum</i>	-----	-----	-----	-----	-----	--1++	+111
<i>Phragmito-Magnocaricetea</i>							
<i>Mentha aquatica</i>	-----	-----111	---2111-11----	11-	+1111-	111+-1	----
<i>Apium nodiflorum</i>	-----	-----	---111	-----	-----	++111	+---
<i>Iris pseudacorus</i>	-----	-----	-----1-	-----	-----	---++	----
<i>Oenanthe fistulosa</i>	-----	-----	-----2-1-	-----	-----	-----	+---
<i>Lythrum salicaria</i>	-----	-----	-----+	-----	+-	111+1+	+---
<i>Lysimachia vulgaris</i>	-----	-----	-----+	-----	-----	-+1+1	+---
<i>Alisma plantago-aquatica</i>	-----	-----	-----	-----	+++--	-----	++++

continued on the next page

Relevé number	111111	1111222222223333	3333333	444444	444455	5555	
	123456	789012345	67890123456789012	3456789	012345	678901	2345
Other species							
<i>Hordeum murinum</i>	1+1++1	-----	-----	-----	-----	-----	----
<i>Gastridium ventricosum</i>	11-+11	-----	-----	-----	-----	-----	----
<i>Stellaria media</i>	111--1	-----	-----	-----	-----	-----	----
<i>Calystegia sepium</i>	111111	----1111	11--11+1+111112-1	1111111	--1---	111111	11++
<i>Galium aparine</i>	222212	----1----	---1-1-1-1+-11+	-----	-----	-----	----
<i>Cirsium vulgare</i>	r-r--r	-----	-----	+-----	+++++	-----	----
<i>Polypogon monspeliensis</i>	+1----	-----	-----	-----	++1--	-----	+---
<i>Daucus carota</i>	-----+	-1-----	-1+-----+1-	-----+	-----	-----	----
<i>Juncus acutus</i>	-----	+--2--+	---+---+1+---+--	-----	-----	-----	++++
<i>Lotus corniculatus</i>	-----	-1---+--1	-----1-1	-----	-----	-----	----
<i>Rubus sanctus</i>	-----	+---+--	211-+---+--1+1+	+-----	-----	-----	----
<i>Myrtus communis</i>	-----	---2++11	---1--rr-r	-----	-----	-----	----
<i>Rubia peregriana</i>	-----	-----	1---111-1-----+	-----	-----	-----	----
<i>Cyperus longus</i>	-----	-----	-----1+-----	-----	+1---	-----	++-1
<i>Epilobium hirsutum</i>	-----	-----	-----+-----+	+1-++1+	---+--+	11----	11--
<i>Poa trivialis</i>	-----	-----	-----1-	-----	-----	-----	1+1-

stands with a *Magnocaricion elatae* physiognomy. It grows on slightly boggy substrates which are flooded at least during winter.

3.6. *Cladietum marisci* Zobrist 1935 (Fig. 2: group VI; Table 2)

This syntaxon is characterized by the dominance of *Cladium mariscus*, which forms dense, impassable stands, together with *Phragmites australis* and *Typha angustifolia*.

The *Cladietum marisci* is found in the marshes of Agia and Kryoneri, in small stands at water depths of 5-25 cm. It is also found in Lake Voulkaria where *Cladium mariscus* forms extensive stands in a 100 m wide zone around the lake, at water depths of less than 80 cm. At water depths exceeding 50-60 cm, which are less favourable for *C. mariscus*, the species forms a floating platform composed of its root network, which may or may not be associated with other species (common reed in particular, see also Devvilez & Iserentant 1981).

In Greece, the above syntaxon is best placed in the *Phragmition communis* alliance, as indicated by the occurrence of *Lysimachia vulgaris*, *Apium nodiflorum*, *Lythrum salicaria*.

3.7. *Schoenoplectus lacustris*-*Cladium mariscus* community (Fig. 2: group VII; Table 2)

Stands of this type are characterized by *Schoenoplectus lacustris* and *Phragmites australis*. *Cladium mariscus* participates in small groups. The community belongs to the *Phragmition* alliance, as shown by the presence of *Typha angustifolia*, *Alisma plantago-aquatica* and *Sparganium erectum*. It is found in Lake Voulkaria, mostly in the north-western and southern areas of the lake, in very wide zones.

Discussion and conclusions

Cladium mariscus is a thermophilous species (Devvilez & Iserentant 1981). It prefers lakes, lagoons, shallow marshes and wet meadows over limestone, all with groundwater levels close to the soil surface. Periods of flooding are possible. The creeping rhizome of *C. mariscus* is only poorly anchored in the soil, and this makes it sensitive to fluctuations in groundwater level. The plant grows in waters of karstic origin that are rich in calcium carbonate (CaCO₃).

Communities of *Cladium mariscus* are found on soils rich in organic matter, as well as on sandy and gravely mineral substrate with sandy-loam and loamy-sand layers. They also grow in water of varied pH, from alkaline to neutral, but generally at depths of less than 80 cm, where their root networks form floating platforms. *C. mariscus* shows a broad ecological tolerance, invading various plant communities. Communities with *C. mariscus* form two principal types as regards their aspect, origin and structure:

(1) “Aquatic” communities. – These develop on the water surface by the centripetal progression of its roots, which form a floating raft. In most cases, the *Cladietum marisci* is initially very dense and paucispecific, even monospecific, and a very significant quantity of litter is accumulated. *Cladium mariscus* may be accompanied by other species with high biomass production, e.g. common reed, which participates in structuring the floating carpet. The presence of *Schoenoplectus lacustris*-*C. mariscus* communities is limited to small inland locations usually bordering reed populations and periodically flooded wetlands.

(2) “Terrestrial” communities. – These originate from invasion of *Cladium mariscus* in pre-existing vegetation. *C. mariscus* tolerates a certain variation in the physicochemical parameters of the vegetation groups into which it penetrates, such as formations of the *Molinio-Holoschoenion* and *Magnocaricion elatae* alliances described above. In the *Saccharum ravennae*-*C. mariscus*- and *Eupatorium cannabinum*-*C. mariscus* communities *C. mariscus* can penetrate species-rich stands with initially a characteristic species composition. It gradually becomes very dense and dominant and finally the species diversity can become extremely poor. In *Dorycnio-Cladietum marisci* and the *Elymus farctus*-*C. mariscus* community a low diversity of species is maintained.

The *Saccharum ravennae*-*Cladium mariscus* community, which grows in wet hollows of sand dunes, forms an original floristic assemblage threatened by possible changes in water level. Like the *Eupatorium cannabinum*-*C. mariscus* community, which grows in mainland basins at intermediate altitudes on wet organic soils, it might be considered as stable.

The *Dorycnio recti-Cladietum marisci* with its different local variants has the widest distribution. We found stands on alluvial soils that remain wet during summer, probably due to capillary rising of ground water. It grows in water bodies along channel banks, wet hollows, or inland areas and forms small, fairly dense populations.

The *Elymus farctus-Cladium mariscus* community is found near the sea in a transitional position between the *Molinio-Holoschoenion* and *Agropyron juncei* alliances of the class *Amphiletea*, as shown by its floristic composition and ecological conditions.

The *Carex riparia-Cladium mariscus* community constitutes a transition between aquatic and terrestrial communities. *Carex riparia* has a Eurasian, Suboceanic and Sub-Mediterranean distribution (Balátová-Tuláčková 1995) where, on wet substrates, it competes successfully with *C. mariscus* while in drier conditions its competitive capacity is reduced.

Cladium mariscus is the principal component of the priority Natura 2000 habitat type 7210: Calcareous fens with *C. mariscus* and *Carex davalliana* (Directive 92/43/EEC, European Commission DG. XI 1996). This habitat type is influenced both by hydrological changes in the main wetland and soil erosion at its borders. *C. mariscus* is sensitive to both changes in hydrological conditions and human influences (land reclamation, extensive cultivation, tourism, fires and water pollution). In many areas populations of *C. mariscus* are reduced dramatically and severely endangered (Lake Cheimaditida, Keri lagoon, and Roumani/Nafplio, Artaki and Kalodiki marshes). It also appears in other habitat types, e.g. ‘humid dune slacks’ (Natura 2000 code: 2190). The healthiest populations were observed at Lake Agras, Voulkaria (where the largest populations were found), Kaiafa, the Saltini and Korission lagoons, and Agios Nikolaos bay.

References

- Balátová-Tuláčková, E. 1963: Zur Systematik der europäischen Phragmitetea. – *Preslia* **35**: 118-122.
 — 1995: L'écologie et le rôle des magnocaricaux dans la succession vers les groupements de l'ordre des Molinietales. – *Colloq. Phytosoc.* **24**: 561-570.
 Benzécri, J. P. & al. 1976: L'analyse des données II: L'analyse des correspondances. – Paris.
 Biondi, E. 1989: The vegetation of sedimentary low coasts in Corfu Island. – *Colloq. Phytosoc.* **19**: 401-426.
 Bornovas, J. & Rondogianni-Tsiambaou, T. 1983: Geological map of Greece, 1: 500 000. – Athens.
 Bouyoucos, G. H. 1951: A recalibration of the hydrometer method for making mechanical analysis of soils. – *Agronomy J.* **43**: 434-438.
 Braun-Blanquet, J. 1964: *Pflanzensoziologie*, ed. 3. – Wien.

- , Roussine, N. & Nègre, R. 1952: Les groupements végétaux de la France Méditerranéenne. – Montpellier.
- Christanis, K. 1999: Geologiki-kitasmatologiki meleti ombrogenon kai topogenon tyrfonon tou Ellinikou chorou. – Patras.
- Davis, P. H. (ed.) 1965-85: Flora of Turkey and the East Aegean Islands **1-9**. – Edinburgh.
- De Sloover, J. R. 1970: Les peuplements de *Cladium mariscus* du district côtier belge, leur origine et leur position phytosociologique. – *Lejeunia*, ser. 2, **51**: 1-26.
- Devvilez, F. & Iserentant, R. 1981: Influence du climat et des conditions mésologiques sur la croissance et le développement de *Cladium mariscus* (L.) Pohl. – *Colloq. Phytosoc.* **10**: 85-114.
- European Commission 1996: Natura 2000. – Interpretation manual of European Union habitats. – Brussels.
- Gradstein, S. R. & Smittenberg, J. H. 1977: The hydrophilous vegetation of Western Crete. – *Vegetatio* **34**: 65-86. [[CrossRef](#)]
- Greuter, W., Burdet, H. M. & Long, G. 1984-89: Med-Checklist **1, 3, 4**. – Genève & Berlin.
- Halacsy, E. 1904: Conspectus florae Graecae **3**. – Lipsiae.
- Horvat, I., Glavač, V., & Ellenberg, H. 1974: Vegetation Südosteuropas. – Stuttgart.
- Mavromatis, G. 1980: [Le bioclimat de la Grèce. Relations entre le climat et la végétation naturelle. Cartes bioclimatiques.]. – Athènes [In Greek]
- Meriaux, J. L. 1981: La classe des Phragmitetea dans le Nord-Ouest de la France. – *Colloq. Phytosoc.* **10**: 139-147.
- Meusel, H., Jäger, E. & Weinert, E. 1965: Vergleichende Chorologie der zentraleuropäischen Flora. – Jena.
- Nelson, D. W. & Sommer, L. E. 1982: Total carbon, organic carbon and organic matter (Walkley-Black procedure). – In: Page, R. H. & al. (ed.), *Methods of soil analysis*, ed. 2. – Madison.
- Pautou, G. & Girel, J. 1981: Les associations végétales à *Cladium mariscus* dominant dans la vallée du Rhône entre Lyon et Genève. – *Colloq. Phytosoc.* **10**: 333-349.
- Rand, M. C., Greenberg, A. E. & Taras, M. G. 1975: Standard methods for the examination of water and wastewater, ed 14. – Washington.
- Rivas-Martínez, S., Costa, M., Castroviejo, S. & Valdés, E. 1980: Vegetación de Doñana (Huelva, España). – *Lazaroa* **2**: 5-189.
- , Fernández-González, F., Loidi, J., Lousã, M. & Penas, A. 2001: Syntaxonomical checklist of vascular plant communities of Spain and Portugal to association level. – *Itin. Geobot.* **14**: 5-341.
- Roux, M. 1997: Le statoscope. Méthodes statistiques pour la biologie. Version 1.6. – Marseille.
- Strid, A. & Tan, K. (ed.) 1997: Flora hellenica **1**. – Koenigstein.
- & — 2002: Flora hellenica **2**. – Ruggell.
- Theocharopoulos, M., Dimitrellos, G., Chochliouros, S. & Georgiadis, T. 2000: Distribution and synecology of *Cladium mariscus* L. in Greece. – Pp. 161-167 in: Anon. (ed.), *Proceedings 8th Congr. Hellenic Bot. Soc.* [in Greek].
- Tutin, T. G., Heywood, V. H., Burges, N. A., Moore, D. M., Valentine, D. H., Walters, S. M. & Webb, D. A. (ed.) 1964-1980: *Flora europaea* **2-5**. – Cambridge, etc.
- , Burges, N. A., Chater, A. O., Edmonson, J. R., Heywood V. H., Moore, D. M., Valentine, D. H., Walters, S. M. & Webb, D. A. (ed.) 1993: *Flora europaea*, ed **2, 1**. – Cambridge, etc.

Address of the authors:

M. Theocharopoulos, T. Georgiadis, G. Dimitrellos, S. Chochliouros & A. Tiniakou, Botanical Institute, Division of Plant Biology, Department of Biology, University of Patras, GR-26500 Patras, Greece; e-mail: georgiad@upatras.gr