



## **Development of Microsatellite Markers For The Semi-Natural Grassland Herb *Veronicastrum japonicum* (Plantaginaceae)**

Authors: Nakahama, Naoyuki, Izuno, Ayako, Arima, Kurumi, and Isagi, Yuji

Source: Applications in Plant Sciences, 4(2)

Published By: Botanical Society of America

URL: <https://doi.org/10.3732/apps.1500096>

---

BioOne Complete ([complete.BioOne.org](https://complete.BioOne.org)) is a full-text database of 200 subscribed and open-access titles in the biological, ecological, and environmental sciences published by nonprofit societies, associations, museums, institutions, and presses.

Your use of this PDF, the BioOne Complete website, and all posted and associated content indicates your acceptance of BioOne's Terms of Use, available at [www.bioone.org/terms-of-use](https://www.bioone.org/terms-of-use).

Usage of BioOne Complete content is strictly limited to personal, educational, and non - commercial use. Commercial inquiries or rights and permissions requests should be directed to the individual publisher as copyright holder.

---

BioOne sees sustainable scholarly publishing as an inherently collaborative enterprise connecting authors, nonprofit publishers, academic institutions, research libraries, and research funders in the common goal of maximizing access to critical research.

## DEVELOPMENT OF MICROSATELLITE MARKERS FOR THE SEMI-NATURAL GRASSLAND HERB *VERONICASTRUM JAPONICUM* (PLANTAGINACEAE)<sup>1</sup>

NAOYUKI NAKAHAMA<sup>2,3</sup>, AYAKO IZUNO<sup>2</sup>, KURUMI ARIMA<sup>2</sup>, AND YUJI ISAGI<sup>2</sup>

<sup>2</sup>Graduate School of Agriculture, Kyoto University, Kyoto 606-8502, Japan

- **Premise of the study:** *Veronicastrum japonicum* (Plantaginaceae) grows in grasslands on Honshu Island, Japan, and is threatened by habitat loss because of rapid land development over recent decades. For the genetic characterization of the remaining populations, microsatellite markers were developed.
- **Methods and Results:** Twelve polymorphic microsatellite loci were developed using next-generation sequencing. The number of alleles per locus ranged from two to 24 (mean 7.7), and the expected heterozygosity per locus ranged from 0.35 to 0.94 (mean 0.68).
- **Conclusions:** These markers can be used for genetic studies in conservation, such as the evaluation of genetic diversity and genetic structure.

**Key words:** conservation; grassland herb; next-generation sequencing; Plantaginaceae; *Veronicastrum japonicum*.

*Veronicastrum japonicum* (Nakai) T. Yamaz. (Plantaginaceae) is a perennial plant species growing in the grasslands of mountainous regions on Honshu Island, Japan (Ohwi, 1978). In recent decades, the number of *V. japonicum* populations has rapidly decreased, and this species is now classified as near threatened or higher in five of the 47 Prefectural Red Lists of Japan (Association of Wildlife Research and EnVision, 2007). The decline of *V. japonicum* populations can be attributed to shrinking grasslands resulting from land development or the abandonment of traditional management. Furthermore, serious grazing damage by deer, which has explosively increased in the past few decades, has threatened *V. japonicum* populations (Takatsuki, 2009; Nagaike et al., 2014). *Veronicastrum japonicum* is also the main larval food plant species of *Melitaea ambigua* (Butler) (Nymphalidae), a rare butterfly species classified as endangered in the Japanese Red List (Ministry of the Environment of Japan, 2012). Accordingly, knowledge of the genetic diversity and structure of *V. japonicum* populations is important for the conservation management of *V. japonicum* and *M. ambigua*. Thus, 12 nuclear microsatellite markers for *V. japonicum* were developed using next-generation sequencing.

### METHODS AND RESULTS

A fresh leaf sample of *V. japonicum* was collected from an individual plant growing in Gunma Prefecture, Japan, and genomic DNA was extracted using

<sup>1</sup>Manuscript received 20 August 2015; revision accepted 1 October 2015.

The authors thank Sohgo Takahashi, Kei Uchida, and Atushi Ushimaru for assistance with the sampling. We also thank Tomohiro Furuta for help with making the voucher specimens.

<sup>3</sup>Author for correspondence: naoyuki.halobates@gmail.com

doi:10.3732/apps.1500096

the DNeasy Plant Mini Kit (QIAGEN, Germantown, Maryland, USA). A DNA fragment library was constructed with an Ion Xpress Plus Fragment Library Kit (Life Technologies, Foster City, California, USA), amplified with an Ion PGM Template OT2 400 Kit, and subsequently sequenced with an Ion PGM Sequencing 400 Kit and an Ion 318 Chip v2 (both from Life Technologies), yielding 161,880 reads (size range 8–618 bp; mean 220 bp). The reads were screened using Primer3 software (Rozen and Skaletsky, 1999) embedded in MSATCOMMANDER version 0.8.2 (Faircloth, 2008) to identify dinucleotide and trinucleotide loci of at least eight and seven repeats, respectively. After screening, 284 putative microsatellite loci were identified. Using these loci, a total of 57 primer pairs with melting temperatures ( $T_m$ ) of 57–61°C, GC content of 35–75%, and fragment sizes of 150–350 bp were designed.

Amplification and specificity of 34 out of the 57 primer pairs were tested using four *V. japonicum* individuals. For all confirmed microsatellite loci, the forward primers were synthesized using one of the three different M13 sequences, 5'-CACGACGTTGTAAAACGAC-3', 5'-CTATAGGGCAGCGTGGT-3', or 5'-TGTGGAATTGTGAGCGG-3' (Boutin-Ganache et al., 2001) as shown in Table 1. Each PCR amplification was performed in a final volume of 5  $\mu$ L, containing approximately 16 ng of template DNA, 2.5  $\mu$ L of 2 $\times$  Multiplex PCR Master Mix (QIAGEN, Valencia, California, USA), 0.01  $\mu$ M of forward primer, 0.2  $\mu$ M of reverse primer, and 0.1  $\mu$ M of M13 (fluorescent-labeled) primer. The PCR thermal profile was as follows: initial denaturation at 95°C for 15 min; followed by 33 cycles of 94°C for 30 s, 57°C for 1.5 min, and 72°C for 1 min; and a final extension at 60°C for 30 min. The PCR products were detected on the ABI Prism 3130 Genetic Analyzer (Applied Biosystems, Waltham, Massachusetts, USA). Fragment lengths were calculated using GeneMapper software (Applied Biosystems). Of the 34 primer pairs tested, 13 did not amplify specific regions and nine did not amplify in all four individuals. Twelve primer pairs amplified in all four individuals, and the lengths of amplified fragments were scored unambiguously. The 12 primers were used for analysis of 21 and 27 individuals sampled from two populations located in Gunma and Nagano prefectures, respectively (Gunma: 36°26'12"N, 138°25'27"E; Nagano: 35°57'25"N, 137°37'49"E). One specimen from each population was collected and deposited in the Kyoto University Museum herbarium (accession numbers: KYO\_00019997 and KYO\_00019998, respectively). Population statistics were calculated using GenAlEx version 6.4 (Peakall and Smouse, 2006). Tests for deviation from Hardy-Weinberg equilibrium (HWE) and linkage disequilibrium (LD) between loci were performed using FSTAT version 2.9.3 (Goudet, 1995).

The number of alleles per locus ranged from two to 24 (mean 7.7) (Table 2). The observed and expected heterozygosities per locus ranged from 0.30 to 1.00

Applications in Plant Sciences 2016 4(2): 1500096; <http://www.bioone.org/loi/apps> © 2016 Nakahama et al. Published by the Botanical Society of America.

This work is licensed under a Creative Commons Attribution License (CC-BY-NC-SA).

TABLE 1. Characteristics of microsatellite loci for *Veronicastrum japonicum*. All values are based on 48 samples from two populations in Gunma and Nagano prefectures in Japan.<sup>a</sup>

Locus	Primer sequences (5'–3')	Repeat motif	Fluorescent label <sup>b</sup>	Allele size range (bp)	GenBank accession no.
Veja004	F: AGGAGCCCAGTTTGCCCTAC R: CAAATGCTAATTGATCTCGACC	(CT) <sub>9</sub>	FAM	211–221	LC030210
Veja005	F: CCCTCTTTCTGGAAGTTTGAGC R: ACGACTCCTCCTTGCTTAGG	(TGT) <sub>6</sub>	VIC	232–237	LC030211
Veja007	F: GGGACATTGGAATGGAGGC R: ATTGATGCTGAACGAGGC	(TCT) <sub>6</sub>	FAM	166–176	LC030212
Veja009	F: CATTAAATACTCTCTCCGCAGG R: GCTCTTCTCAGAGTTTCC	(CT) <sub>9</sub>	NED	233–253	LC030213
Veja011	F: TGATGCTTGCCATCGTTCC R: AACTCAGCTGTTCTCCCG	(GCT) <sub>9</sub>	VIC	200–221	LC030214
Veja012	F: CTTCTTCCACCAACCAAGGG R: AACAGGACTCGCTCGTCAG	(GT) <sub>9</sub>	NED	308–332	LC069371
Veja015	F: CTGTCCCACCTTTAGATTCCC R: GTCCCTTCAAGCTTCCC	(TC) <sub>9</sub>	NED	187–193	LC030215
Veja018	F: CACCAATCATTTCCGCAAACC R: ACAACCGCCCTTTATGATCTAC	(GA) <sub>12</sub>	NED	251–321	LC030216
Veja021	F: TGGACACATCAACCAGATTTC R: TCCTGGATTCAACGAGAACC	(TAT) <sub>7</sub>	NED	194–209	LC030217
Veja025	F: AGGCCAATCAAATAATTACGC R: ACGCTTGTAGAGAGGTTAGGC	(CA) <sub>9</sub>	FAM	173–192	LC030218
Veja029	F: GCGCACTGTTAAGGACAGG R: CGTGAACAACGGCATAACC	(TGT) <sub>6</sub>	VIC	178–196	LC030219
Veja032	F: CAACGCCAGCCATCCATAC R: TTGGAGGAGAATGACACCCA	(GT) <sub>10</sub>	VIC	238–265	LC030220

<sup>a</sup>Annealing temperature was 57°C for all loci.

<sup>b</sup>Sequence of the fluorescent labels: FAM = 5'-CACGACGTTGTAAAACGAC-3', NED = 5'-CTATAGGGCACGCGTGGT-3', VIC = 5'-TGTG-GAATTGTGAGCGG-3'.

(mean 0.66) and from 0.35 to 0.94 (mean 0.68), respectively (Table 2). Only one locus (Veja009) in the Gunma population showed significant deviation from HWE ( $P < 0.05$ ). Significant LD was found in the Nagano population for only one pair of loci (Veja004 and Veja021;  $P < 0.05$ ).

## CONCLUSIONS

The microsatellite markers described in this study will be useful for conservation genetic studies of *V. japonicum*, such as those evaluating genetic diversity and structure within and between populations. Assessment of their genetic information will also contribute to elucidating how *V. japonicum* genetic diversity and structure has been affected by declining populations

and habitat management practices. Understanding the impacts of habitat management will help the conservation of both *V. japonicum* and the endangered butterfly *M. ambigua*, because maintenance of *V. japonicum* population size is crucial for survival of this butterfly species.

## LITERATURE CITED

- ASSOCIATION OF WILDLIFE RESEARCH AND ENVISION. 2007. Search system of Japanese Red Data. Website <http://www.jpnrdb.com/index.html> [accessed 1 February 2015] [in Japanese].  
BOUTIN-GANACHE, I., M. RAPOSO, M. RAYMOND, AND C. F. DESCHÉPPER. 2001. M13-tailed primers improve the readability and usability of

TABLE 2. Results for primer screening of all samples for 12 microsatellite loci in two populations of *Veronicastrum japonicum*.<sup>a,b</sup>

Locus	Gunma ( $N = 21$ )				Nagano ( $N = 27$ )			
	$A$	$H_o$	$H_e$	$F_{IS}$	$A$	$H_o$	$H_e$	$F_{IS}$
Veja004	7	0.81	0.75	−0.09	5	0.48	0.59	0.19
Veja005	2	0.43	0.39	−0.11	3	0.37	0.60	0.39
Veja007	6	0.62	0.71	0.13	6	0.70	0.65	−0.08
Veja009	8	<b>0.48</b>	0.70	0.32	8	0.70	0.80	0.12
Veja011	6	0.62	0.53	−0.17	5	0.59	0.64	0.08
Veja012	13	0.86	0.82	−0.05	12	0.78	0.77	−0.01
Veja015	4	0.52	0.67	0.22	4	0.30	0.35	0.15
Veja018	18	0.95	0.91	−0.04	24	1.00	0.94	−0.06
Veja021	7	0.62	0.73	0.15	6	0.67	0.71	0.06
Veja025	9	0.57	0.67	0.14	10	0.93	0.85	−0.09
Veja029	5	0.52	0.52	−0.01	3	0.70	0.60	−0.17
Veja032	5	0.86	0.73	−0.17	9	0.63	0.67	0.06

Note:  $A$  = number of alleles;  $F_{IS}$  = fixation index;  $H_e$  = expected heterozygosity;  $H_o$  = observed heterozygosity;  $N$  = sample size.

<sup>a</sup>Populations: Gunma Prefecture: 36°26'12"N, 138°25'27"E; Nagano Prefecture: 35°57'25"N, 137°37'49"E. Vouchers deposited in the Kyoto University Museum herbarium (accession numbers: KYO\_00019997 and KYO\_00019998, respectively).

<sup>b</sup>Numbers in boldface show deviations from Hardy–Weinberg equilibrium ( $P < 0.05$ ).

- microsatellite analyses performed with two different allele-sizing methods. *BioTechniques* 31: 24–26.
- FAIRCLOTH, B. C. 2008. MSATCOMMANDER: Detection of microsatellite repeat arrays and automated, locus-specific primer design. *Molecular Ecology Resources* 8: 92–94.
- GOUDET, J. 1995. FSTAT: A computer program to calculate *F*-statistics, version 1.2. *Journal of Heredity* 86: 485–486.
- MINISTRY OF THE ENVIRONMENT OF JAPAN. 2012. The red list of insects of Japan. Web site [http://www.biodic.go.jp/rdb/rdb\\_f.html](http://www.biodic.go.jp/rdb/rdb_f.html) [accessed 1 February 2015] [in Japanese].
- NAGAIKE, T., E. OHKUBO, AND K. HIROSE. 2014. Vegetation recovery in response to the exclusion of grazing by sika deer (*Cervus nippon*) in seminatural grassland on Mt. Kushigata, Japan. *ISRN Biodiversity* 2014: 493495.
- OHWI, J. 1978. Flora of Japan. Shibundo, Tokyo, Japan [in Japanese].
- PEAKALL, R., AND P. E. SMOUSE. 2006. GenAlEx 6: Genetic analysis in Excel. Population genetic software for teaching and research. *Molecular Ecology Notes* 6: 288–295.
- ROZEN, S., AND H. SKALETISKY. 1999. Primer3 on the WWW for general users and for biologist programmers. In S. Misener and S. A. Krawetz [eds.], *Methods in molecular biology*, vol. 132: Bioinformatics methods and protocols, 365–386. Humana Press, Totowa, New Jersey, USA.
- TAKATSUKI, S. 2009. Effects of sika deer on vegetation in Japan: A review. *Biological Conservation* 142: 1922–1929.