

Updated Distribution of the Pink Fairy Armadillo *Chlamyphorus truncatus* (Xenarthra, Dasypodidae), the World's Smallest Armadillo

Authors: Borghi, Carlos E., Campos, Claudia M., Giannoni, Stella M., Campos, Valeria E., and Sillero-Zubiri, Claudio

Source: Edentata, 12(1) : 14-19

Published By: IUCN/SSC Anteater, Sloth and Armadillo Specialist Group

URL: <https://doi.org/10.5537/020.012.0103>

BioOne Complete (complete.BioOne.org) is a full-text database of 200 subscribed and open-access titles in the biological, ecological, and environmental sciences published by nonprofit societies, associations, museums, institutions, and presses.

Your use of this PDF, the BioOne Complete website, and all posted and associated content indicates your acceptance of BioOne's Terms of Use, available at www.bioone.org/terms-of-use.

Usage of BioOne Complete content is strictly limited to personal, educational, and non - commercial use. Commercial inquiries or rights and permissions requests should be directed to the individual publisher as copyright holder.

BioOne sees sustainable scholarly publishing as an inherently collaborative enterprise connecting authors, nonprofit publishers, academic institutions, research libraries, and research funders in the common goal of maximizing access to critical research.

Updated distribution of the pink fairy armadillo *Chlamyphorus truncatus* (Xenarthra, Dasypodidae), the world's smallest armadillo

CARLOS E. BORGHÍ^{A,1}, CLAUDIA M. CAMPOS^B, STELLA M. GIANNONI^A,
VALERIA E. CAMPOS^A & CLAUDIO SILLERO-ZUBIRÍ^C

^A INTERBIODES (Research Group on Biological Interactions in the Desert), Department of Biology and Institute and Museum of Natural Sciences & CONICET (National Council for Scientific and Technical Research, Argentina), FCEFN, National University of San Juan, Av. Ignacio de la Roza y Meglioli, San Juan J5402DCS, Rivadavia; San Juan, Argentina. E-mail: cborghi@unsj-cuim.edu.ar

^B GIB – IADIZA – CCT-MENDOZA CONICET, Av. Ruiz Leal s/n, Parque General San Martín, Mendoza, Argentina. CP 5500.
E-mail: ccampos@mendoza-conicet.gob.ar

^C Wildlife Conservation Research Unit, Zoology Department, University of Oxford. The Recanati-Kaplan Centre, Tubney House, Tubney OX13 5QL, United Kingdom. E-mail: claudio.sillero@zoo.ox.ac.uk

¹ Corresponding author

Abstract The distribution and ecology of the pink fairy armadillo *Chlamyphorus truncatus*, the world's smallest armadillo, are poorly understood due largely to the strictly subterranean habits of this species. We compiled records from the literature, museum and private collections, information from rural people, and researchers to update the species' current distribution and conservation threats. Recent records suggest that populations of *C. truncatus* persist in localities along most of the species' historical range, including at least 12 sites with recurrent pink fairy armadillo records. We added new registers outside the historical distribution; south of Mendoza province, north of Río Negro province and south of Buenos Aires province, effectively extending the species range. The rate at which new records are accrued has increased, arguably reflecting a greater interest in the species and greater presence of field workers in the species range. The conservation of the pink fairy armadillo depends on the maintenance of the arid and semiarid habitats where this species occurs, with commercial agriculture, pesticides, extensive livestock farming, and predation by domestic dogs and cats, posing the threats of greatest concern.

Keywords: Argentina, *Chlamyphorus truncatus*, conservation, dryland, pichiciego, rare species, subterranean mammal

Distribución actualizada del pichiciego menor *Chlamyphorus truncatus* (Xenarthra, Dasypodidae), el armadillo más pequeño del mundo

Resumen La distribución y ecología del pichiciego menor, *Chlamyphorus truncatus*, el armadillo más pequeño del mundo, han sido poco estudiadas debido a los hábitos estrictamente subterráneos de la especie. Recogimos registros de la literatura, de museos y colecciones privadas e información provista por pobladores rurales e investigadores a los fines de actualizar la distribución actual de la especie y sus problemas de conservación. Los registros recientes sugieren que las poblaciones de *C. truncatus* persisten en localidades a lo largo del rango de distribución histórica, incluyendo al menos 12 sitios con registros recurrentes de la especie. Agregamos nuevos registros fuera de su distribución histórica, al sur de la provincia de Mendoza, al norte de Río Negro y al sur de la provincia de Buenos Aires, extendiendo así el rango de distribución. La tasa de acumulación de nuevos registros ha incrementado en los últimos tiempos, lo cual refleja un interés creciente por la especie y un aumento de trabajos de campo en su rango de distribución. La conservación del pichiciego menor depende del mantenimiento de los hábitats áridos y semiáridos que ocupa. Las amenazas más preocupantes para la especie son la agricultura y ganadería extensiva, el uso de pesticidas y la depredación por animales domésticos.

Palabras clave: Argentina, *Chlamyphorus truncatus*, conservación, especie rara, mamífero subterráneo, pichiciego, zonas áridas

INTRODUCTION

The pink fairy armadillo *Chlamyphorus truncatus* was described by Harlan (1835), who used its native name “pichiciago”, mentioned by Mr. William Colesberry when presenting a specimen to the Philadelphia Museum. The pink fairy armadillo is a desert-adapted, strictly subterranean armadillo (Ceí, 1967; Borghi *et al.*, 2002) endemic to central Argentina, where it inhabits sandy plains, dunes, and scrubby grasslands dominated by sparse shrubs of creosotebush (*Larrea* spp.) and by *Portulaca* plants in the wet season (spring and summer).

This species is the smallest living armadillo (Dasypodidae), and is amongst the least known subterranean mammals. It has minute eyes; its fine silky fur is yellowish white and its flexible dorsal shell of about 24 bands is attached to the body only by a thin dorsal membrane. A vertical plate at the rear of the carapace gives it a rather blunt end (to which the name *truncatus* refers), from which a spatula-shaped tail protrudes (Minoprio, 1945; Guiñazú Rawson de Arentsen, 1956).

Yepes (1928, 1929, 1931, 1932), Minoprio (1945), Guiñazú Rawson de Arentsen (1956), Ceí (1967), and Meritt (1985) have reported on the natural history and geographical distribution of pink fairy armadillos. They have nocturnal and solitary habits and a diet largely composed of insects (mainly ants and beetles), worms, snails, and small amounts of roots and other plant parts (Minoprio, 1945; Guiñazú Rawson de Arentsen, 1956; Ceí, 1967).

The conservation status of the pink fairy armadillo is uncertain. It is listed as Data Deficient by the IUCN Red List of Threatened Species (Superina *et al.*, 2009) but a decline in distribution and abundance is generally accepted (Lagiglia, 1956; Roig, 1995; Ojeda & Diaz, 1997; Diaz & Ojeda, 2000; Superina, 2006; Aguiar & Fonseca, 2008; Superina *et al.*, 2009). This perceived decline has been associated with farming activities, particularly large-scale plowing, as well as with predation by domestic dogs and cats (Minoprio, 1945; Lagiglia, 1956; Fonseca & Aguiar, 2004; Aguiar & Fonseca, 2008; Superina *et al.*, 2009; Abba & Superina, 2010).

Here we review the historical and current distribution of the pink fairy armadillo, and address the following questions: 1. Are there sites with recurrent records of the species, indicating long-term persistence?, and 2. Do the current location data show a reduction of its historical range?

MATERIALS AND METHODS

We obtained data on distribution and collection date from the literature, museum and private collections, carcasses, and live animals observed or captured by rural people and field researchers.

Specimen data were collated from the following collections: Museo Argentino de Ciencias Naturales “Bernardino Rivadavia”, Buenos Aires, Argentina (MACN); Instituto Argentino de Investigaciones de las Zonas Áridas, CCT CONICET Mendoza, Argentina (CMI); and Museo Municipal de Historia Natural, San Rafael, Mendoza, Argentina (MMHN).

In addition, informal, unstructured, and open-ended interviews were conducted throughout the species range, targeting key informants, including park rangers, field biologists, farm hands (locally referred as “puesteros”), and rural school students and teachers. Interviewees were asked for details of any observations of pink fairy armadillos, including location, habitat, date and natural history, and were confronted with photos of the species to confirm the correct identification (Rabinowitz, 2003). Whenever there was more than one record in a single location, independent of the time scale involved, we considered it to be a recurrent record. Records were grouped into four categories (date of last record for recurrent records): those without a precise date, prior to 1980 (i.e., historical distribution), between 1981 and 1993 (i.e., recent records), and from 1994 to present (i.e., current records). Only the records assigned to a precise location were plotted on the map. The historical distribution of the species is represented using the map in Yepes (1929).

RESULTS

A total of 96 records were collected, of which 91 included date of collection or observation. Of these, 69 records had detailed locality information (**TABLE 1**). Most records (58%) corresponded to museum collections, 26% to key informants, 14% were collated from the literature and from observed or captured live animals, and 1% from a field record of a carapace. Records came mainly from the provinces of Mendoza (54%), San Luis (10%), Buenos Aires (10%), La Pampa (9%), and San Juan (7%) (**TABLE 1**).

We added new registers to the historical distribution of the species, in the south of Mendoza, north of Río Negro and the south of Buenos Aires (**FIG. 1**). We found 12 sites with recurrent pink fairy armadillo records (**FIG. 1 AND TABLE 1**): seven in Mendoza, three in Buenos Aires, one in La Pampa, and one in San Juan. Of the 91 records precisely dated, 45 were classified as historical (1896–1980), 14 as recent (1981–1993), and 32 as current (1994 to present). The rate at which these records were accrued was 0.5 records/year prior to 1980, 1.2 records/year between 1981 and 1993, and 1.9 records/year since 1994.

Several sites with recurrent records were registered across the range, but were more abundant in Mendoza and Buenos Aires provinces. Recurrent records from 1994 to date were found in Telteca, Divisadero, Ñacuñán, Monte Comán, El Nihuil,

Canalejas (Mendoza), Carro Quemado (La Pampa), Monte Hermoso, Sauce Grande and Chasicó (Buenos Aires), and Valle Fértil (San Juan) (**TABLE 1**). Eight of these registers correspond to sites with recent (post-2000) records. **TABLE 1** indicates the differences between the first and the last register for all recurrent records.

DISCUSSION

Populations of the pink fairy armadillo persist throughout most of its historical range, as indicated by the records we collated (**FIG. 1**). The distribution of reports and recurrent records presented here reflects a close relationship between frequency of registers and presence of field researchers specifically interested in the species (V. Roig, C. E. Borghi, and M. Superina in Mendoza, C. E. Borghi in San Juan). Due to their nocturnal and subterranean habits, these animals are extremely difficult to observe; consequently, there is a paucity of population data, and the conservation status of extant populations remains unknown. Nevertheless, the rate at which records are collated has increased over the last century, with the current accrual rate being the highest. This may be due to various reasons, such as the existence of more roads/trails, greater vehicle access, or more people working in rural environments (e.g., rangers, NGO personnel, rural school teachers). Thus, even though the populations of the species might be declining, greater field activities in the species range




FIGURE 1. Updated distribution of *Chlamyphorus truncatus*. Records were grouped into three categories: ● prior to 1980; ■ between 1981 and 1993; and ▲ from 1994 to date. Grey symbols indicate only one record and black symbols indicate recurrent records in the same location. Numbers correspond to location reference numbers (**SEE TABLE 1**). Dashed line: Distribution of *C. truncatus* redrawn from Yepes (1929), indicated for comparative purposes.

TABLE 1. Locations of *Chlamyphorus truncatus* examined in this study, including province, locality, type of record and number of reference or source, years between recurrent records, location coordinates, and reference numbers (see Figure 1). Type of records: collection (C), reports from the literature (R), living or dead individuals found in the field or recorded during interviews during this study (L), mention in newspapers with a photograph (N).

| Reference number | Province | Locality | Type of record (number of individuals–year) | Reference or source | Maximum interval between recurrent records (years) | Coordinate Locations |
|------------------|--------------|--------------------|---|--|--|----------------------|
| 1 | Buenos Aires | Bahía Unión | L (1–1995) | This study | --- | 39°33'S; 62°05'W |
| 2 | Buenos Aires | Laguna Chasicó | C (1–1929), L (1–2008) | Collection MACN & A. Abba pers. comm. | 79 | 38°35'S; 63°09'W |
| 3 | Buenos Aires | Monte Hermoso | L (1–1960s, 1–1970s, 1–1990) | This study | 30 | 38°58'S; 61°17'W |
| 4 | Buenos Aires | Sauce Grande | C (1–1978), L (1–1995, 1–n.d.) | 1 in the Private Collection of C. Sillero & this study | 17 | 38°59'S; 61°10'W |
| 5 | Buenos Aires | La Colorada Chica | R (1–2009), | Morici (2011) | --- | 38°59'S; 61°10'W |
| 6 | Catamarca | La Guardia | C (1–1918) | Collection MACN | --- | 29°34'S; 65°26'W |
| 7 | Córdoba | La Nacional | C (1–1994) | Collection MACN | --- | 34°46'S; 64°51'W |
| 8 | Córdoba | Vicuña Mackenna | L (1–1980s) | This study | --- | 33°54'S; 64°23'W |
| 9 | La Pampa | Carro Quemado | C (1–1971); L (2–2008) | Collection MACN & A. Abba pers. comm. | 37 | 36°30'S; 65°20'W |
| 10 | La Pampa | Parque Luro | R (1–1997) | Massoia & Tiranti (1997) | --- | 36°55'S; 64°14'W |
| 11 | La Pampa | Paso del Algarrobo | L (1–1993) | C. Sillero, this study | --- | 36°42'S; 66°54'W |

TABLE 1 CONTINUED ON NEXT PAGE

TABLE 1, CONTINUED FROM PREVIOUS PAGE

| Reference number | Province | Locality | Type of record (number of individuals–year) | Reference or source | Maximum interval between recurrent records (years) | Coordinate Locations |
|------------------|-----------|--|--|---|--|----------------------|
| 12 | La Pampa | Santa Isabel | C (1–1980) | Collection MMHN | --- | 36°32'S; 66°54'W |
| 13 | La Pampa | Colonia Butaló | C (1–1977) | Collection MMHN | --- | 36°12'S; 67°04'W |
| 14 | La Pampa | Victorica | C (1–1929) | Collection MACN | --- | 36°13'S; 65°26'W |
| 15 | La Rioja | Patquia | C (1–1931) | Collection MACN | --- | 30°03'S; 66°53'W |
| 16 | Mendoza | Pichi Ciego | L (1–1996) | This study | --- | 33°23'S; 68°04'W |
| 17 | Mendoza | Alto del Algarrobal | C (1–1971) | Collection MMHN | --- | 34°37'S; 68°10'W |
| 18 | Mendoza | Carmensa | L (1–2000) | This study | --- | 35°08'S; 67°39'W |
| 19 | Mendoza | San Carlos | C (1–1978) | Collection MMHN | --- | 33°46'S; 69°02'W |
| 20 | Mendoza | Colonia Tabanera | C (1–1976) | Collection MMHN | --- | 34°31'S; 68°13'W |
| 21 | Mendoza | Corral de Lorca | R (1–2000s) | Superina (2006) | --- | 34°41'S; 67°04'W |
| 22 | Mendoza | El Divisadero | C (2–1989), L (1–2001) | Collection CMI & this study | 12 | 33°45'S; 67°40'W |
| 23 | Mendoza | El Nihuil | C(1–1976), R (1–2000s) | Collection MMHN & Superina (2006) | 24 | 35°03'S; 68°38'W |
| 24 | Mendoza | El Sosneado | C (1–1971) | Collection MMHN | --- | 35°04'S; 69°34'W |
| 25 | Mendoza | Santa Rosa | L (1–1999) | This study | --- | 33°16'S; 68°05'W |
| 26 | Mendoza | La Paz | C (1–1906) | Collection MACN | --- | 33°27'S; 67°32'W |
| 27 | Mendoza | Las Catitas | L (1–2002) | C. Borghi, this study | --- | 33°18'S; 68°02'W |
| 28 | Mendoza | Las Malvinas | C (1–1977) | Collection MMHN | --- | 34°50'S; 68°14'W |
| 29 | Mendoza | Las Paredes, El Usillal | C (1–1990) | Collection MMHN | --- | 34°29'S; 68°25'W |
| 30 | Mendoza | Monte Comán | R (3–2004, 2005, 2006) | Superina (2006) | 2 | 34°35'S; 67°51'W |
| 31 | Mendoza | Ñacuñán | C (2–1980s, 1–1988), R (1–2000s, 1–2006), L (1–2000) | Collection (CMI), Superina (2006) & this study (L) | 20 | 34°02'S; 67°56'W |
| 32 | Mendoza | Puesto Cortez | L (1–2009) | This study | --- | 35°36'S; 69°02'W |
| 33 | Mendoza | Reserva Telteca | C (1–1993), L (1–1997) | Collection IADIZA & this study | 4 | 32°21'S; 67°59'W |
| 34 | Mendoza | San Miguel | C (2–1987) | Collection IADIZA | 0 | 32°21'S; 67°24'W |
| 35 | Mendoza | San Pedro del Atuel | C (1–1980) | Collection MMHN | --- | 35°09'S; 67°38'W |
| 36 | Mendoza | Canalejas | L (1–2009), N (1–2011) | This study & Newspaper: "Diario San Rafael, 29/07/2011" | 2 | 35°12'S; 66°32'W |
| 37 | Río Negro | Villa Regina | C (1–1957) | Collection MACN | --- | 39°04'S; 67°04'W |
| 38 | San Juan | Agua Cercada | L (1–2008) | This study | --- | 30°46'S; 67°22'W |
| 39 | San Juan | Cochagual | C (1–1915) | Collection MACN | --- | 31°55'S; 68°23'W |
| 40 | San Juan | Encón | L (1–2000s) | This study | --- | 32°10'S; 67°43'W |
| 41 | San Juan | San Agustín de Valle Fertil | L (1–2008), L & N (1–2010) | This study & Newspaper: "Diario de Cuyo, 07/03/2010" | 2 | 30°38'S; 67°27'W |
| 42 | San Juan | Zonda | R (1–1920) | Yepes (1928) | --- | 31°28'S; 68°45'W |
| 43 | San Luis | Altos Negros | L (1–1996) | This study | --- | 34°08'S; 66°06'W |
| 44 | San Luis | Buena Esperanza | R (1–1967) | Rood (1970) | --- | 34°45'S; 65°14'W |
| 45 | San Luis | Chischaca | C (1–1926) | Collection MACN | --- | 33°52'S; 66°14'W |
| 46 | San Luis | Nueva Escocia to Batavia (20 km south Nueva Escocia) | L (1–1996) | This study | --- | 32°22'S; 68°02'W |
| 47 | San Luis | Pedernera, Bagual | C (1–1914) | Collection MACN | --- | 33°47'S; 65°17'W |
| 48 | San Luis | Pisco Yacú | C (1–1931) | Collection MACN | --- | 32°20'S; 65°12'W |

might explain the higher number of pink fairy armadillo encounters observed in recent times. Recent recurrent records of the species (post-2000) suggest that, at least in some localities across the historical range, populations of *C. truncatus* still persist.

There are no recent records on the southern, northern, and eastern fringes of the species range, largely due to the paucity of field research undertaken in those areas since 1980. Thus, no assumptions can be made about the persistence of the species there. In contrast, we show new registers in areas outside the historical distribution described by Yepes (1929) (FIG. 1).

Ultimately, the conservation of the pink fairy armadillo would depend on the persistence of arid and semiarid habitats where the species occurs. The main threats to pink fairy armadillos are related to degradation and loss of habitats due to land clearing for commercial agriculture and overgrazing (Morello, 1985; Ojeda *et al.*, 2002; Tabeni & Ojeda, 2005; Vilela *et al.*, 2009), with the rapid expansion of agriculture in the western and northern parts of the species' range being of greatest concern (Abba & Superina, 2010). Current development policy in this region includes a rapid increase in land devoted to vineyards, olive groves, and other intensive crops suited to sandy soils (Vilela *et al.*, 2009). Additionally, predation by feral and farm dogs and cats has been suggested as a major direct threat to pink fairy armadillos (Minoprio, 1945; Lagiglia, 1956), but its real impact remains unknown. The presence of wild boar (*Sus scrofa*), an exotic species introduced to Argentina in the last century (Novillo & Ojeda, 2008; Cuevas *et al.*, 2010), might also have a negative impact on pink fairy armadillos due to their disruptive foraging habits. Although there are recurrent records of pink fairy armadillos in Ñacuñán, Telteca, and other protected areas, it is uncertain whether there is a fully protected viable population in these reserves. Unfortunately, even in these protected areas the pink fairy armadillo is vulnerable to predation by the ever present dogs and cats.

Field studies are needed to better understand the ecology of this rare, little-known species, and the effects on its populations of agricultural activities, extensive livestock farming, interactions with wild boar, and predation by domestic predators. Also, as feeders of soil fauna and plant matter, it has been suggested that armadillos may be exposed to agricultural toxins and pesticides (Herbst *et al.*, 1989). Although the effects of these compounds on pink fairy armadillos have not been evaluated, organophosphate insecticides are used in vineyards and olive crops to control leaf-cutting ants (M. Battistella, pers. comm.). Since pink fairy armadillos chiefly eat ants (Redford, 1985), the use of agricultural pesticides potentially poses one of the greatest threats to their persistence.

ACKNOWLEDGEMENTS

We thank the Mendoza Provincial Park Rangers Mario Villegas and Julián Baronetto (Telteca and La Payunia Reserves), Néstor Maceira and Enrique Fontemachi (INTA, Instituto Nacional de Tecnología Agropecuaria, Argentina), Ricardo Ojeda and Paola Sassi (CONICET), Verónica Allayme (Village of Ñacuñán, Mendoza), Agustín M. Abba (University of La Plata), Leif Larssen (Monte Hermoso), Lisandro Crego (La Pampa), and Gustavo Aprile (Fundación Vida Silvestre Argentina) for field records of pink fairy armadillos. Maximiliano Battistella (INTA San Juan, Argentina) advised us on pest control of ants in vineyards and olive crops. Erika Cuellar (Oxford University, UK) kindly commented on an earlier manuscript, and Nelly Horak assisted with an earlier English version of our manuscript. We also like to thank two anonymous reviewers for their valuable suggestions. The work of C. E. Borghi was supported in part by the Brookfield Zoo Chicago Zoological Society. C.E.B., C.M.C., S.M.G., and V.E.C. are supported by CONICET.

REFERENCES

- Abba, A. M. & M. Superina. 2010. The 2009/2010 armadillo Red List assessment. *Edentata* 11: 135–184.
- Aguiar, J. M. & G. A. B. Fonseca. 2008. Conservation status of the *Xenarthra*. Pp. 215–231 in: *The Biology of the Xenarthra* (S. F. Vizcaíno & W. J. Loughry, eds.). University Press of Florida, Gainesville.
- Borghi, C. E., S. M. Giannoni & V. G. Roig. 2002. Eye reduction in subterranean mammals and eye protective behavior in *Ctenomys*. *Mastozoología Neotropical* 9: 123–134.
- Cei, J. M. 1967. Pichiciego and portulaca. *Animals* (London) 10: 176–177.
- Cuevas, M. F., A. Novillo, C. Campos, M. A. Dacar & R. A. Ojeda. 2010. Food habits and impact of rooting behaviour of the invasive wild boar, *Sus scrofa*, in a protected area of the Monte Desert, Argentina. *Journal of Arid Environments* 74: 1582–1585.
- Diaz, G. & R. Ojeda. 2000. *Libro Rojo de los mamíferos amenazados de la Argentina*. Sociedad Argentina para el Estudio de los Mamíferos, Buenos Aires, Argentina. 221 pp.
- Fonseca, G. A. B. & J. M. Aguiar. 2004. The 2004 Edentate Species Assessment workshop. *Edentata* 6: 1–26.
- Guinazú Rawson de Arentsen, B. V. 1956. El pichi hormiguero: contribución a su conocimiento y estudio. *Revista Científica de Investigación*

- del Museo de Historia Natural de San Rafael 1: 25–32.
- Harlan, R. 1835. Description of *Chlamyphorus truncatus*, a new genus of mammiferous quadrupeds, of the Order Edentata. Pp 31–46 in: Medical and physical researches: or original memoirs in medicine, surgery, physiology, geology, zoology, and comparative anatomy. Lydia R. Bailey, Philadelphia.
- Herbst, L. H., A. I. Webb, R. M. Clemmons, M. R. Dorsey-Lee & E. E. Storrs. 1989. Plasma and erythrocyte cholinesterase values for the common long-nosed armadillo, *Dasypus novemcinctus*. Journal of Wildlife Diseases 25: 364–369.
- Lagiglia, H. A. 1956. Nota sobre la retrocesión y conservación del *Chlamyphorus truncatus* Harl. dentro de la fauna mendocina. Revista Científica de Investigación del Museo de Historia Natural de San Rafael 1: 33–34.
- Massoia, E. & I. Tiranti. 1997. Nuevos datos sobre los Dasypodidae vivientes y fósiles argentinos (Mammalia, Cingulata) – 3– *Chlamyphorus truncatus* de la Provincia de La Pampa. APRONA 31: 28–31.
- Meritt, D. A. Jr. 1985. The fairy armadillo, *Chlamyphorus truncatus* Harlan. Pp. 393–395 in: The evolution and ecology of armadillos, sloths, and vermilinguas (G. G. Montgomery, ed.). Smithsonian Institution Press, Washington and London.
- Minoprio, J. D. L. 1945. Sobre el *Chlamyphorus truncatus*, Harlan. Acta Zoologica Lilloana 3: 5–58.
- Morello, J. A. 1985. Grandes ecosistemas de Sudamérica. Textos para discusión. Fundación Bariloche, Argentina. 116pp.
- Morici, A. 2011. Nuevo registro de pichiciego menor (*Chlamyphorus truncatus*) para la provincia de Buenos Aires, Argentina. Biológica: Naturaleza, Conservación y Sociedad 14: 77–78.
- Novillo, A. & R. A. Ojeda. 2008. The exotic mammals of Argentina. Biological Invasions 10: 1333–1344.
- Ojeda, R. A. & G. B. Diaz. 1997. Sección II. Mamíferos. Pp. 73–219 in: Libro Rojo de los mamíferos y aves amenazados de la Argentina (F. J. García, R. A. Ojeda, R. M. Fraga, G. B. Diaz & R. J. Baigún, eds.). Administración de Parques Nacionales, Buenos Aires, Argentina.
- Ojeda, R. A., C. E. Borghi & V. G. Roig. 2002. Mamíferos de Argentina. Pp. 23–63 in: Diversidad y conservación de los mamíferos neotropicales (G. Ceballos & J. A. Simonetti, eds.), CONABIO–UNAM, México, D.F.
- Rabinowitz, A. 2003. Manual de capacitación para la investigación de campo y la conservación de la vida silvestre. Wildlife Conservation Society, FAN, Bolivia. 310pp.
- Redford, K. H. 1985. Food habits of armadillos (Xenarthra: Dasypodidae). Pp. 429–437 in: The evolution and ecology of armadillos, sloths, and vermilinguas (G. G. Montgomery, ed.). Smithsonian Institution Press, Washington and London.
- Roig, V. G. 1995. Situación de conservación, biología y ecología de *Chlamyphorus truncatus*. Edentata 2: 19.
- Rood, J. P. 1970. Notes on the behavior of the pygmy armadillo. Journal of Mammalogy 51: 179.
- Superina, M. 2006. New information on population declines in pink fairy armadillos. Edentata 7: 48–50.
- Superina, M., A. M. Abba & V. Roig. 2009. *Chlamyphorus truncatus*. In: IUCN 2011. The IUCN Red List of Threatened Species 2011.1. International Union for Conservation of Nature and Natural Resources. <<http://www.iucnredlist.org>>. Downloaded on 31 October 2011.
- Tabeni, S. & R. A. Ojeda. 2005. Ecology of the Monte Desert small mammals in disturbed and undisturbed habitats. Journal of Arid Environments 63: 244–255.
- Vilela, A., M. L. Bolkovic, P. Carmanchahi, M. Cony, D. Delamo & D. Wassner. 2009. Past, present and potential uses of native flora and wildlife of the Monte Desert. Journal of Arid Environments 73: 238–243.
- Yepes, J. 1928. Los Edentata argentinos. Revista de la Universidad de Buenos Aires 24: 461–515.
- Yepes, J. 1929. Notas sobre la distribución geográfica del “pichi ciego menor” (*Chlamyphorus truncatus*) y “pichi llorón” (*Chaetophractus vellerosus*). Physis 9: 439–446.
- Yepes, J. 1931. El escudete cefálico del pichi ciego menor (*Chlamyphorus truncatus* Harl.). Revista Chilena de Historia Natural 35: 107–112.
- Yepes, J. 1932. Las formas geográficas del “pichi ciego menor” (*Chlamyphorus truncatus* Harl.). Physis 11: 9–18.

Received: 15 August 2011; Accepted: 24 November 2011